



GENERALIZED RENORMALIZABLE GAUGE FORMULATION
OF SPONTANEOUSLY BROKEN GAUGE THEORIES

KAZUO FUJIKAWA[†]
The Enrico Fermi Institute
The University of Chicago, Chicago, Illinois 60637

BENJAMIN W. LEE^{*} and A. I. SANDA
National Accelerator Laboratory, Batavia, Illinois 60510

[†] Work supported in part by NSF Contract No. NSF GP 32904X

^{*} On leave of absence from Institute for Theoretical Physics, SUNY
Stony Brook, New York 11790



ABSTRACT

The spontaneously broken gauge theory is formulated in the generalized renormalizable gauge (R_ξ -gauge). A parameter ξ can be adjusted to include existing gauges, U-gauge, R-gauge, and 't Hooft-Feynman gauge as special cases. Three applications of the R_ξ -gauge formulation are given. First we compute the weak correction to the muon magnetic moment unambiguously in the existing models for leptons. Secondly, we discuss the large momentum transfer limit of the Pauli-magnetic form factor of the muon. Finally, we discuss the static charge of the neutrino, and show that an appropriate regularization makes it vanish.

I. INTRODUCTION

The possibility of constructing a unified theory of weak and electromagnetic interactions in terms of a spontaneously broken gauge symmetry has attracted a great deal of attention lately, following the works of Weinberg¹ and 't Hooft.² In this paper we shall present a formulation of spontaneously broken gauge theories (SBGT) which is particularly suited for practical calculations. In this formulation the gauge condition one adopts is a generalization of the one used by 't Hooft and depends on a parameter ξ which can vary continuously from 0 to ∞ . In this gauge, which we shall call generically as the R_ξ -gauge, the massive vector boson propagator is precisely the one invented by Lee and Yang in their discussion³ of the ξ -limiting process

$$\begin{aligned} \Delta_{\mu\nu}(p, \xi) &= -i \left[g_{\mu\nu} - \left(1 - \frac{1}{\xi}\right) \frac{p_\mu p_\nu}{p^2 - \frac{1}{\xi} M^2} \right] \frac{1}{p^2 - M^2} \\ &= -i \left(g_{\mu\nu} - \frac{1}{M^2} p_\mu p_\nu \right) \frac{1}{p^2 - M^2} - i \frac{1}{M^2} p_\mu p_\nu \frac{1}{p^2 - \frac{1}{\xi} M^2} \quad (1.1) \end{aligned}$$

The difference between the R_ξ -gauge formulation of SBGT and the ξ -limiting process applied to the electrodynamics of massive vector bosons is this: in the former, the negative metric scalar boson pole of the vector boson propagator at $p^2 = M^2/\xi$ is cancelled by the pole of the unphysical scalar boson propagator

$$\frac{i}{p^2 - \frac{1}{\xi} M^2} \quad (1.2)$$

in the S-matrix, and the S-matrix of the former is independent of the parameter ξ and is unitary, whereas in the latter, one recovers the unitarity of the S-matrix only in the limit $\xi \rightarrow 0$. The ξ -independence of the S-matrix⁴ in the former is a direct consequence of the nonabelian gauge invariance of the relevant Lagrangian.

It is worthwhile to note the connection between the R_ξ -gauge and other gauges discussed in the literature.

(1) The R-Gauge: In the proof of renormalizability of SGGT by Lee and Zinn-Justin⁵, and also in the discussion of Salam and Strathdee⁶, a generalization of the Landau gauge in quantum electrodynamics, the so-called R-gauge, was used. The R-gauge is obtained from the R_ξ -gauge for $\xi = \infty$.

(2) The 't Hooft-Feynmann Gauge: This gauge, which was discussed by 't Hooft, is obtained when we set $\xi = 1$. In this gauge the vector boson propagator is proportional to $g_{\mu\nu}$, and the unphysical scalar boson propagator of Eq. (1.2) has a pole at $p^2 = M^2$.

(3) The U-Gauge: In this formulation, the unphysical scalar bosons are absent and the vector boson propagator is the canonical one.

$$\Delta_{\mu\nu}(p) = -i \left(g_{\mu\nu} - \frac{p_\mu p_\nu}{M^2} \right) \frac{1}{p^2 - M^2} \quad (1.3)$$

In this gauge, the unitarity of the S-matrix is manifest since there are no spurious singularities at $p^2 = M^2/\xi$. However, Green's functions are unrenormalizable in this gauge: it is only the S-matrix that can be defined in this gauge. The U-gauge is formally equivalent to the R_ξ -gauge in the limit $\xi \rightarrow 0$. The equivalence here is "formal", in the sense that Feynman amplitudes in the two formulations are equal if the limit $\xi \rightarrow 0$ is taken before the Feynman integral is performed.

The U-gauge formulation of SGBT deserves some more discussion. Because the quantization of SGBT in this gauge is most straightforward, most of the existing calculations were performed in this formulation, despite the divergence difficulties unique to this gauge. The cancellation of divergences in the S-matrix (but not in Green's functions) has been demonstrated by various authors in a number of cases⁷. However, the isolation of the finite part of an S-matrix element in this gauge may prove ambiguous. In fact, Jackiw and Weinberg and Bars and Yoshimura⁸ have commented on an ambiguity that exists in the calculation of the weak-interaction contribution to the anomalous magnetic moment of the muon. We claim that, based on our own experiences, the computation of Feynman amplitudes is enormously simplified in the R_ξ -gauge. It is also easier to check the ξ -independence of the S-matrix (thereby verifying the unitarity of the S-matrix) in the R_ξ -gauge, than to establish the cancellation of higher order divergences. When there are ambiguities in fixing the finite part of an S-matrix element in the U-gauge, the R_ξ -gauge formulation provides a gauge invariant (with respect to the nonabelian gauge group) way of circumventing such difficulties. In fact, we shall resolve the ambiguity in the computation of the anomalous

magnetic moment of the muon by evaluating it in the R_ξ -gauge. Our study explains also why the ξ -limiting process used by Jackiw and Weinberg, and Bars and Yoshimura yields the correct result.⁹

This paper is organized as follows: In Section II we formulate the generalized renormalizable gauge (R_ξ -gauge). In Section III, we apply the R_ξ -gauge to the calculation of weak correction to the magnetic moment of the muon. We will present unambiguous answers for three existing models of Weinberg,⁷ of Georgi and Glashow¹⁰ and of B. Lee, and Prentki and Zumino.¹¹ In Section IV, we show that the naive calculation of the neutrino static charge gives a non-vanishing result and we discuss how to remedy this situation. In Appendix A, we give details of Section III. In Appendix B we point out the reasons for ambiguities present in the U-gauge calculations of the weak correction to the muon magnetic moment. Finally, in Appendix C, we give the Lagrangians and necessary Feynman rules for our calculations.

After the completion of this paper, we received a paper by Y. P. Yao in which a formulation similar to ours is discussed in the context of an abelian gauge theory.

II. FORMULATION OF THE R_ξ -GAUGE.

In this Section we shall discuss the formulation of SGBT in a general class of covariant linear gauge conditions. We shall consider, for definiteness, the Georgi-Glashow model based on the $O(3)$ gauge symmetry without fermions. In Appendix, we will extend our considerations of this Section to all three models mentioned in the Introduction, with fermions.

In the absence of fermions, the Georgi-Glashow model consists of a triplet of gauge bosons and a triplet of scalar mesons. The Lagrangian is of the form

$$\begin{aligned} \mathcal{L} = & -\frac{1}{4} \left(\partial_\mu \vec{B}_\nu - \partial_\nu \vec{B}_\mu + g \vec{B}_\mu \times \vec{B}_\nu \right)^2 \\ & + \frac{1}{2} \left[(\partial_\mu + g \vec{B}_\mu \times) \vec{\phi} \right]^2 - V(\vec{\phi}) \end{aligned} \quad (2.1)$$

where $V(q)$ is an isospin invariant quartic polynomial of the scalar fields $\vec{\phi}$. The potential is assumed to have an absolute minimum at $\vec{\phi} = \vec{v} \neq 0$. We can always choose the isospin z-axis to coincide with the direction of \vec{v} .

It is convenient to define a unit vector $\hat{\eta}$ along the z-axis:

$$\vec{v} = v \hat{\eta}.$$

We also define $\vec{\phi}_t$ and ϕ by

$$\begin{aligned} \vec{\phi} &= \vec{\phi}_t + \hat{\eta}(v + \phi), \\ \vec{\phi}_t \cdot \hat{\eta} &= 0. \end{aligned} \quad (2.2)$$

The gauge condition we shall adopt is (see also Appendix B)

$$\partial^\mu \vec{B}_\mu - \frac{1}{\xi} g v \hat{\eta} \times \vec{\phi} = 0 \quad (2.3)$$

where ξ is a non-negative real parameter.

It was shown by 't Hooft¹² that the gauge condition (2.3) may be taken care of by defining the effective action¹³:

$$\begin{aligned} S[\vec{B}_\mu, \vec{\phi}] \\ = \int d^4x \left[\mathcal{L}(x) - \frac{\xi}{2} \left(\partial^\mu \vec{B}_\mu - \frac{1}{\xi} g \vec{v} \times \vec{\phi} \right)^2 - \frac{1}{2} \left(\frac{1}{\alpha} - \xi \right) \left(\partial^\mu B_\mu^3 \right)^2 \right] \\ - i \text{Tr} \ln [1 + g \mathcal{G} \gamma] \end{aligned} \quad (2.4)$$

Where \mathcal{G} is defined by

$$\begin{aligned} [(-\partial^2 + i\epsilon) \delta_{ab} - \frac{1}{\xi} (g v)^2 (\delta_{ab} - \eta_a \eta_b)] \langle b, x | \mathcal{G} | c, y \rangle \\ = \delta_{ac} \delta^4(x-y) \end{aligned} \quad (2.5)$$

and γ is defined by

$$\begin{aligned} \langle a, x | \gamma | b, y \rangle \\ = \left\{ \epsilon_{abc} \frac{\partial}{\partial x_\mu} B_\mu^c(x) + \frac{1}{\xi} g v [\phi_{ta}(x) \eta_b - \phi (\delta_{ab} - \eta_a \eta_b)] \right\} \delta^4(x-y) \end{aligned} \quad (2.6)$$

and Tr denotes the trace operation over the space-time variables x, y , as well as over the isospin indices a, b . In Eq. (2.4), ξ and α are in general arbitrary non-negative real numbers.

The effective action S of Eq. (2.4) is to be used in defining the generating functional of connected Green's functions. Thus, if we define

$$\begin{aligned} \exp i Z[\vec{J}_\mu, \vec{K}] \\ = \int [d\vec{B}_\mu] [d\vec{\phi}] \exp \left\{ i S[\vec{B}_\mu, \vec{\phi}] \right. \\ \left. + i \int d^4x [\vec{K}(x) \cdot \vec{\phi}(x) - \vec{J}_\mu(x) \cdot \vec{B}^\mu(x)] \right\}, \end{aligned} \quad (2.7)$$

functional derivatives of Z at $\vec{J}_\mu = \vec{K} = 0$ give connected Green's functions of the theory. The generating functional Z depends on two parameters α' and ξ . Note that the choice $\alpha'^{-1} = \xi = \infty$ leads to the R-gauge discussed in references 5 and 6.

The effective Lagrangian can be written as

$$\begin{aligned}
 \mathcal{L} &= \frac{\xi}{2} \left(\partial^\mu \vec{B}_\mu - \frac{1}{\xi} \partial_\mu \vec{v} \times \vec{\phi} \right)^2 - \frac{1}{2} \left(\frac{1}{\alpha} - \xi \right) \left(\partial^\mu B_\mu^3 \right)^2 \\
 &= \frac{1}{2} \left[\partial_\mu \vec{\phi} + g \vec{B} \times (\vec{\phi}_t \times \phi \hat{\eta}) \right]^2 - \frac{(g v)^2}{2 \xi} \vec{\phi}_t^2 \\
 &\quad + g^2 v \hat{\eta} \times \vec{B} \cdot \vec{B} \times (\vec{\phi}_t + \phi \hat{\eta}) - V(\vec{\phi}) \\
 &\quad - \frac{1}{4} (\partial_\mu \vec{B}_\nu - \partial_\nu \vec{B}_\mu + g \vec{B}_\mu \times \vec{B}_\nu)^2 + \frac{(g v)^2}{2} (\vec{B}_\mu^t)^2 \\
 &\quad - \frac{\xi}{2} (\partial^\mu \vec{B}_\mu^t)^2 - \frac{1}{2\alpha} (\partial^\mu B_\mu^3)^2.
 \end{aligned} \tag{2.8}$$

The terms proportional to $\vec{B}_\mu \cdot \partial^\mu \vec{\phi}$ have disappeared from Eq. (2.8).

The propagators for various fields are obtained by inverting the matrix of the quadratic form L_0 of the above expression:

$$\begin{aligned}
 \mathcal{L}_0 &= \frac{1}{2} \left[(\partial_\mu \vec{\phi})^2 - \frac{M^2}{\xi} \vec{\phi}_t^2 - \mu_\phi^2 \phi^2 \right] \\
 &\quad - \frac{1}{4} (\partial_\mu \vec{B}_\nu - \partial_\nu \vec{B}_\mu)^2 - \frac{\xi}{2} (\partial^\mu \vec{B}_\mu^t)^2 + \frac{M^2}{2} (\vec{B}_\mu^t)^2 - \frac{1}{2\alpha} (\partial^\mu B_\mu^3)^2
 \end{aligned}$$

where

$$M^2 = g v, \quad \mu_\phi^2 = 2 \partial V(v \hat{\eta}) / \partial v^2 \tag{2.9}$$

and they are ¹⁴

$$S^\pm \equiv \frac{1}{\sqrt{2}} (\phi^1 \pm i\phi^2) \quad : \quad i \frac{1}{k^2 - \frac{1}{\xi} M^2 + i\epsilon}$$

$$\psi \equiv \phi^3 - \sigma \quad : \quad i \frac{1}{k^2 - \mu_\psi^2 + i\epsilon}$$

$$W_\mu \equiv \frac{1}{\sqrt{2}} (B_\mu^1 + i B_\mu^2) \quad : \quad -i \left[g_{\mu\nu} - k_\mu k_\nu \frac{1}{k^2 - \frac{1}{\xi} M^2} \left(1 - \frac{1}{\xi}\right) \right] \frac{1}{k^2 - M^2 + i\epsilon}$$

(2.10)

and

$$A_\mu \equiv B_\mu^3 \quad : \quad -i \left(g_{\mu\nu} - \frac{k_\mu k_\nu}{k^2} (1-\alpha) \right) \frac{1}{k^2 + i\epsilon}.$$

We see that our gauge interpolates between the R-gauge ($\xi \rightarrow \infty$) and the U-gauge ($\xi \rightarrow 0$). For $\xi = 1$, we recover the 't Hooft-Feynman gauge, in which the vector boson propagators are proportional to $g_{\mu\nu}$.

In the Weinberg model (and also in the model of B. Lee, and Prentki and Zumino) we have another gauge boson Z_μ . We fix the gauge for this boson by adding the following term to the Lagrangian

$$\mathcal{L}^{c'} = - \frac{\eta}{2} \left(\partial_\mu Z^\mu + \frac{G\nu}{\eta} \chi \right)^2, \quad G = \sqrt{g^2 + g'^2} \quad (2.11)$$

where η is a parameter that can vary over the range

$$0 \leq \eta \quad (2.12)$$

and G and χ are the coupling constant of Z_μ and the corresponding unphysical neutral scalar boson, respectively. We must also modify the last term $-i \text{Tr} \ln [1 + g \mathcal{O} \gamma]$ in Eq. (2.4) accordingly. The propagator for Z_μ and χ are given by

$$Z_\mu : -i \frac{\left(\delta_{\mu\nu} - \frac{k_\mu k_\nu (1-\eta)}{M_Z^2 - \eta k^2} \right)}{k^2 - M_Z^2 + i\epsilon} \quad (2.13)$$

$$\chi : i \frac{1}{k^2 - \frac{1}{\eta} M_Z^2 + i\epsilon} \quad (2.14)$$

where $M_Z \equiv Gv$.

In an S-matrix element, the pole at $k^2 = \frac{1}{\xi} M^2$ of the vector boson propagator is cancelled by the similar pole at m^2/ξ of the s^\pm -propagator. Neither scalar particles implied by these poles are physical. (In the R-gauge formulation the s^\pm are the would-be Goldstone fields). In fact the couplings of s^\pm to other particles can be determined based on the above considerations. As an example, let us determine the coupling of s^- to the $e \nu$ pair. We write the coupling of $W_\mu^- \left[\equiv \frac{1}{\sqrt{2}} (B_\mu^1 + i B_\mu^2) \right]$ to the $e \nu$ pair as

$$\mathcal{L}_{We\nu} = g \bar{e} (1 - \gamma_5) \nu W_\mu^-.$$

Now consider the T-matrix element for the process

$$e(p) + \nu(q) \rightarrow \nu(q') + e(p').$$

To lowest order, the W-exchange gives

$$(ig)^2 (-i) \left[\bar{u}_e(p') \gamma^\mu (1 - \gamma_5) u_\nu(q) \right] \left[\bar{u}_\nu(q') \gamma^\lambda (1 - \gamma_5) u_e(p) \right] \\ \times \left[\left(g^{\mu\lambda} - \frac{k^\mu k^\lambda}{M^2} \right) \frac{1}{k^2 - M^2} + \frac{k^\mu k^\lambda}{M^2} \frac{1}{k^2 - \frac{1}{\xi} M^2} \right]$$

where $k = p' - q = p - q'$, and we have used the vector boson propagator of Eq.

(2.10). The pole term at $k^2 = \frac{1}{\xi} M^2$:

$$\left(\frac{i g m_e}{M} \right)^2 (-i) \left[\bar{u}_e(p') (1 - \gamma_5) u_\nu(q) \right] \left[\bar{u}_\nu(q') (1 + \gamma_5) u_e(p) \right] \frac{1}{k^2 - \frac{1}{\xi} M^2},$$

$$M^2 = (g v)^2$$

must be cancelled by the s^- exchange contribution. This requires the $s^- e \nu$ coupling to be

$$\mathcal{L}_{s^- e \nu} = \pm \frac{m_e}{v} \bar{e} (1 - \gamma_5) \nu s^-$$

(The sign ambiguity is superficial, since the sign of v is indeterminate .

Once a definite sign convention is made here, all other couplings are uniquely determined).

We note that the above cancellation is one of the consequences of the following two fundamental relations:

(i) The S-matrix is gauge independent, namely

$$\frac{\partial S}{\partial \xi} \equiv 0$$

(ii) The propagator $D_{\mu\nu}$ for W_μ and the propagator D for s^\pm satisfy the identity

$$\frac{\partial}{\partial \xi} D_{\mu\nu}(k) \equiv - \frac{k_\mu k_\nu}{M^2} \frac{\partial}{\partial \xi} D(k)$$

III. MAGNETIC FORM FACTOR OF THE MUON

a. Weak Correction to the Magnetic Moment of the Muon.

In a unified theory of weak and electromagnetic interactions, one-loop contributions to the anomalous magnetic moment a_μ of the muon are formally of order α , whether they derive from photon exchange or weak vector boson exchanges. It can be written to this order as

$$a_\mu = \frac{\alpha}{2\pi} \left\{ 1 + \left(\frac{\mu}{M} \right)^2 f \left(\left(\frac{\mu}{M} \right), \left(\frac{m_{Y^0}}{\mu} \right) \right) \right\} \quad (3.1)$$

where μ = muon mass, M = W boson mass, f is a function of the mass ratios (μ/M) and (m_{Y^0}/μ) , and m_{Y^0} is the mass of a neutral heavy lepton Y^0 that might exist in such a theory. The second term in Eq. (3.1) is in magnitude of order $(\alpha/M^2) \mu^2 \sim G_F \mu^2$ and we shall call it the weak correction to a_μ and denote it by

$$a_\mu^w = \frac{\alpha}{2\pi} \left(\frac{\mu}{M} \right)^2 f \left(\left(\frac{\mu}{M} \right), \left(\frac{m_{Y^0}}{\mu} \right) \right) \quad (3.2)$$

In addition, there are contributions of massive Higgs scalar bosons in such a theory to a_μ . However, they are of order $(\mu/m_\phi)^2$ compared to Eq. (3.2), where m_ϕ is a typical Higgs scalar mass, and since the masses of these scalars are presumably very large, we shall ignore them in the following discussion.

The weak correction to a_μ^w has been computed by several authors⁸ in the U-gauge previously. In this gauge, the electromagnetic vertex of the muon is quadratically divergent, so that its separation into the electric and anomalous (Pauli) magnetic form factors is ambiguous. As a consequence, one finds that a_μ^w computed in this gauge depends on the way the internal momentum is routed in a diagram, even though it is finite.

In the R_ξ -gauge, the electromagnetic vertex of the muon is only logarithmically divergent, and there is no such ambiguity in evaluating a_μ^W . In order to verify the gauge invariance we have evaluated it in three different gauges: $\xi = \infty, 1$ and 0 . In this section we will present the results of these calculations. In Appendix A, we will present a general proof that the value of a_μ^W is independent of ξ . In the following we shall refer to the result obtained in the limit $\xi \rightarrow 0$ after the Feynman integration as the U-gauge result. For those diagrams involving unphysical scalars in this particular case, the limit $\xi \rightarrow 0$ and the integration commute. This explains why the procedure used by Jackiw and Weinberg⁸, and Bars and Yoshimura⁸, of replacing the vector propagator (1.3) by the regularized one (1.1) yielded the correct result, even though this replacement per se is not a gauge invariant procedure.

Our results are given for three different models. These are the model of Weinberg⁹ based on $SU(2) \times U(1)$, that of Georgi and Glashow¹⁰ based on $O(3)$ and that of Lee and Prentki and Zumino (LPZ) based on $O(3) \times O(2)$. The diagrams shown in Figs. 1 and 2 contribute to the models given by Weinberg and LPZ, and the diagrams shown in Figs. 1 and 3 contribute to the model of Georgi and Glashow. In these figures s^- and χ are unphysical scalars and Y^0 is a neutral lepton. For the purpose of illustration let us evaluate the diagram shown in Fig. 2b. It gives the contribution

$$-e \frac{\mu^2}{M_Z^2} \frac{(\xi^2 + \xi'^2)}{4} \int \frac{d^4 k}{(2\pi)^4} \frac{\bar{u}(p') \gamma'_5 (\not{p}' - \not{k} + \mu) \not{\epsilon}_\mu (\not{p} - \not{k} + \mu) \gamma'_5 u(p)}{[(p-k)^2 - \mu^2][(p'-k)^2 - \mu^2][k^2 - M_Z^2/\xi]} \quad (3.3)$$

where p and p' are the incident and final muon momenta respectively and k is the internal momentum of χ . We separate this expression into the charge and anomalous magnetic form factors:

$$-ie \bar{u}(p') \left[F_1(q^2) \gamma_\mu + F_2(q^2) \frac{i}{2\mu} \sigma_{\mu\nu} q^\nu + \text{parity violating terms} \right] u(p),$$

$$q = p' - p. \quad (3.4)$$

After the k integration $F_2(q^2)$ of Eq. (3.4) is found to be

$$F_2(q^2) = - \left(\frac{g \mu^2}{4M^2} \right) \frac{\mu^2}{8\pi^2} \int_0^1 dx \int_0^{1-x} dy \frac{(x+y)^2}{\mu^2(x+y)^2 - g^2 xy + M_Z^2/\eta(1-x-y)}. \quad (3.5)$$

Integrations over x and y yield the desired result in the gauge characterized by η . For example, to obtain the R-gauge result, we take the limit $\eta \rightarrow \infty$. Then

$$a_\mu^w(F_{12} b)_R = - \frac{G_F \mu^2}{8\pi^2 \sqrt{2}} \quad (3.6)$$

To get from Eq. (3.3) to Eq. (3.6) we have used the relations

$$g^2 + g'^2 = g^2 \frac{M_Z^2}{M^2} \quad \text{and} \quad \frac{g^2}{8M^2} = G_F/\sqrt{2}$$

To obtain U-gauge result, we let $\eta \rightarrow 0$ and we find that the diagram does not contribute.

$$a_\mu^w(F_{12} b)_U = 0 \quad (3.7)$$

Finally in a t'Hooft-Feynman gauge, we let $\eta = 1$ and we see that

$$a_\mu^w(F_{12} b)_{tH-F} = O(\mu^2/M_Z^2) \quad (3.8)$$

Details of all other diagrams can be found in Appendix A. Tables 1, 2 and 3 give contributions from Figs 1, 2 and 3 respectively. For example, contribution of Fig. 2b can be found in Table 2 column b. We neglected terms of order μ/M . In Table 3, terms of order m_{Y0}/M , μ/m_{Y0} were also neglected. It is amusing to see that individual diagrams are quite gauge dependent, but, as it must, the diagrams always add up to give a gauge independent result not only to the leading order in $(\mu/M)^2$, $(\mu/m_Y)^2$ but to all orders. (See Appendix A).

To obtain the result of Weinberg's model, we add the results of Tables 1 and 2

$$a_\mu^W = \frac{G_F \mu^2}{8\pi^2 \sqrt{2}} \left\{ \frac{10}{3} + \frac{1}{3} \left[(3 - 4 \cos^2 \theta)^2 - 5 \right] \right\} \quad (3.9)$$

Note that this result is the same as those obtained previously in the U-gauge by the ϵ -limiting regularization.⁸

The result for the model proposed by LPZ can be obtained from that of Weinberg's model by merely changing the definition for the coupling constants. We obtain

$$a_\mu^W = \frac{G_F \mu^2}{8\pi^2 \sqrt{2}} \left\{ \frac{10}{3} - \frac{4}{3} \frac{u^2 + v^2}{u^2} (1 + \sin^2 \theta \cos^2 \theta) \right\} \quad (3.10)$$

where θ is related to the physical quantities by

$$\frac{G_F}{\sqrt{2}} = \frac{e^2}{4M^2 \sin^2 \theta} \quad \text{and} \quad \frac{u^2 + v^2}{u^2} = 2 \frac{M_1^2}{M_Z^2} (1 + \tan^2 \theta) \quad (3.11)$$

u and v are the vacuum expectation values of Higgs' scalars in this model.

In Eq. (3.10), a_μ^W can be about the same order of magnitude as that of Eq. (3.9) if $(M^2/M_Z^2) \tan^2 \theta \sim O(1)$. In that case both of these models predict the weak correction to the muon magnetic moment to be of order 10^{-8} , below the experimental detectability. In any case, the strong interaction correction a_μ^S has been estimated from the colliding beam experiment to be of order $(6.5 \pm 0.5) \times 10^{-8}$,¹⁵ so that the latter seems to be bigger than the former.

The Georgi-Glashow model receives contributions from Figs. 1 and 2.

The result of the calculation in the special case $m_{Y^0} \ll M$ are given in Table 3 and Table 1. Note that Fig. 3 gives contributions which are much larger than the previous two models by the factor m_{Y^+}/μ . We will thus concentrate our attention on the contributions of order $G_F \mu m_{Y^0}$ in Fig. 3. Evaluating for arbitrary value of m_{Y^0}/M we obtain

$$a_\mu^W = - \frac{G_F (m_{Y^+} + \mu) \mu}{8\sqrt{2} \pi^2 \sin^2 \theta} \left[1 + \frac{b}{(1-y)^3} \left(\frac{1}{2} - 2y + \frac{3}{2} y^2 - y^2 \ln y \right) \right] \quad (3.12)$$

where $y = m_{Y^0}^2/M^2$ and $\sin^2 \theta = 4G_F M^2 / \sqrt{2} e^2$.

Note also the relation

$$2m_{Y^0} \cos \theta = m_{Y^+} + \mu.$$

In Fig. 4 we have plotted the prediction of Eq. (3.12). Two sets of curves correspond to contours of constant a_μ^W and m_{Y^0} at various values of m_{Y^+} and M .

The present status on experimental and theoretical uncertainties in a_μ places the limit¹⁵

$$2 \times 10^{-8} \leq a_\mu^W + a_\mu^S \leq 6 \times 10^{-7} \quad (3.13)$$

where a_μ^S corresponds to the hadronic correction to the muon magnetic moment.

Using the value for a_μ^S quoted above, it seems quite safe to guess that

$$-2 \times 10^{-7} \leq a_\mu^W \leq 0.$$

If we further demand, for example $m_{Y^+} \geq 0.5 \text{ GeV}$ Fig. 4 readily

gives the allowed range of m_{Y^0} and M . The generous lower limit

$-2 \times 10^{-7} = a_\mu^W$ gives $m_{Y^+} \leq 5 \text{ GeV}$. If we use $|a_\mu^W| \leq +1.1 \times 10^{-7}$, we get $m_{Y^+} \leq 2 \text{ GeV}$. In these estimates we assume $m_{Y^+} m_\mu / M_\Psi^2 \ll 1$.

A charged heavy lepton of this mass range can be detected in the near future.

A pair of Y^+ and Y^- can be produced in reactions such as

$$\gamma + (Z) \rightarrow Y^+ Y^- + (Z)$$

or

$$e^+ + e^- \rightarrow Y^+ + Y^-$$

The detection of coincident e_μ^- from the decays

$$Y^- \rightarrow \bar{\nu}_\mu + \bar{\nu}_e + e^-$$

$$Y^+ \rightarrow \bar{\nu}_\mu + \bar{\nu}_\tau + \mu^+$$

is a signature of the $Y^+ Y^-$ pair production.

It is important to recognize that a_μ^W in Eq. (3.12) does not vanish even in the limit $m_{Y^0} \rightarrow \infty$. If one performs a naive U-gauge calculation the first term in Eq. (3.12) is absent and $a_\mu^W \rightarrow 0$ for $m_{Y^0} \rightarrow \infty$. (i.e. we can make a_μ^W arbitrarily small by letting m_{Y^0} be large).

b. Asymptotic Behavior of $F_2(q^2)$.

In this sub-section we discuss the behavior of the Pauli magnetic form factor $F_2(q^2)$ as $|q^2| \rightarrow \infty$. We caution the reader that $F_2(q^2)$ for $q^2 \neq 0$ is not an on-shell S-matrix element (i.e., not measurable) and is not invariant under nonabelian gauge group (i.e., depends on the gauge). Fig. 5a gives a process in which $F_2(q^2)$ is relevant. But Fig. 5b is of the same order. We obtain the gauge invariant answer only when Fig. 5a and 5b and all other diagrams of the same order are added. $F_2(q^2=0) = a_\mu^W$ is available for experimental measurements only because of the pole due to the photon propagator. Still, the knowledge of the asymptotic behavior of $F_2(q^2)$, though gauge dependent, is important in the question of renormalizability when diagrams of the type Fig. 5a is inserted in more complicated diagrams.

Our conclusion is that $F_2(q^2) \rightarrow 0$ as $|q^2| \rightarrow \infty$ in all gauges except the U-gauge i.e. for all combinations $\eta \neq 0, \xi \neq 0$. This can be easily seen at least for off mass-shell muons, as follows: for $\eta \neq 0, \xi \neq 0$, the triangle diagram that we consider has the degree of divergence at most zero. Thus due to the kinematical factor $\sigma_{\mu\nu} q_\nu$, the integral for $F_2(q^2)$ has the degree of divergence -1 . Therefore, by Weinberg's theorem¹⁶, $F_2(q^2) \leq O(1/\sqrt{q^2})$.

We have also done the calculation for the on-mass-shell ^{muon} amplitude and verified that $F(q^2) \rightarrow 0$ as $|q^2| \rightarrow \infty$ in all gauges except the U-gauge in the Weinberg model. In order to obtain the result for the U-gauge, we let $\xi \rightarrow 0$ and then let $q^2 \rightarrow -\infty$. The result is (for the Weinberg model)

$$F_2(q^2) \rightarrow \frac{G_F \mu^2}{\sqrt{2} 8\pi^2} \left(4 - \frac{\mu^2}{M^2}\right) \ln(-q^2) + \text{constant for } q^2 \rightarrow -\infty.$$

These results indicate the gauge dependence of the off-shell amplitude. In particular, for the renormalizable gauge (i.e. for $\xi \neq 0$), $F_2(q^2)$ shows a manifestly renormalizable behavior. On the other hand, $F(q^2)$ in the U-gauge exhibits a divergent behavior at $q^2 = -\infty$. As remarked in reference 17, the logarithmic growth of $F_2(q^2)$ for large q^2 does not necessarily imply any trouble with S-matrix elements for physical processes. When all diagrams of the same order for a physical process are added the bad behavior of $F_2(q^2)$ can be cancelled by those of other diagrams.

The dispersion relation for F_2 in the U-gauge requires a subtraction (which cannot be determined a priori) while its absorptive part may be computed by the standard Landau-Cutkosky rule. On the other hand, F_2 in the R_ξ -gauge has an absorptive contribution from unphysical states, while it requires no unknown subtraction.

[After the completion of this paper, we received a preprint of W. A. Bardeen et al, CERN 1485, in which they evaluate a_μ in the Weinberg model using the n regularization method of 't Hooft and Veltman (G. 't Hooft and M. T. Veltman, Nuclear Physics, to be published). Their answer agrees with ours. H. R. Quinn and J. Primack have computed a_μ for the Georgi-Glashow model. We appreciate Professor Quinn's explaining their result to us].

IV. STATIC CHARGE OF THE NEUTRINO.

As in the case of Pauli-magnetic form factor discussed in Section III, the notion of the electric form factor of a (muon -) neutrino is an unphysical one in the present theory: the electric form factor, for nonzero momentum transfer, is not an element of the S-matrix and, in the present theory, is not gauge invariant nor unitary for arbitrary ξ . However, the electric charge, i.e., the value of the electric form factor F_1 at zero momentum transfer, is an element of the S-matrix and measurable. It must be zero if due care is exercised in evaluating Feynman integrals.

In the Georgi-Glashow model, there are altogether 10 diagrams contributing to the electric charge of the neutrino. In Fig. 6, we show 5 of them which involve internal muon lines. The other five are similar and involve internal γ^+ lines. We shall evaluate the Feynman integrals in the R-gauge ($\xi = \infty$) for convenience. (We have also checked the ξ independence of our results). The contribution of each of the five diagrams in Fig. 6 is:

$$a: \quad F_1^{(a)}(0) = -i \frac{3}{2} \left(\frac{g}{2} \right)^2 \int \frac{d^4 k}{(2\pi)^4} \frac{1}{(k^2 - M^2)^2} \\ - i \frac{3}{2} \left(\frac{g}{2} \right)^2 \int \frac{d^4 k}{(2\pi)^4} \frac{\mu^2}{k^2 - \mu^2} \frac{1}{(k^2 - M^2)^2} \quad (4.1)$$

$$b: \quad F_1^{(b)}(0) = i \frac{3}{2} \left(\frac{g}{2} \right)^2 \int \frac{d^4 k}{(2\pi)^4} \frac{1}{k^2} \frac{\mu^2}{k^2 - \mu^2} \frac{1}{k^2 - M^2}$$

$$c: \quad F_1^{(c)}(0) = -i \frac{1}{2} \left(\frac{g}{2} \right)^2 \int \frac{d^4 k}{(2\pi)^4} \frac{1}{k^2} \frac{1}{k^2 - \mu^2}$$

$$d: \quad F_1^{(d)}(0) = -i \left(\frac{3}{2} \right) \left(\frac{g}{2} \right)^2 \int \frac{d^4 k}{(2\pi)^4} \frac{\mu^2}{(k^2 - M^2) (k^2 - \mu^2)^2}$$

$$e: \quad F_1^{(e)}(0) = -i \frac{1}{2} \left(\frac{f_\mu}{2} \right)^2 \int \frac{d^4 k}{(2\pi)^4} \frac{k^2 - 2\mu^2}{k^2 (k^2 - \mu^2)^2}$$

where μ and M are the masses of the muon and the W boson, and f_μ is the coupling constant of the scalar meson to the muon and neutrino:

$$f_\mu = g\mu/M.$$

In Eq. (4.1) we have written $F_1^{(a)}(0)$ as a sum of logarithmically divergent and convergent integrals. A simple computation shows that the sum of the second term of $F_1^{(a)}$, $F_1^{(b)}$ and $F_1^{(d)}$ are zero, after the integration so that if the sum $F_1^{(c)} + F_1^{(e)}$ vanishes, then the muon contribution to the electric charge of the neutrino is

$$- \frac{3}{2} \left(\frac{g}{2} \right)^2 \int \frac{d^4 k}{(2\pi)^4} \frac{1}{(k^2 - M^2)^2}$$

and is independent of the muon mass. The γ^+ contribution to the electric charge of the neutrino is exactly opposite to the above mentioned μ^- -contribution, so the net charge of the neutrino is zero as it should be.

Thus, the matter hinges entirely on whether the sum $F_1^{(c)} + F_1^{(e)}$ is zero. A naive evaluation of these two terms gives

$$\begin{aligned} F_1^{(c)} + F_1^{(e)} &= \frac{f_\mu^2}{8} I \\ I &= i \int \frac{d^4 k}{(2\pi)^4} \left[\frac{k^2 - 2\mu^2}{k^2 (k^2 - \mu^2)^2} - \frac{1}{k^2 (k^2 - \mu^2)} \right] \\ &= - \frac{i}{8\pi^2}, \end{aligned} \quad (4.2)$$

which is not zero; it is significant that the value of I is independent of the muon mass μ . Note that $F_1^{(c)} + F_1^{(e)}$ cannot be cancelled by the similar contribution of γ^+ , since the latter is proportional to $(f_{Y^+})^2$, f_{Y^+} being the coupling constant of the scalar meson to the muon-neutrino pair. That is, a naive evaluation of Feynman integrals leads to a nonzero electric charge of the neutrino!

The above paradox has nothing to do with the nonabelian gauge invariance of the theory or the massless nature of the neutrino. The offending diagrams, Figs. 5c and 5e, are characteristic of a theory in which fermions are coupled to a scalar meson. The sum of the two diagrams shown in Fig. 7 may be written as

$$F_\mu \sim - \frac{\partial}{\partial q^r} \int \frac{d^4 k}{(2\pi)^4} \gamma_5 \left[\frac{1}{(\not{Y} + \not{K} - m_P)(k^2 - m_\pi^2)} - \frac{1}{(\not{K} - m_P)[(k-q)^2 - m_\pi^2]} \right] \gamma_5 \quad (4.3)$$

and if we can shift the contour of integration $k \rightarrow k+q$ in the second term of the integrand, the integral vanishes identically. The integral is, however, linearly divergent, so that the change of the variable of integration is legitimate only after the integral is suitably regularized in a gauge invariant manner. A simple regularization scheme is to replace the pion propagators in (4.3) by

$$\begin{aligned} \frac{1}{k^2 - m_\pi^2} &\rightarrow \frac{1}{k^2 - m_\pi^2} - \frac{1}{k^2 - \Lambda_c^2} \\ \frac{1}{(k-q)^2 - m_\pi^2} &\rightarrow \frac{1}{(k-q)^2 - m_\pi^2} - \frac{1}{(k-q)^2 - \Lambda_o^2} \end{aligned} \quad (4.4)$$

In this case, the naive evaluation of the charge of the neutron gives a result independent of the mass m^2 internal pion lines so that the regularization implied by Eq. (4.4) yields to a zero neutron charge. The result here is counter to the folklore which says that convergent Feynman integrals need not be regulated: if we perform the differentiation with respect to q prior to integration in Eq. (4.3), as one would to recover the original Feynman integral, then the integral becomes convergent and the conventional wisdom would say that it is not necessary to regulate the integral. What we have learned is that to keep the charge of a neutral fermion zero, it is necessary to regularize Feynman integrals in a gauge invariant way, even if the integrals are convergent.

Let us return now to our problem. We can regularize the scalar meson line in a gauge invariant manner as in the σ' -model: we insert in the Lagrangian the regulator term:

$$-\frac{1}{2}[(D_\mu \vec{\phi}')^2 - \Lambda_0^2 \phi'^2] = -\frac{1}{2} \left\{ [(\partial_\mu + g \vec{B}_\mu) \times \vec{\phi}']^2 - \Lambda_0^2 \phi'^2 \right\}$$

and replace the untranslated scalar fields ϕ by the sum $\phi + \phi'$ in all interaction terms.^{18,19} This modification of the Lagrangian is clearly gauge invariant (with respect to the nonabelian gauge group), and the integral I in Eq. (4.2) is now regulated to read

$$I_{reg} = i \int \frac{d^4 k}{(2\pi)^4} \left[\frac{k^2 - 2\mu^2}{(k^2 - \mu^2)^2} \left(\frac{1}{k^2} - \frac{1}{k^2 - \Lambda_0^2} \right) - \frac{k^2}{k^2 - \mu^2} \left(\frac{1}{(k^2)^2} - \frac{1}{(k^2 - \Lambda_0^2)^2} \right) \right]$$

which is zero for all values of Λ_b^2 . Thus, a gauge invariant regularization of the Feynman integral does give the physically correct result $F_1^{(c)} + F_1^{(e)} = 0$. We remind the reader that $F_1^{(c)} + F_1^{(e)}$ is nonabelian gauge invariant by itself. Thus the regularization procedure stated above is sufficient to remove the neutrino static charge for arbitrary gauge.

V CONCLUSION

We have given a formulation of the convenient gauge for actual applications of SBGT (R_ξ -gauge). Based on this formulation, we have verified the gauge independence of several simple S-matrix elements. This indicates that the ghost killing mechanism is indeed working in examples we have considered. It is important to show that the gauge independence properties of the S-matrix are preserved at every stage of the renormalization program.

In our formulation, the finite part of the S-matrix is uniquely determined. Results of our calculation of weak correction to the muon magnetic moment agree with U-gauge calculations with the ξ -limiting regularization procedure. An experimental implication of our results is that the charged heavy lepton in the Georgi-Glashow model is required to be small (of the order of $.5 \sim 5$ GeV.) It is therefore worthwhile searching for this lepton in the existing accelerators.

A naive calculation of the neutrino static charge in SBGT gives a non-vanishing result. This difficulty, however, also exists in the abelian gauge theory. A prescription to remove the static charge in a manifestly gauge invariant (non-abelian) manner was given. To check the self-consistency of SBGT, it is desirable to evaluate other lower order diagrams and to confirm the absence of any other unexpected "anomalous" behaviors.

APPENDIX A: Formulas for the Muon Magnetic Form Factor and the Proof of the Gauge Independence of the g_μ -Factor.

In this appendix we give the general formulae for the weak corrections to the anomalous magnetic moment of the muon based on our R_ξ -gauge in Section II. We calculate the neutral vector meson contribution in the Weinberg model and the massive neutral lepton contribution in the Georgi-Glashow model. The Feynman rules are given in Appendix C. From these two results one can easily derive the magnetic moment for other schemes of lepton interactions.

A-I, Neutral Vector Meson in the Weinberg Model.

We have two diagrams shown in Figs. 2a and 2b for the neutral vector and scalar meson contribution in the Weinberg model. Fig. 2b has been discussed in Section III. (See Eq. (3.5)).

For Fig. 2a we have

$$-e \bar{u}(p') \left\{ \gamma_\beta \left[a \left(\frac{1+\gamma_5}{2} \right) + b \left(\frac{1-\gamma_5}{2} \right) \right] (\not{\epsilon} + \not{\mu}) \gamma_\mu (\not{\epsilon} + \not{\mu}) \gamma_\alpha \left[a \left(\frac{1+\gamma_5}{2} \right) + b \left(\frac{1-\gamma_5}{2} \right) \right] \right\} u(p) \times \frac{P^{\alpha\beta}(k)}{k^2 - M_Z^2} \quad (A-1)$$

where

$$a = \frac{g'^2}{\sqrt{g^2 + g'^2}}, \quad b = \frac{1}{2} \frac{g'^2 - g^2}{\sqrt{g^2 + g'^2}} \quad (A-2)$$

$$P^{\alpha\beta}(k) = g^{\alpha\beta} - \frac{k^\alpha k^\beta (1-\eta)}{M_Z^2 - \eta k^2}$$

(A-3)

The result is given by

$$F(q^2) = - \frac{\mu^2}{8\pi^2} (a^2 + b^2) \int dx dy \frac{(x+y-2)(x+y-1)}{Q^2 + M_Z^2(1-x-y)}$$

(A-4)

$$- \frac{4\mu^2}{8\pi^2} ab \int dx dy \frac{(x+y-1)}{Q^2 + M_Z^2(1-x-y)}$$

$$+ \frac{\mu^2}{8\pi^2} \frac{(a-b)^2}{2M_Z^2} \left\{ \int dx dy [3(x+y)-2] \ln \left[\frac{Q^2 + \Lambda(1-x-y)}{Q^2 + M_Z^2(1-x-y)} \right] \right.$$

$$\left. - \int dx dy [Q^2(x+y) + 2q^2xy] \left[\frac{1}{Q^2 + M_Z^2(1-x-y)} - \frac{1}{Q^2 + \Lambda(1-x-y)} \right] \right\}$$

where $Q^2 = \mu^2(x+y)^2 - 8^2 xy$, $\Lambda = M_Z^2/\eta$.

Note the relation

$$\frac{(a-b)^2}{M_Z^2} = \frac{1}{4} \frac{g^2}{M^2} \quad (\text{A-5})$$

with M the mass of the W-boson.

Based on Eqs. (3.5), (A-4) and (A-5) we prove the gauge independence condition for the physical S-matrix element, $\frac{\partial}{\partial \eta} F_2(0) \equiv 0$ or equivalently

$$\frac{\partial}{\partial \Lambda} F_2(0) \equiv 0 \quad (\text{A-6})$$

Eq. (A-6) demands the following relation

$$\int_0^1 dt \, t \int_{-1}^1 dz \left\{ (3t-2) \frac{(1-t)}{\mu^2 t^2 + \Lambda(1-t)} - \frac{\mu^2 t^3 (1-t)}{[\mu^2 t^2 + \Lambda(1-t)]^2} + \frac{2\mu^2 t^2 (1-t)}{[\mu^2 t^2 + \Lambda(1-t)]^2} \right\} \equiv 0 \quad (\text{A-7})$$

where we changed the Feynman variables to

$$\begin{aligned} t &= (x+y) \\ z &= (x-y)/t \end{aligned} \quad (\text{A-8})$$

Eq. (A-7) is indeed satisfied if one notices the relation

$$\frac{\partial}{\partial t} \left[\frac{1-t}{\mu^2 t^2 + \Lambda(1-t)} \right] = -\mu^2 \frac{2t-t^2}{[\mu^2 t^2 + \Lambda(1-t)]^2} . \quad (\text{A-9})$$

Therefore the sum of the neutral vector and scalar meson contributions is gauge independent, and consequently it is free of ghost contributions. Eq. (A-6) allows us to use any gauge we want to calculate the matrix element. In particular if one takes the limit $\eta \rightarrow 0$ in Eq. (A-4) one recovers the results based on the ξ -limiting procedure (scalar meson contribution vanishes in this limit).

In passing we note that the Higgs neutral scalar meson (which is independent of the gauge) in the Weinberg model (see Fig. 7) gives the magnetic form factor

$$-\frac{\mu^2}{8\pi^2} \left(\frac{g^2 \mu^2}{4M^2} \right) \int dx dy \frac{(x+y)(x+y-2)}{\mu^2(x+y)^2 - g^2(xy) + M_\psi^2(1-x-y)} \quad (\text{A-10})$$

This gives a small contribution to $F_2(0)$ for $(\mu^2/M_\psi^2) \ll 1$.

A-II. Neutral Massive Lepton in the Georgi-Glashow Model.

The neutral massive lepton in the Georgi-Glashow scheme contains four diagrams. See Figs. 3a - d. Fig. 3d gives

$$\begin{aligned}
& \frac{e^3}{2M^2} \left[(\mu - m \cos \theta)^2 + (\mu \cos \theta - m)^2 \right] \bar{u}(p') \not{k} u(p) \frac{(\ell + \ell')_\mu}{(k^2 - m^2)(\ell^2 - \Lambda)(\ell'^2 - \Lambda)} \\
& + \frac{e^3}{M^2} (\mu - m \cos \theta)(\mu \cos \theta - m) \bar{u}(p') u(p) \frac{(\ell + \ell')_\mu}{(k^2 - m^2)(\ell^2 - \Lambda)(\ell'^2 - \Lambda)} \\
& \hspace{15em} (A-11)
\end{aligned}$$

and the result is

$$\begin{aligned}
F(q^2) = & - \frac{e^2 m \mu}{8 \pi^2 M^2} [(\mu - m \cos \theta)(\mu \cos \theta - m)] \int dx dy \frac{(1-x-y)}{f(\Lambda, \Lambda)} \\
& - \frac{e^2 \mu^2}{16 \pi^2 M^2} [(\mu - m \cos \theta)^2 + (\mu \cos \theta - m)^2] \int dx dy \frac{(x+y)(1-x-y)}{f(\Lambda, \Lambda)} \\
& \hspace{15em} (A-12)
\end{aligned}$$

where we defined

$$\Lambda = M^2 / g$$

$$m = \text{mass of the neutral massive lepton } Y^0 \hspace{10em} (A-13)$$

$$f(a, b) = Q^2 + x(a - \mu^2) + y(b - \mu^2) + (1-x-y)m^2$$

$$Q^2 = \mu^2(x+y)^2 - g^2 xy$$

Note that $F(q^2)$ in Eq. (A-12) vanishes in the U-gauge limit, $\xi \rightarrow 0$ (or $\Lambda \rightarrow \infty$).

Figs. 3b and 3c give rise to

$$\begin{aligned}
 & - \frac{e^3}{2} [\mu(1+\cos^2\theta) - 2m\cos\theta] \left\{ \bar{u}(p') \not{k} \gamma_\alpha u(p) \frac{P_\mu^\alpha(l)}{(k^2-m^2)(l'^2-\Lambda)(l^2-\Lambda)} \right. \\
 & \quad \left. + \bar{u}(p') \gamma_\beta \not{k} u(p) \frac{P_\mu^\beta(l')}{(k^2-m^2)(l^2-\Lambda)(l'^2-\Lambda)} \right\} \\
 & + \frac{e^3}{2} m [m(1+\cos^2\theta) - 2\mu\cos\theta] \left\{ \bar{u}(p') \gamma_\alpha u(p) \frac{P_\mu^\alpha(l)}{(k^2-m^2)(l'^2-\Lambda)(l^2-\Lambda)} \right. \\
 & \quad \left. + \bar{u}(p') \gamma_\beta u(p) \frac{P_\mu^\beta(l')}{(k^2-m^2)(l^2-\Lambda)(l'^2-\Lambda)} \right\} \quad (A-14)
 \end{aligned}$$

where

$$P_\mu^\alpha(l) = g_\mu^\alpha + \frac{k_\mu k^\alpha (\frac{1}{\xi} - 1)}{l^2 - \Lambda}$$

The result is given by

$$\begin{aligned}
 F(q^2) &= \frac{e^2 \mu}{16 \pi^2 M^2} [\mu(1+\cos^2\theta) - 2m\cos\theta] \left\{ 2M^2 \int dx dy \frac{y}{f(\Lambda, M^2)} \right. \\
 & \quad + \int dx dy [3(x+y)-2] \ln \left(\frac{f(\Lambda, \Lambda)}{f(\Lambda, M^2)} \right) \\
 & \quad \left. - \int dx dy (x+y-1) [Q^2 - \mu^2(x+y)] \left[\frac{1}{f(\Lambda, M^2)} - \frac{1}{f(\Lambda, \Lambda)} \right] \right\} \quad (A-15)
 \end{aligned}$$

For the first group in Eq. (A-14). The second group in Eq. (A-14) gives

$$F(q^2) = \frac{1}{16\pi^2} \frac{e^2 m \mu^2}{M^2} \left[m(1 + \omega^2 \theta) - 2\mu \omega \theta \right] \times$$

$$\int dx dy (x+y-1)^2 \left[\frac{1}{f(\lambda, M^2)} - \frac{1}{f(\lambda, \lambda)} \right] \quad (\text{A-16})$$

Note that $F(q^2)$ in Eqs. (A-15) and (A-16) both vanishes in the U-gauge limit $\lambda \rightarrow \infty$.

Finally Figs. 3a gives the expression

$$- e^3 m \omega \theta \bar{u}(p) \gamma_\alpha \gamma_\alpha u(p) \frac{P^{\beta\tau}(\ell') V_{\tau\mu\sigma} P^{\tau\alpha}(\ell)}{(k^2 - m^2)(\ell'^2 - M^2)(\ell^2 - M^2)} \quad (\text{A-17})$$

$$- \frac{e^3}{2} (1 + \omega^2 \theta) \bar{u}(p) \gamma_\beta k_\beta \gamma_\alpha u(p) \frac{P^{\beta\tau}(\ell') V_{\tau\mu\sigma} P^{\tau\alpha}(\ell)}{(k^2 - m^2)(\ell'^2 - M^2)(\ell^2 - M^2)}$$

where

$$V_{\tau\mu\sigma} \equiv g_{\sigma\tau} (\ell + \ell')_\mu - \ell_\tau g_{\mu\sigma} - \ell'_\sigma g_{\mu\tau}$$

$$+ g_{\sigma\mu} \ell_\tau - g_{\tau\mu} \ell_\sigma \quad (\text{A-18})$$

and

$$q = \ell' - \ell$$

The result is rather lengthy. The first group in Eq. (A-17) gives rise to

$$\begin{aligned}
 F(q^2) = & -\frac{1}{8\pi^2} \frac{e^2 \mu m \cos \theta}{M^2} \left\{ 3M^2 \int dx dy \frac{x+y}{f(M^2, M^2)} \right. \\
 & + 3 \int dx dy (x-y) \ln \frac{f(\Lambda, M^2)}{f(M^2, M^2)} \quad (A-19) \\
 & + \int dx dy \left[(1-x+y) [Q^2 - 2\mu^2(x+y) + \mu^2] + 2q^2 y (x-1) \right] \left[\frac{1}{f(M^2, M^2)} - \frac{1}{f(\Lambda, M^2)} \right] \\
 & - \frac{1}{2} \frac{q^2}{M^2} \int dx dy [3(x+y)-2] \ln \left(\frac{f(\Lambda, M^2) f(M^2, \Lambda)}{f(M^2, M^2) f(\Lambda, \Lambda)} \right) \\
 & + \frac{1}{2} \frac{q^2}{M^2} \int dx dy (x+y-1) [Q^2 - 2\mu^2(x+y) + \mu^2] \times \\
 & \left. \times \left[\frac{1}{f(M^2, M^2)} - \frac{1}{f(M^2, \Lambda)} - \frac{1}{f(\Lambda, M^2)} + \frac{1}{f(\Lambda, \Lambda)} \right] \right\}
 \end{aligned}$$

and the second group in Eq. (A-17) gives

$$\begin{aligned}
F(q^2) = & \frac{i}{8\pi^2} \frac{e^2 \mu^2}{4\pi^2} (1 + \cos^2 \theta) \left\{ 2M^2 \int dx dy \frac{(x+y)[2(x+y)+1]}{f(M^2, M^2)} \right. \\
& - 2 \int dx dy [4(x+y)^2 - 9(x+y) + 2 + 6y] \ln \left(\frac{f(\lambda, M^2)}{f(M^2, M^2)} \right) \\
& + 2 \int dx dy \left[\mu^2(x+y-1)^2 [2y + (x+y)(x+y-1)] - g^2 xy [2y + (1-x-y)(2-x-y)] \right] \times \\
& \times \left[\frac{1}{f(M^2, M^2)} - \frac{1}{f(\lambda, M^2)} \right] \\
& + 4 \int dx dy \left\{ g^2(xy-y) \left[\frac{1}{f(M^2, M^2)} - \frac{1}{f(\lambda, M^2)} \right] \right. \\
& + \left(\frac{g^2}{M^2} \right) \int dx dy [4(x+y)^2 - 9(x+y) + 4] \ln \left(\frac{f(\lambda, M^2) f(M^2, \lambda)}{f(M^2, M^2) f(\lambda, \lambda)} \right) \\
& - \left(\frac{g^2}{M^2} \right) \int dx dy (x+y-1) [(x+y-2)Q^2 + \mu^2(x+y)] \times \\
& \times \left[\frac{1}{f(M^2, M^2)} - \frac{1}{f(M^2, \lambda)} - \frac{1}{f(\lambda, M^2)} + \frac{1}{f(\lambda, \lambda)} \right] \left. \right\} \quad (A-20)
\end{aligned}$$

We would like to discuss the gauge independence condition $\frac{\partial}{\partial \xi} F(o) \equiv 0$

$$\text{or } \frac{\partial}{\partial \lambda} F(o) \equiv 0 \quad (A-21)$$

From Eqs. (A-12), (A-15), (A-16), (A-19) and (A-20) we readily recognize that $F(q^2)$ consists of two groups; one of which is proportional to $m \cos \theta$ and the other proportional to $(1 + \cos^2 \theta)$. These two groups separately satisfy Equation (A-21). The proof of Eq. (A-21) can be made as in Eqs. (A-7) - (A-9) by the repeated use of partial integration. We note that the contributions with $f(\Lambda, M^2)$ factor and the contribution with $f(\Lambda, \Lambda)$ factor in Eqs. (A-12)-(A-20) separately satisfy Equation (A-12). We do not write down this lengthy but straightforward proof.

Eq. (A-21) ensures the absence of the ghost contribution to the anomalous magnetic moment. Eq. (A-21) also allows us to use the most convenient gauge when we calculate numerical values. We also note that in the limit $\xi \rightarrow 0$ we recover the result based on the ξ -limiting process.

The neutrino contribution is obtained from the above result by setting $m=0$. The coupling constant should be adjusted according to the specific model one uses.

A-III. Large q^2 Behavior of $F(q^2)$

From the above general results for $F(q^2)$ it is easy to see that all the contributions to $F(q^2)$ from scalar mesons vanish in the U-gauge limit, $\xi = 0$ and $\eta = 0$. They also vanish at $q^2 = -\infty$ (i.e., large space-like momentum transfer). It is also not difficult to see that

Σ -contribution in Eq. (A-4) also vanishes at $q^2 = -\infty$ independently of the value of η .

In the following we discuss the large q^2 -behavior of the W-contribution in Eqs. (A-19) and (A-20). Those equations show that all the contributions to $F(q^2)$ vanishes at $q^2 = -\infty$ for $\xi = 1$ (i.e., 't-Hooft-Feynman gauge). However the gauge independence of the off-shell amplitude cannot be proved. We expect that $F(q^2)$ at $q^2 = -\infty$ may depend on the gauge one chooses. We show this explicitly in the case of the neutrino contribution in the Weinberg model. We thus put $m=0$ in Eqs. (A-19) and (A-20).

We first note the following relations

$$\begin{aligned}
 & -q^2 \int dx dy \frac{xy}{Q^2 + x(\Lambda - \mu^2) + y(M^2 - \mu^2)} \\
 & \xrightarrow{q^2 \rightarrow -\infty} \ln(-q^2) + \int_0^1 dt \ln\left(\frac{t}{\mu^2 t + M^2 - \mu^2}\right)
 \end{aligned} \tag{A-22}$$

and

$$\begin{aligned}
 & -q^2 \int dx dy \rho(x, y) \ln f(M^2, \Lambda) \\
 & \xrightarrow{q^2 \rightarrow -\infty} -q^2 \int_0^1 dt \rho(t) t \ln t \\
 & \quad + (\Lambda + M^2 - 2\mu^2) \int_0^1 dt \rho(t) \ln(-q^2 t) \\
 & \quad + \frac{1}{2} (\Lambda - M^2) \int_0^1 dt \rho(t) \ln \left[\frac{\mu^2 t + M^2 - \mu^2}{\mu^2 t + \Lambda - \mu^2} \right] \\
 & \quad - \frac{1}{2} \int_0^1 dt \rho(t) \left[\mu^2 t - \mu^2 + \frac{\Lambda + M^2}{2} \right] \ln \left\{ (\mu^2 t + M^2 - \mu^2)(\mu^2 t + \Lambda - \mu^2) \right\} \\
 & \quad + (\Lambda + M^2 - 2\mu^2) \int_0^1 dt \rho(t)
 \end{aligned} \tag{A-23}$$

where $f(x, y) \equiv 4(x+y)^2 - q(x+y) + 4$

and $g(t) \equiv 4t^2 - qt + 4.$

Using these two relations in Eq. (A-20) we can readily show that

$$\begin{aligned}
 F(q^2) &\rightarrow 0 \quad \text{for } q^2 \rightarrow -\infty \quad \text{and } \xi \neq 0 \\
 F(q^2) &\rightarrow \frac{1}{8\pi^2} \frac{g\mu^2}{8M^2} \left(4 - \frac{\mu^2}{M^2}\right) \ln(-q^2) + \text{const} \quad (\text{A-24}) \\
 &\quad \text{for } q^2 \rightarrow -\infty \quad \text{after } \xi = 0
 \end{aligned}$$

(A-25)

Eqs. (A-24) and (A-25) indicate the gauge independence of the off-shell amplitude.

We also point out an interesting large q^2 -behavior obtained if one uses (incorrect) regularization schemes other than the ξ -limiting procedure in the U-gauge. The last two terms in Eq. (A-20) show that the linear divergence in q^2 at large q^2 could exist. For the gauge invariant calculation this linear divergence cancels. But if one uses other regularization schemes such as the "proper time" (see Appendix B) or the "Pauli-Villars" regularization with a massive neutrino in the U-gauge, this linear divergence indeed survives.

APPENDIX B. Ambiguities in the U-gauge

In this appendix we briefly review the ambiguity Jackiw and Weinberg, and also Bars and Yoshimura⁸ encountered in their calculation of the muon g-factor.

The logarithmic term in the parametric integral for $F(0)$ in the U-gauge limit (i.e. $\xi \rightarrow 0$) causes an ambiguity: Eq. (A-20) in Appendix A contains the following logarithmic term

$$F(0) = -\frac{1}{8\pi^2} \frac{g^2 \mu^2}{4M^2} \int dx dy [4(x+y)^2 - 9(x+y) + 2 + 6y] \times \\ \times \ln \left[\frac{x}{\mu^2(x+y)^2 + (x+y)(M^2 - \mu^2)} \right] + \text{other terms.} \quad (\text{A-1})$$

This is the correct answer. On the other hand if one regulates the neutrino propagator we get

$$F(0) = -\frac{1}{8\pi^2} \frac{g^2 \mu^2}{4M^2} \int dx dy [4(x+y)^2 - 9(x+y) + 2 + 6y] \times \\ \times \ln \left[\frac{1-x-y}{\mu^2(x+y)^2 + (x+y)(M^2 - \mu^2)} \right] + \text{other terms.} \quad (\text{A-2})$$

If one first exponentiates the Feynman amplitude (a sort of "proper time") and performs a loop integral, the following result is obtained

$$F(0) = -\frac{1}{8\pi^2} \frac{g^2 \mu^2}{4M^2} \int dx dy [4(x+y)^2 - 9(x+y) + 2 + 6y] \times \\ \times \ln \left[\frac{1}{\mu^2(x+y)^2 + (x+y)(M^2 - \mu^2)} \right] + \text{other terms.} \quad (\text{A-3})$$

All these expressions give rise to different answers. This kind of ambiguity is absent in the general R-gauge calculation. Naive U-gauge calculations are plagued by this kind of ambiguity. Some existing proofs of cancellations of divergences in higher order diagrams are based on the exponential parametrization of propagators that leads to (B-3). While such a method is acceptable in establishing the absence of divergences, it will not provide a reliable finite part.

APPENDIX C. Lagrangians and Feynman Rules.

In this Appendix we present Lagrangians for the existing models of leptons mentioned in the Introduction. We also write^{down} necessary Feynman rules for our calculations in Sections III and IV. In the R_ξ -gauge

discussed in Section II, we have unphysical scalars in the Lagrangian. The Feynman rules are therefore more complicated than those in the U-gauge limit. The Feynman rules for those unphysical scalars and also the relative signs for various amplitudes can be conveniently checked based on the gauge independence of the T-matrix element as we discussed in Section II.

CI. Georgi-Glashow Model

This model is based on the group $O(3)$. We have a triplet of leptons and also a singlet of neutral massive lepton. The mass is generated by a triplet of real scalars. A part of this Lagrangian has been given in Section II (We follow the notation of Bjorken and Drell¹⁶). The total Lagrangian has the following form:

$$\mathcal{L} = \mathcal{L}_I + \mathcal{L}_{II} + \mathcal{L}_{III} - V$$

$$\begin{aligned} \mathcal{L}_I \equiv & \bar{Y}^+ (i\not{\partial} - m_{Y^+}) Y^+ + \bar{Y}^0 (i\not{\partial} - m_{Y^0}) Y^0 \quad (C-1) \\ & + \bar{\mu} (i\not{\partial} - m_\mu) \mu + \bar{\nu}_L i\not{\partial} \nu_L + e \bar{L} \gamma^\mu (\vec{T} \cdot \vec{B})_\mu L \\ & + e \bar{R} \gamma^\mu (\vec{T} \cdot \vec{B})_\mu R - \frac{e(m_{Y^0} \cos \theta - m_\mu)}{M} [\bar{L} (\vec{T} \cdot \vec{S}) R + h.c.] \\ & - \frac{e m_{Y^0} \sin \theta}{M} \left\{ [\sin \theta \bar{Y}_L^0 - \cos \theta \bar{\nu}_L] [S^- Y_R^+ + S^+ \mu_R + \psi Y_R^0] + h.c. \right\} \end{aligned}$$

where

$$L \equiv \begin{bmatrix} Y_L^+ \\ \cos\theta Y_L^0 + \sin\theta Y_L^- \\ \mu_L \end{bmatrix}, \quad R \equiv \begin{bmatrix} Y_R^+ \\ Y_R^0 \\ \mu_R \end{bmatrix} \quad (C-2)$$

with

$$\mu_{R/L} \equiv \left(\frac{1 \pm \gamma_5}{2} \right) \mu, \quad \text{etc.} \quad (C-3)$$

$$\vec{T} \cdot \vec{B}_\mu = \begin{bmatrix} A_\mu & -W_\mu^+ & 0 \\ -W_\mu^- & 0 & W_\mu^+ \\ 0 & W_\mu^- & -A_\mu \end{bmatrix}, \quad \vec{T} \cdot \vec{S} = \begin{bmatrix} \psi & -s^+ & 0 \\ -s^- & 0 & s^+ \\ 0 & s^- & -\psi \end{bmatrix} \quad (C-4)$$

where ψ = Higgs scalar

$$S^\pm = \text{unphysical scalars} \quad (\text{C-5})$$

$$\text{and } G_F = (e^2 \sin^2 \theta / 4 M^2)$$

$$\text{There is a constraint : } 2 m_{Y^0} \cos \theta = m_{Y^+} + m_{\mu} \quad (\text{C-6})$$

Note that the covariant derivative is given by $\nabla_\mu \psi \equiv [\partial_\mu + i(\vec{T} \cdot \vec{B})_\mu] \psi$

with $g \equiv -e$.

$$\begin{aligned} \mathcal{L}_{II} \equiv & \frac{1}{2} \left| \partial_\mu \psi + i e [\psi_\mu^- s^+ - W_\mu^+ s^-] \right|^2 \\ & + \left| \partial_\mu s^+ + i M W_\mu^+ - i e A_\mu s^+ + i e \psi W_\mu^+ \right|^2 \end{aligned} \quad (\text{C-7})$$

The quadratic term of \mathcal{L}_{II} is given by

$$\begin{aligned} \mathcal{L}_{II}^{\text{quad.}} = & \left| \partial_\mu s^+ \right|^2 + \frac{1}{2} (\partial_\mu \psi)^2 + M^2 |W_\mu^+|^2 \\ & - i M [\partial_\mu s^+ W^{\mu -} - \partial_\mu s^- W^{\mu +}] \end{aligned} \quad (\text{C-8})$$

This $\mathcal{L}_{II}^{\text{quad.}}$ suggests the following gauge term

$$\begin{aligned} \mathcal{L}^c = & - \frac{1}{2\alpha} (\partial_\mu A^\mu)^2 \\ & - \xi \left| \partial_\mu W^{\mu +} + \frac{iM}{\xi} s^+ \right|^2 \end{aligned} \quad (\text{C-9})$$

See also Section II.

$$\begin{aligned} \mathcal{L}_{\text{III}} = & -\frac{1}{2} \left| \partial_\mu W_\nu^+ - \partial_\nu W_\mu^+ + ie [W_\mu^+ A_\nu - W_\nu^+ A_\mu] \right|^2 \\ & - \frac{1}{4} \left| \partial_\mu A_\nu - \partial_\nu A_\mu - ie [W_\mu^+ W_\nu^- - W_\mu^- W_\nu^+] \right|^2 \end{aligned} \quad (\text{C-10})$$

The mass of W_μ^\pm is given in Eq. (C-8).

The potential is given by

$$\begin{aligned} V(\phi) &= \frac{1}{2} \mu_0^2 |\phi|^2 + \lambda |\phi|^4 \\ &= \frac{1}{2} m_\psi^2 \psi^2 + \lambda \left[4v\psi(2s^+s^- + \psi^2) + (2s^+s^- + \psi^2)^2 \right] \\ &\quad + [\mu_0^2 + 4v^2\lambda] [s^+s^- + v\psi] \end{aligned} \quad (\text{C-11})$$

where

$$m_\psi^2 = \mu_0^2 + 12\lambda v^2$$

(C-12)

$\mu_0^2 < 0$ and we have the condition $\mu_0^2 + 4v^2\lambda = 0$.

The field ϕ is the unshifted field, and it is expressed as

$$\phi = \begin{pmatrix} \phi^+ \\ \phi^0 \\ \phi^- \end{pmatrix} = \begin{pmatrix} s^+ \\ v+\psi \\ s^- \end{pmatrix}, \quad \langle \phi \rangle_0 = \begin{pmatrix} 0 \\ v \\ 0 \end{pmatrix}. \quad (\text{C-13})$$

The gauge compensating term for Eq. (C-9) is given by (see also Section II)

$$-i T_R \ln [1 + \mathcal{Q} \gamma] \quad (\text{C-14})$$

where

$$\mathcal{Q} = \begin{bmatrix} -(\partial^2 + \frac{M^2}{\xi} - i\epsilon) & 0 & 0 \\ 0 & -\partial^2 + i\epsilon & 0 \\ 0 & 0 & -(\partial^2 + \frac{M^2}{\xi} - i\epsilon) \end{bmatrix} \quad \mathcal{Q} = \int^4 (x-y) \quad (\text{C-15})$$

and

$$\gamma = \begin{bmatrix} ie\partial_\mu A^\mu - \frac{eM}{\xi}\psi & -ie\partial_\mu W^{+\mu} + \frac{eM}{\xi}s^+ & 0 \\ -ie\partial_\mu W^{-\mu} & 0 & ie\partial_\mu W^{+\mu} \\ 0 & ie\partial_\mu W^{-\mu} + \frac{eM}{\xi}s^- & -ie\partial_\mu A^\mu - \frac{eM}{\xi}\psi \end{bmatrix} \delta^4(x-y) \quad (\text{C-16})$$

The divergence in this matrix stands for an operator, e.g., $\partial_\mu A^\mu \equiv (\partial_\mu A^\mu) + A^\mu \partial_\mu$. The lowest order contribution from Eq. (C-14) is a ψ -vacuum tadpole diagram. For $\xi = \infty$ (Landau-gauge), for example, Eq. (C-14) becomes

$$-iT_R \ln \left[1 + \frac{1}{-\partial^2 + i\epsilon} \partial_\mu (ie \vec{T} \cdot \vec{B}^\mu) \right] \quad (\text{C-17})$$

C II Weinberg Model

The detailed form of this Lagrangian is found in references 1, 6. We briefly summarize it in the following

$$\mathcal{L} \equiv \mathcal{L}_I + \mathcal{L}_{II} + \mathcal{L}_{III} - V$$

$$\begin{aligned} \mathcal{L}_I = & \bar{\nu}_L i \not{\partial} \nu_L + \bar{\mu} (i \not{\partial} - m_\mu) \mu \\ & - e \bar{\mu} \not{A} \mu - \frac{g}{i2} \left[\bar{\nu}_L \not{X}^\dagger \mu + \text{h.c.} \right] - \frac{G}{2} \bar{\nu}_L \not{\nu}_L \\ & + G \bar{\mu} \not{\chi} \left[\frac{\cos 2\theta}{2} \left(\frac{1-\gamma_5}{2} \right) - \sin^2 \theta \left(\frac{1+\gamma_5}{2} \right) \right] \mu \\ & - \frac{g m_\mu}{2M} \left[(\sqrt{2} \bar{\mu} \nu_L S^- + \text{h.c.}) + \bar{\mu} \mu \psi + \bar{\mu} i \gamma_5 \mu \chi \right] \end{aligned} \quad (\text{C-18})$$

where

S^\pm and χ = unphysical scalar, ψ = Higgs scalar

$$G = \sqrt{g^2 + g'^2}$$

$$\cos \theta = \frac{g}{G} = \frac{M}{M_Z}, \quad e = -\frac{g g'}{G}, \quad G_F = \frac{g^2}{8M^2} \quad (\text{C-19})$$

For notational convenience we defined the charge e with an extra minus sign.

$$\begin{aligned} \mathcal{L}_{II} = & \left| \partial_\mu s^+ + i M W_\mu^+ + \frac{i}{\sqrt{2}} g W_\mu^+ s^0 + i \left(-e A_\mu + \frac{G \cos 2\theta}{2} Z_\mu \right) \right|^2 \\ & + \left| \partial_\mu s^0 - i \frac{M_Z}{\sqrt{2}} Z_\mu - i \frac{G Z_\mu}{2} s^0 + \frac{i}{\sqrt{2}} g W_\mu^- s^+ \right|^2 \end{aligned} \quad (C-20)$$

where $s^0 \equiv \frac{1}{\sqrt{2}} (\psi + i\chi)$.

The quadratic part of \mathcal{L}_{II} is given by

$$\begin{aligned} \mathcal{L}_{II}^{\text{quad}} = & |\partial_\mu s^+|^2 + \frac{1}{2} [(\partial_\mu \psi)^2 + (\partial_\mu \chi)^2] + M^2 |W_\mu^+|^2 \\ & + \frac{1}{2} M_Z^2 (Z_\mu)^2 - i M [\partial_\mu s^+ W^{-\mu} - \partial_\mu s^- W^{+\mu}] - M_Z \partial_\mu \chi Z^\mu \end{aligned} \quad (C-21)$$

The gauge term is given as

$$\begin{aligned} \mathcal{L}^G = & -\frac{1}{2\alpha} (\partial_\mu A^\mu)^2 \\ & - \xi \left| \partial_\mu W^{+\mu} + i \frac{M}{\xi} s^+ \right|^2 \\ & - \frac{\eta}{2} \left(\partial_\mu Z^\mu + \frac{M_Z}{\eta} \chi \right)^2 \end{aligned} \quad (C-22)$$

The gauge compensating term will be discussed later.

$$\begin{aligned}
\mathcal{L}_{III} = & -\frac{1}{2} \left| \partial_\mu W_\nu^+ - \partial_\nu W_\mu^+ + ie [W_\mu^+ A_\nu - W_\nu^+ A_\mu] - iG \cos^2 \theta [W_\mu^+ Z_\nu - W_\nu^+ Z_\mu] \right|^2 \\
& -\frac{1}{4} \left| \partial_\mu A_\nu - \partial_\nu A_\mu - ie [W_\mu^+ W_\nu^- - W_\nu^+ W_\mu^-] \right|^2 \\
& -\frac{1}{4} \left| \partial_\mu Z_\nu - \partial_\nu Z_\mu + iG \cos^2 \theta [W_\mu^+ W_\nu^- - W_\nu^+ W_\mu^-] \right|^2
\end{aligned}
\tag{C-23}$$

The masses for W_μ and Z_μ are given in Eq. (C-21).

$$\begin{aligned}
V = & \lambda \left[S^+ S^- + \left(S^0 + \frac{v}{\sqrt{2}} \right) \left(\bar{S}^0 + \frac{v}{\sqrt{2}} \right) - \frac{v^2}{2} \right]^2 \\
= & \frac{1}{2} m_\psi^2 \psi^2 + \lambda \left\{ \sqrt{\frac{2}{\lambda}} m_\psi \psi (S^+ S^- + S^0 \bar{S}^0) + [S^+ S^- + S^0 \bar{S}^0]^2 \right\}.
\end{aligned}
\tag{C-24}$$

The gauge compensating term is given by

$$-i T_n \ln [1 + \mathcal{G} \cdot \gamma]
\tag{C-25}$$

where

$$\left[\begin{array}{cccc}
-(\partial^2 - \frac{M^2}{\xi} - i\epsilon) & 0 & 0 & 0 \\
0 & -(\partial^2 + \frac{M^2}{\xi} - i\epsilon) & 0 & 0 \\
0 & 0 & -\partial^2 + i\epsilon & 0 \\
0 & 0 & 0 & -(\partial^2 + \frac{M^2}{\eta} - i\epsilon)
\end{array} \right] \mathcal{G} = \int^4 (x-y)
\tag{C-26}$$

and the matrix γ is defined by

$$\begin{bmatrix} \delta f^+ \\ \delta f^- \\ \delta f^A \\ \delta f^Z \end{bmatrix} \equiv [\gamma] \begin{bmatrix} \Omega^+ \\ \Omega^- \\ \Omega^A \\ \Omega^Z \end{bmatrix} \equiv [\gamma] \begin{bmatrix} \frac{1}{\sqrt{2}} (\Omega^+ - i \Omega^Z) \\ \frac{1}{\sqrt{2}} (\Omega^+ + i \Omega^Z) \\ \frac{g'}{g} \Omega^3 + \frac{2g}{g'} \Omega^0 \\ \Omega^3 - 2 \Omega^0 \end{bmatrix} \quad (C-27)$$

with

$$\begin{aligned} \delta f^\pm &\equiv \mp i g \partial^\mu \left\{ [Z_\mu \cos \theta + A_\mu \sin \theta] \Omega^\pm \right\} - \frac{g M}{2 \xi} (\psi \pm i \chi) \Omega^\pm \\ &\quad \mp i e \cos \theta \partial^\mu \left\{ W_\mu^\pm [\Omega^A + \cot \theta \Omega^Z] \right\} + \frac{e M \cos \theta}{2 \xi} s^\pm \left[2 \Omega^A - \frac{G}{e} \cos 2\theta \Omega^Z \right] \\ \delta f^A &\equiv i G \sin \theta \partial^\mu [W_\mu^- \Omega^+ - W_\mu^+ \Omega^-] \\ \delta f^Z &\equiv i g \partial^\mu [W_\mu^- \Omega^+ - W_\mu^+ \Omega^-] \\ &\quad + \frac{G M_Z}{2 \eta} [-\psi \Omega^Z + s^+ \Omega^- + s^- \Omega^+] \end{aligned} \quad (C-28)$$

Another way to take care of this compensating term is to introduce four auxiliary complex scalar fields ¹² φ_a , $a = (+, -, A, Z)$. We add the following extra piece to the effective Lagrangian

$$\tilde{\mathcal{L}} \equiv \varphi_a^\dagger [\mathcal{L}^{-1} + \gamma]^{a,b} \varphi_b \quad (C-29)$$

The ordinary perturbative treatment of $\tilde{\mathcal{L}}$ with an extra (-) sign for each closed loop of the fictitious scalar particles φ_a gives rise to Eq. (C-25)

C III The Lepton Model by Lee, Prentki and Zumino

The group structure of this model is close to that of the Weinberg model. We need two sets of scalar triplets to accomodate "quarks" in this model. However only one triplet of complex scalars are sufficient for the lepton model. We present here this simplified lepton model.

$$\begin{aligned}
 \mathcal{L} &= \mathcal{L}_I + \mathcal{L}_{II} + \mathcal{L}_{III} - V \\
 \mathcal{L}_I &= \bar{M}^+ (\not{\partial} - m_{M^+}) M^+ + \bar{\nu}_L \not{\partial} \nu_L + \bar{\mu} (\not{\partial} - m_\mu) \mu \\
 &\quad - \frac{g m_{M^+}}{M} [(\bar{L} \cdot \tilde{S}) M_R^+ + h.c.] - \frac{g m_\mu}{M} [(\bar{L} \cdot S) \mu_R + h.c.] \\
 &\quad - e [\bar{\mu} \not{A} \mu - \bar{M}^+ \not{A} M^+] \\
 &\quad - g \left\{ (\bar{\nu} \not{X}^+ \mu - \bar{M}^+ \not{X}^+ \nu) + h.c. \right\} \\
 &\quad + G \bar{\mu} \not{Z} \left\{ \cos^2 \theta \left(\frac{1-\gamma_5}{2} \right) - \sin^2 \theta \left(\frac{1+\gamma_5}{2} \right) \right\} \mu \\
 &\quad - G \bar{M} \not{Z} \left\{ \cos^2 \theta \left(\frac{1-\gamma_5}{2} \right) - \sin^2 \theta \left(\frac{1+\gamma_5}{2} \right) \right\} M^+
 \end{aligned} \tag{C-30}$$

where

$$L \equiv \begin{bmatrix} M^+ \\ \nu_L \\ \mu_L \end{bmatrix}, \quad S \equiv \begin{bmatrix} s^{++} \\ s^+ \\ s^0 \end{bmatrix}, \quad \tilde{S} \equiv \begin{bmatrix} \bar{s}^0 \\ s^- \\ s^{--} \end{bmatrix} \tag{C-31}$$

$$G = \sqrt{g^2 + g'^2}, \quad e = (-) \frac{g g'}{G}, \quad G_F = \frac{g^2}{4M^2},$$

$$\cos \theta = \frac{g}{G}, \quad M = g u, \quad M_Z = \sqrt{2} G u.$$

$$\begin{aligned}
\mathcal{L}_{II} = & \left| \partial_\mu S^{++} - i g W_\mu^+ S^+ + i (-2eA_\mu + G \cos 2\theta Z_\mu) S^{++} \right|^2 \\
& + \left| \partial_\mu S^+ + i M W_\mu^+ + i g W_\mu^+ S^0 - i g W_\mu^- S^{++} - i [eA_\mu + G \sin^2 \theta Z_\mu] S^+ \right|^2 \\
& + \left| \partial_\mu S^0 - i \frac{M_Z}{\sqrt{2}} Z_\mu + i g W_\mu^- S^+ - i G Z_\mu S^0 \right|^2
\end{aligned}$$

(C-32)

The quadratic part of \mathcal{L}_{II} is given by

$$\begin{aligned}
\mathcal{L}_{II}^{quad} = & \left| \partial_\mu S^{++} \right|^2 + \left| \partial_\mu S^+ \right|^2 + \frac{1}{2} \left[(\partial_\mu \psi)^2 + (\partial_\mu \chi)^2 \right] \\
& + M^2 |W_\mu^+|^2 + \frac{1}{2} M_Z^2 (Z_\mu)^2 \\
& - i M \left[\partial^\mu S^+ W_\mu^- - \partial^\mu S^- W_\mu^+ \right] - M_Z \partial_\mu \chi \cdot Z^\mu
\end{aligned}$$

(C-33)

where we defined

$$S^0 \equiv \frac{1}{\sqrt{2}} (\psi + i\chi)$$

S^\pm and χ = unphysical scalars

(C-34)

$S^{\pm\pm}$ and ψ = Higgs' scalars.

The gauge term is the same as Eq. (C-22) in the Weinberg Model. The Yang-Mills Lagrangian is also given by Eq. (C-23).

Finally we discuss the potential.

$$\begin{aligned}
 V &= \mu_0 (\vec{\xi}^\dagger \vec{\xi}) + \frac{\nu}{4} (\vec{\xi}^\dagger \times \vec{\xi})^2 + \lambda (\vec{\xi}^\dagger \vec{\xi})^2 \\
 &= \frac{1}{2} m_\psi^2 \psi^2 + \frac{1}{2} m_{S^{++}}^2 |S^{++}|^2 + 2\sqrt{2} \lambda u \psi [S^0 \bar{S}^0 + S^+ S^- + S^{++} S^{--}] \\
 &\quad + \lambda [S^0 \bar{S}^0 + S^+ S^- + S^{++} S^{--}]^2 - \frac{\nu}{4} \left\{ 2\sqrt{2} u \psi (S^0 \bar{S}^0 + S^{++} S^{--}) \right. \\
 &\quad + 2u S^+ (S^0 \bar{S}^- - S^+ S^{--}) + 2u S^- (\bar{S}^0 S^+ - S^- S^{++}) \\
 &\quad \left. + (S^0 \bar{S}^0 - S^{++} S^{--})^2 + 2(\bar{S}^0 S^+ - S^- S^{++})(S^0 \bar{S}^- - S^+ S^{--}) \right\} \quad (C-35)
 \end{aligned}$$

where $\vec{\xi}$ is the unshifted complex scalar field

$$\vec{\xi} = \begin{bmatrix} S^{++} \\ S^+ \\ u + S^0 \end{bmatrix} ; \quad \langle \vec{\xi} \rangle_0 = \begin{bmatrix} 0 \\ 0 \\ u \end{bmatrix} \quad (C-36)$$

$$m_\psi^2 = -2\mu_0 > 0, \quad m_{S^{++}}^2 = \nu u^2 > 0$$

There is ^{the} following constraint:

$$\lambda = \frac{\nu}{4} + \left(-\frac{\mu_0}{2u} \right) = \frac{i}{4u^2} [m_\psi^2 + m_{S^{++}}^2] \quad (C-37)$$

We can make m_ψ and $m_{S^{++}}$ arbitrarily large. The gauge compensating term is also similar to that of the Weinberg model. We do not discuss it here.

C IV Feynman Rules

We summarize several Feynman rules we use in Sections III and IV. Propagators for vector bosons and also for scalars are found in Section II. The Fermion propagator has the standard form

$$\frac{i}{\not{p} - m} \quad (C-38)$$

In the following we give Feynman rules for the lepton models due to Georgi and Glashow (GG), Weinberg (W), and Lee, Prentki and Zumino (LPZ). We write the Feynman rules for the R_ξ -gauge of Section II.

All of these models give the identical Feynman rules for Figs. 8a ~ 8d.

They are

$$(8a) \quad (-ie) \bar{\mu} \gamma_\alpha \mu \quad (C-39)$$

$$(8b) \quad (-ie) (\ell' + \ell)_\mu \quad (C-40)$$

$$(8c) \quad (ie) \left\{ (g_{\alpha\beta} (\ell' + \ell)_\mu - \ell_\beta g_{\mu\alpha} - \ell'_\alpha g_{\mu\beta}) + (g_{\alpha\mu} g_{\beta\mu} - g_{\beta\mu} g_{\alpha\mu}) \right\} \quad \text{when } q = \ell' - \ell \quad (C-41)$$

$$(8d) \quad (-ie) M g_{\alpha\mu} \quad (C-42)$$

Note that we normalized the sign of the charge by Fig. 8a. The magnetic form factor appears as a coefficient of $(-ie) \bar{u} [i \sigma_{\mu\nu} q^\nu / 2m] u$

For other diagrams Feynman rules depend on the model. We just list them below

$$(8e) \quad \begin{aligned} GG: & \quad (ie \sin \theta) \bar{\nu} \gamma_\alpha \left(\frac{1-\gamma_5}{2} \right) \mu \\ W: & \quad \left(-\frac{ig}{2} \right) \bar{\nu} \cdot \gamma_\alpha \left(\frac{1-\gamma_5}{2} \right) \mu \\ LPZ: & \quad (-ig) \bar{\nu} \gamma_\alpha \left(\frac{1-\gamma_5}{2} \right) \mu \end{aligned} \quad (C-43)$$

$$(8f) \quad \begin{aligned} GG: & \quad \left(\frac{ie \sin \theta m_\mu}{M} \right) \bar{\nu} \left(\frac{1+\gamma_5}{2} \right) \mu \\ W: & \quad \left(-\frac{ig m_\mu}{\sqrt{2} M} \right) \bar{\nu} \left(\frac{1+\gamma_5}{2} \right) \mu \\ LPZ: & \quad \left(-\frac{ig m_\mu}{M} \right) \bar{\nu} \left(\frac{1+\gamma_5}{2} \right) \mu \end{aligned} \quad (C-44)$$

$$(8g) \quad GG : (ie) \quad \bar{\gamma}_0 \left[\cos \theta \left(\frac{1+\gamma_5}{2} \right) + \left(\frac{1-\gamma_5}{2} \right) \right] \gamma_\alpha \mu$$
(C-45)

$$(8h) \quad GG : \left(\frac{ie}{M} \right) \bar{\gamma}_0 \left[(m_\mu \cos \theta - m_{\gamma_0}) \left(\frac{1+\gamma_5}{2} \right) + (m_\mu - m_{\gamma_0} \cos \theta) \left(\frac{1-\gamma_5}{2} \right) \right] \mu$$
(C-46)

$$(8i) \quad W : i G \bar{\mu} \left[\frac{\cos 2\theta}{2} \left(\frac{1+\gamma_5}{2} \right) - \sin^2 \theta \left(\frac{1-\gamma_5}{2} \right) \right] \gamma_\alpha \mu$$

$$LPZ : i G \bar{\mu} \left[\cos^2 \theta \left(\frac{1+\gamma_5}{2} \right) - \sin^2 \theta \left(\frac{1-\gamma_5}{2} \right) \right] \gamma_\alpha \mu$$
(C-47)

$$(8j) \quad W : \left(-i \frac{g m_\mu}{2 M} \right) \bar{\mu} i \gamma_5 \mu$$

$$LPZ : \left(-i \frac{g m_\mu}{4 M} \right) \bar{\mu} i \gamma_5 \mu$$
(C-48)

FOOTNOTES AND REFERENCES

1. S. Weinberg, Phys. Rev. Letters 19, 1264 (1967); 27, 1688 (1972).
2. G. 't Hooft, Nuclear Phys. B35, 167 (1971).
3. T. D. Lee and C. N. Yang, Phys. Rev. 128, 885 (1962).
4. The R_ξ -gauge was formulated independently by J. Zinn-Justin.
The proof of renormalizability of SGGT in this gauge is presently pursued by Zinn-Justin and one of the present authors (P.W.L.)
What is to be proved is that divergences in higher orders of Green's functions can be removed by renormalization in such a way that the S-matrix is independent of ξ .
5. B. W. Lee and J. Zinn-Justin, Phys. Rev. D, to be published;
B. W. Lee and J. Zinn-Justin, Phys. Rev. D, to be published.
6. A. Salam and J. Strahdee, to be published.
7. R. Jackiw and S. Weinberg, Phys. Rev. D5, 2396 (1972); S. Y. Lee
UCSD-10P10-97-99 (1972). To be published in Phys. Rev.; T. W.
Appelquist, J. R. Primack, and H. R. Quinn, Harvard preprint (1972).
8. R. Jackiw and S. Weinberg, Phys. Rev. D5, 2396 (1972); I. Bars
and M. Yoshimura, Berkeley Preprint to be published in Phys. Rev.
9. Some of the numerical values for the weak correction to the muon
magnetic moment have been previously evaluated without using the
lepton models based on SGGT. T. Burnett and J. J. Levine,
Phys. Letters 24, 467 (1967); S. J. Brodsky and J. Sullivan,
Phys. Rev. 156, 1644 (1967); T. Kinoshita et al., Phys. Rev. D2,
910 (1970).

Footnotes and References (continued)

10. H. Georgi and S. L. Glashow, Phys. Rev. Letters 28, 1494 (1972).
11. B. W. Lee, Phys. Rev. D to be published; J. Prentki and B. Zumino, Nuclear Phys. B to be published.
12. G. 't Hooft, Nuclear Phys. B33, 173, (1971).
13. For details see, for example, Ref. 5.
14. The physical scalar bosons are called Higgs' scalars, and denotes by Ψ , as opposed to unphysical scalars which are denotes by s^{\pm} .
15. For summer, the review of the recent advances in this field, see S. Brodsky Proceedings 1971 International Symposium on Electron and Photon Interaction at High Energy, Editor N. Mistry Laboratory of Unclear Studies Cornell University (1971).
16. J. D. Bjorken and S. D. Drell Relativistic Quantum Fields McGraw-Hill Book Company (1965).
17. G. Altarelli, N. Cabibbo and L. Maiani ISS 72/3.
18. J. L. Gervais and B. W. Lee, Nuclear Phys. B12, 627, (1969).
19. S. L. Adler and W. A. Bardeen, Phys. Rev. 182, 1517 (1969).

FIGURE CAPTIONS

- Fig. 1. W boson and scalar boson contributions to weak correction to the muon magnetic moment. These give contributions in all three models.
- Fig. 2. Z and χ boson contribution to the weak correction to the muon magnetic moment. These diagrams are for Weinberg's model. Similar sets of diagrams exist for the LPZ model.
- Fig. 3. Y^0 lepton contribution to the weak correction to the muon magnetic moment. These diagrams are for Georgi-Glashow model.
- Fig. 4. Predictions of Eq. 3.12. Sets of contours correspond to constant a_μ^W and constant M_{Y^0} on the M_{Y^+}, M plane. If we take $-2 \times 10^{-7} \leq a_\mu^W \leq 0$, for example, the experimentally allowed region lies below the line labeled -2×10^{-7} . If we further take $M_{Y^+} \geq .5 \text{ GeV}$, the allowed region is bounded from below. The upper bound for M_{Y^+} , in any case, is approximately 5 GeV.
- Fig. 5. Example of diagrams contributing to muon elastic scattering.
- Fig. 6. Diagrams which contribute to the static charge of the neutrino in the Georgi-Glashow model.
- Fig. 7. Diagrams contributing to the neutron static charge.
- Fig. 8. Several vertex diagrams for lower order calculations.

Tables

1. Contributions of diagrams shown in Fig. 1. To obtain the answer these numbers should be multiplied by $G\mu^2/8\pi^2\sqrt{2}$.
2. Contributions of diagrams shown in Fig. 2. To obtain the answer these numbers should be multiplied by $G\mu^2/8\pi^2\sqrt{2}$.
3. Contributions of diagrams shown in Fig. 3. To obtain the answer these numbers should be multiplied by $G\mu^2/2\pi^2\sqrt{2}\sin^2\theta$.







Figure 1




Figure 2







FIGURE 3







Figure 4



(a)



(b)

Figure 5








Figure 6




Figure 7













Figure 8




Diagram Gauge	 (a)	 (b) + (c)	 (d)	Total
U-Gauge	$\frac{10}{3}$	0	0	$\frac{10}{3}$
't Hooft-Feynman-Gauge	$\frac{7}{3}$	1	0	$\frac{10}{3}$
R-Gauge	$\frac{4}{3}$	1	1	$\frac{10}{3}$

Table 1



Diagram Gauge	 (a)	 (b)	Total
U-Gauge	$\frac{1}{3} [(3-4\cos^2 \theta)^2 - 5]$	0	$\frac{1}{3} [(3-4\cos^2 \theta)^2 - 5]$
't Hooft-Feynman-Gauge	$\frac{1}{3} [(3-4\cos^2 \theta)^2 - 5]$	0	$\frac{1}{3} [(3-4\cos^2 \theta)^2 - 5]$
R-Gauge	$\frac{1}{3} [(3-4\cos^2 \theta)^2 - 5] + 1$	-1	$\frac{1}{3} [(3-4\cos^2 \theta)^2 - 5]$

Table 2





Diagram Gauge	 (a)	 (b) +  (c)	 (d)	Total
U - Gauge	-1	0	0	-1
't Hooft- Feynman- Gauge	$-\frac{3}{4}$	$-\frac{1}{4}$	0	-1
R - Gauge	$-\frac{1}{2}$	$-\frac{1}{4}$	$-\frac{1}{4}$	-1

Table 3