



# Fermi National Accelerator Laboratory

FERMILAB-Pub-76/53-THY  
June 1976

## Second-Class Currents and Very Light Quarks

A. HALPRIN

University of Delaware, Newark, Delaware 19711

AND

B. W. LEE and P. SORBA<sup>†</sup>

Fermi National Accelerator Laboratory, Batavia, Illinois 60510

### ABSTRACT

We consider induced second-class nucleon currents in a light quark model generated by unequal  $u$  and  $d$  quark masses. The effect produced by gluon vertex corrections is strikingly large without confinement effects. Restricting the quarks to the limited spatial region of a physical nucleon changes the energy scale from the quark mass to the quark-bound state energy and thereby reduces the second-class form factors well below their first class counterparts.

---

<sup>†</sup>On leave of absence from Centre de Physique Theorique -CNRS, Marseille, France.

## I. INTRODUCTION

The results of recent angular correlation measurements in the  $\beta$  decay of some light nuclei have opened a new chapter in the tortured history of second-class weak interactions. The experiments of Sugimoto, et al., on the mass 12 system and the experiment of Calaprice, et al., on the mass 19 system support the existence of second-class currents.<sup>1,2</sup> In contrast, the experiment of Nathan, et al., on the mass 8 system does not find a comparable second class effect, in agreement with older experiments on this system.<sup>3,4,5</sup> The experimental situation is, therefore, in a very unsettled state.<sup>6</sup>

It is well known that second-class currents are alien to conventional quark models with degenerate  $u$  and  $d$  quark masses.<sup>7</sup> However, it is not clear if an appreciable second-class current can be induced in a model in which  $(m_d - m_u)/(m_d + m_u)$  is of order unity. Such a mass relation has been suggested by Leutwyler<sup>8</sup> and can also be accommodated in the MIT bag model.<sup>9</sup> In this paper we examine the second-class currents induced by gluon exchange effects in a model with light, unequal quark masses. In the next section, we consider induced second-class current matrix elements taken between free quark states. In Section III, we consider the modifications due to confining both quarks and gluons to the interior of a physical nucleon. Section IV summarizes

our results. In the main body of the text, we shall neglect the Cabibbo admixing of the  $d$  and  $s$  quarks in weak interactions. We shall comment on the induced second-class current effects in  $|\Delta S| = 1$   $\beta$  decays in the summary.

## II. SECOND-CLASS CURRENTS WITH FREE QUARKS

The weak charged current matrix element between  $u$  and  $d$  free quark states (we consider only one color) can be expressed as

$$\begin{aligned} \langle u | j_\beta(0) | d \rangle = & \bar{u} \left[ \gamma_\beta (f_V + f_A \gamma_5) + (q_\beta/2m) (f_S + f_P \gamma_5) \right. \\ & \left. + i \sigma_{\beta\alpha} (q^\alpha/2m) (f_M + f_\Pi \gamma_5) \right] d \end{aligned} \quad (1)$$

where  $u$  and  $d$  are the 4-spinor functions, and  $q = p_u - p_d$  is the momentum transfer. The G-parities of, first and second-class vector (axial vector) currents are, by definition, even (odd) and odd (even), so that in the limit of equal quark masses only  $f_S$  and  $f_\Pi$  are second-class form factors.

In this section, we consider the form factors in a weak interaction theory of the convention type, in which the quarks are coupled to the charged intermediate vector boson field by

$$L_W(x) = g_W \bar{\psi}_U(x) \gamma_\beta (1 + \gamma_5) \psi_D(x) W^\beta(x). \quad (2)$$

The strongest quark interaction is assumed to be with a massless Yang-Mills field (gluon) whose coupling is

$$L_S(x) = g_S \bar{\psi}(x) \lambda_i \gamma_\beta \psi(x) A_i^\beta(x) \quad (3)$$

where  $i$  is the color SU(3) index. The strong coupling constant is assumed to satisfy  $g_S^2/4\pi \cong 0.5$  at a subtraction point characterized by the mass scale  $\mu^2 \leq 1 \text{ GeV}^2$  in accordance with the ideas of asymptotic freedom<sup>10</sup> or the MIT bag model,<sup>9</sup> so that it makes sense to treat  $L_S$  perturbatively. We shall distinguish between  $m_u$  and  $m_d$  but will otherwise ignore electromagnetic interactions.

To first order in  $g_w$  and zero order in  $g_S$ ,  $f_V = f_A = 1$  and the remaining form factors in Eq. 1 vanish. To second order in  $g_S$ , we have the gluon exchange correction shown in Figure 1. The second class form factors generated by this diagram can be calculated without renormalization difficulties by standard methods. After Feynman parameterization and some algebraic manipulation, the integrals for  $f_S$  and  $f_{II}$  can be written as

$$\frac{f_j}{2m} = -\frac{32}{3} i g_S^2 \int dz_1 dz_2 dz_3 \frac{d^4 k}{(2\pi)^4} \delta(1-z_1-z_2-z_3) F_j H^{-3} \quad (4)$$

where

$$j = s \text{ and } II, \text{ and}$$

$$H = k^2 + q^2 z_2 z_3 - m^2 (1-z_1)^2 - 2m\Delta m z_3 (1-z_1) - (\Delta m)^2 z_3 (1-z_1) + i\epsilon$$

$$F_S = 2m \left[ 2z_2 (z_1 - 2) + (1-z_1)^2 + (1-z_1) \right] + 2\Delta m \left[ 2z_2^2 + z_2 (3z_1 - 4) + (1-z_1)^2 + (1-z_1) \right]$$

(cont.)

$$F_{II} = mZ_1(1-Z_1-2Z_2) + \Delta mZ_1(1-Z_1-Z_2)$$

$$m = m_u \quad , \quad m + \Delta m = m_d \quad . \quad (4)$$

To first order in  $\Delta m/2m$  , we find

$$f_s(0)/2m = -(4\alpha_s/9\pi m)(\Delta m/2m) \quad (5a)$$

$$f_{II}(0)/2m = -(4\alpha_s/3\pi m)(\Delta m/2m) \quad (5b)$$

where  $\alpha_s \equiv g_s^2/4\pi$  . Further details and expressions to all orders in  $\Delta m/m$  , together with  $\Delta m \neq 0$  corrections to first class form factors are given in Appendix I.

In addition to  $f_{II}$  , the gluon exchange mechanism therefore generates a nonzero  $f_s$  , which violates the CVC hypothesis. This is a consequence of our assumption  $m_u \neq m_d$  , and is experimentally acceptable, since the present experimental tests of CVC are essentially restricted to the first class current.<sup>7</sup>

We now turn to the analogous nucleon form factors,  $g_s$  and  $g_{II}$  associated with neutron beta decays through the  $n \rightarrow p$  second class current matrix element

$$\langle p | j_\beta^{2nd}(0) | n \rangle = (2M)^{-1} \bar{u}_p \left[ q_\beta g_s + i \sigma_{\beta\alpha} q^\alpha g_{II} \right] u_n \quad (6)$$

where  $u_n$  and  $u_p$  are free nucleon spinors and  $M$  is the nucleon mass. Ignoring quark confinement and gluon exchange effects between different quarks within a nucleon, the nucleon form factors are given by

$$g_S(0) = -\frac{4\alpha_S}{9\pi} \left(\frac{M}{m}\right) \frac{\Delta m}{2m} \quad (7a)$$

$$g_{II}(0) = -\frac{4\alpha_S}{3\pi} \frac{M}{m} \frac{\Delta m}{2m} \quad (7b)$$

If we use quark masses like those suggested by Leutwyler,<sup>8</sup> say  $m_u = 4\text{MeV}$ ,  $m_d = 6\text{ MeV}$ , and if we use  $\alpha_S = 0.5$ , then  $g_{II}(0) = +3g_S(0) = -11$ . Such a value of the tensor form factor,  $g_{II}$ , is comparable to the weak magnetism term as suggested by the experiments of Refs. 1 and 2.

### III. EFFECTS OF CONFINEMENT

In this section we estimate the effects of confining quarks and gluons to a nucleon of finite size. The estimates are made within the framework of the MIT bag model, but we believe they are substantially model independent. The effects are of two types. The first is the generation of a second class current to zero order in  $\alpha_S$ , which owes its existence not only to  $m_u \neq m_d$ , but also to the mixing of upper and lower components in the transition matrix element and reflects the relativistic nature of light quarks. The same phenomenon allows zero mass quarks to produce a non-vanishing nucleon magnetic moment to zero order in  $\alpha_S$ .<sup>11</sup> The second type of confinement effect consists of those modifications of the order  $\alpha_S$  terms of Eq. 7 that result from quark and gluon confinement.

We begin by reminding the reader that the nucleon magnetic moment form factor,  $g_M(0)$ , can be expressed in the Breit frame as

$$i \frac{g_M(0)}{2M} U_f^\dagger \vec{\sigma} U_i (2\pi)^3 \delta^3(0) = \int \langle p_f | \vec{r} \times \vec{V}(\vec{r}, 0) | p_i \rangle d^3r \quad (8)$$

where  $V_\nu$  is the vector current field operator,  $|p_i\rangle$  and  $|p_f\rangle$  are initial and final zero momentum nucleon states, and  $U$  is a two component spinor. The analogous expression for the second class current tensor coefficient,  $g_{II}(0)$ , is

$$i \frac{g_{II}(0)}{2M} U_f^\dagger \vec{\sigma} U_i (2\pi)^3 \delta^3(0) = \int \langle p_f | \vec{r} A_0(\vec{r}, 0) | p_i \rangle d^3r \quad (9)$$

where  $A_\nu$  is the axial vector current operator.

In the rigid sphere approximation to the MIT bag model, the lowest energy single quark wave function is given in the rest frame by<sup>9</sup>

$$q(r) = \frac{N}{\sqrt{4\pi}} \begin{pmatrix} i\frac{\sqrt{\omega+m}}{\omega} j_0(kr)U \\ -\frac{\sqrt{\omega-m}}{\omega} j_1(kr)\vec{\sigma} \cdot \hat{r}U \end{pmatrix} \quad (10)$$

where

$$N^{-2} = j_0(kR)^2 R^3 \left[ 2\omega(\omega R - 1) + m \right] / \omega(\omega - m) \quad (11)$$

$U$  is a 2-spinor, and  $j_0$  and  $j_1$  are spherical Bessel functions.  $R$  is the bag radius, and  $\omega$  is the single quark energy which is related to  $k$  by

$$\omega = \sqrt{k^2 + m^2} \quad (12)$$

The eigenvalue equation for the wave number is

$$\tan kR = kR \left[ 1 - mR - R\sqrt{k^2 + m^2} \right]^{-1} \quad (13)$$

The wave function has been normalized to  $\int q^\dagger q d^3r = 1$ . Thus, to zero order in  $\alpha_s$

$$\begin{aligned} \frac{g_{II}^{(0)}(0)}{2M} &= \int r_i q^\dagger(r) \gamma^0 \gamma_5 q(r) d^3r \\ &= \frac{N^2}{3} \frac{\Delta m}{\omega} \int_0^R r^3 j_0(kr) j_1(kr) \\ &= \frac{\Delta m}{\omega} \frac{N^2}{3} \frac{R^4}{4\beta^3} \left[ \cos 2\beta + 2 - \frac{3}{2\beta} \sin 2\beta \right]. \quad (14) \end{aligned}$$

where we have introduced

$$\beta \equiv kR.$$

In the above, we have ignored the difference between the energy of a quark when it is in a neutron bag as opposed to a proton bag. The second line of Eq. (14) shows the cross term between upper and lower components of the nucleon. For  $m \ll \omega = 400$  MeV,  $R = (200 \text{ MeV})^{-1}$ , we find  $g_{II}(0) \approx \frac{\Delta m}{\omega}$ , so that for  $\Delta m$  equal to a few MeV,  $g_{II}$  is very small. This zero order second class effect is, therefore, not of any practical interest at the present time.

For the calculation of confinement effects on order  $\alpha_s$  terms we can neglect the nonabelian nature and it will be convenient to treat the quark-gluon interaction in the radiation gauge, where we can write the interaction Hamiltonian as<sup>12</sup>

$$\begin{aligned} H_S &= H_S^{(1)} + H_S^{(2)} \\ H_S^{(1)} &= -\sqrt{4\pi\alpha_s} \int \bar{\psi}(x) \vec{\gamma} \lambda_i \psi(x) \cdot \vec{A}_i(x) d^3x \end{aligned} \quad (\text{cont.})$$

$$H_s^{(2)} = \frac{\alpha_s}{2} \frac{\psi^\dagger(\mathbf{x}) \lambda_i \psi(\mathbf{x}) \psi^\dagger(\mathbf{y}) \lambda_i \psi(\mathbf{y})}{|\vec{\mathbf{x}} - \vec{\mathbf{y}}|} d^3x d^3y. \quad (15)$$

The associated old fashioned, time-ordered perturbation diagrams are of two kinds:  $\alpha_s$  modifications of a single quark within the bag, and modifications involving more than one of the three quarks. Representative examples are shown in Figures (2) and (3). Self-energy insertions have been omitted, since we are using the "physical" quark mass. It should also be noted that except for quark self energy insertions,  $H_s^{(2)}$  does not contribute to modification of a single quark line.

We single out Figure (2a) as the lowest energy intermediate state contribution to the single quark vertex correction, when intermediate quarks and gluons are in their lowest modes, and the bag within which they are contained is at rest. This contribution to  $g_{II}(0)$  is given by

$$\begin{aligned} g_{II}^{(2a)}(0) U_f^\dagger \frac{\vec{\sigma}}{2M} U_i = & \frac{16}{3} g_s^2 \int \vec{y} \bar{q}_u(z) \gamma^i q_{u\lambda}(z) \\ & \times \bar{q}_{u\lambda}(y) \gamma^0 \gamma_5 q_{d\rho}(y) \bar{q}_{d\rho}(x) \gamma^j q(x) G_m^i(z) G_m^{j*}(x) \\ & \times k_0^{-1} (k_0 + E_w)^{-1} d^3x d^3y d^3z \end{aligned} \quad (16)$$

where  $E_w$  is the W-boson energy and  $k_0$  is the energy of the lowest gluon mode in the cavity. The gluon wave functions are<sup>13</sup>

$$\vec{G}_m(\vec{r}) \propto \vec{r} Y_{1m}(\theta, \phi) j_1(kr)$$

(cont.)

$$= N_m \left[ \hat{\theta} i m - \hat{\phi} (|m| \cos \theta - \delta_{m0} \sin \theta) \right] j_1(kr) e^{im\phi}$$

and

$$N_m^{-2} = 2k_0 \frac{4\pi}{3} (1 + |m|) \left[ j_1(kR)^2 - j_0(kR)j_2(kR) \right] R^3 \quad (17)$$

so as to normalize to

$$i \int G_m^j(r, t) \overleftrightarrow{\frac{\partial}{\partial t}} G_n^j(r, t) d^3r = \delta_{mn} \quad \text{where: } G_m^j(r, t) = G_m^j(r) e^{-ik_0 t} .$$

Here the smallest wave number is  $k \equiv k_0 = 2.73/R$ . Since the  $\vec{y}$  dependent factors in Eq. (16) are identical to those appearing in the evaluation of  $g_{II}^{(0)}(0)$  (see Eq. (14)), we may write

$$g_{II}^{(0)} U_f^{\dagger \sigma} U_i = \frac{16}{3} g_{II}^{(0)}(0) g_s^2 \int \bar{q}_u(z) \gamma^j q_{u\lambda}(z) U_\lambda^{\dagger \sigma} U_\rho \bar{q}_{d\rho}(x) \gamma^k q_d(x) \times G_m^j(z) G_m^{k*}(x) k_0^{-1} (k_0 + E_w)^{-1} d^3x d^3z \quad (18)$$

We can make an order of magnitude estimate of the integral by noting that the quarks are highly relativistic, so that  $\int \bar{q} \gamma q d^3x \sim 1$ . The normalization factor for  $\vec{G}_m$  implies  $|\vec{G}_m| \sim (4\pi k_0 R^3)^{-1}$ . Thus,

$$g_{II}^{(2a)}(0) \sim g_{II}^{(0)}(0) \frac{5\alpha_s}{(k_0 R)^3} \sim g_{II}^{(0)}(0) \frac{\alpha_s}{4} \quad (19)$$

This estimate is certainly within an order of magnitude, and provided that the  $\alpha_s$  expansion of  $g_{II}^{(0)}$  converges, makes

it clear that setting  $g_{II}(0) = g_{II}^{(0)}(0)$  does not give a misleading picture of confinement effects. To strengthen this conclusion, we note that a factor of  $g_{II}^{(0)}(0)$  will arise from the  $q\bar{q}W$  vertex in every diagram, including those in Fig. 3, in which only ground state quarks are included; inclusion of higher modes of excitation will produce a similar factor at that vertex proportional to  $\Delta m$ . A similar reduction of the scalar form factor  $g_S(0)$ , will obviously also result from confinement.

In order to convince the reader (and ourselves) that this result is essentially model-independent, we attempt to modify the free quark vertex correction of Eq. (4) without the introduction of explicit bound states. Examination of the  $k$  integration in Eq. (4) shows that the major contribution comes from small Euclidean  $k^2$  values. But confinement of the gluons to a finite spatial region implies that Euclidean  $k^2$  can be no smaller than the square of the lowest gluon bound state energy,  $k_0^2$ . We have evaluated Eq. (4) for  $f_{II}(0)$  with this lower limit for  $k^2$ . The modification to Eq. (5b) is essentially to replace  $m$  by  $k_0$ . Since gluons and light quarks confined to a radius  $R$  will each have an energy of order  $R^{-1}$ , the numerical result obtained in this way is about the same as that of the  $\alpha_s$  bag model contribution. More specifically we find

$$g_{II}^{(\alpha_s)}(0) = + \frac{\alpha_s}{9\pi} \frac{M\Delta m}{k_0^2} .$$

Translated into coordinate space, this estimate recognizes that the radius of the nucleon,  $R$ , places an upper limit on the wavelength of gluons that can communicate with quarks confined to this region.

#### IV. CONCLUSION

We have shown that light quark models of the nucleon in which the  $u$ - $d$  quark mass difference is taken into account predict second class form factors of order  $M\Delta m/m^2$  ( $M$ = nucleon,  $m$ =quark mass), if confinement is totally ignored. The first effect of confinement is to replace the quark mass by the much larger single quark energy, which reduces the second class form factors to values well below those of the first class form factors. The lesson to be learned is that free quark results are totally misleading when the quarks are, in fact, confined and highly relativistic. Finally, we note that in  $|\Delta s| = 1$  baryon  $\beta$ -decays the relevant  $\Delta m$  is  $m_s - m_{u,d}$ , so that here  $\Delta m/\omega$  is of order unity. We expect  $g_{II}(|\Delta s|=1)/g_V \sim g_S(|\Delta s|=1)/g_V \sim 1$  as a consequence of a mechanism analogous to that described here.

#### ACKNOWLEDGEMENTS

The authors would like to thank K. Johnson and J. Kiskis for helpful communication, and one of us (A.H.) would like to thank the Fermilab for hospitality during periodic visits. One of the authors (BWL) acknowledges the crucial comment of

S. L. Glashow, which prompted our investigation of this problem.

Another author (P.S.) appreciated discussions with H. Kluberg-Stern and R. Shrock, and the kind hospitality at the Fermilab theoretical physics department.

APPENDIX

In this appendix we give details associated with the free quark vertex correction of Eq. (4). As in the analogous vertex correction in QED,<sup>14</sup> the vertex diagram can be written as

$$\begin{aligned}
 \Lambda_\mu(p, p') &= (-ig_s)^2 \int \frac{d^4k}{(2\pi)^4} \frac{-i}{k^2 - \lambda^2 + i\epsilon} \gamma_\nu \frac{i}{\not{p}' - \not{k} - m + i\epsilon} \\
 &\quad \times \gamma_\mu (1 + \gamma_5) \frac{i}{\not{p} - \not{k} + m + \Delta m + i\epsilon} \gamma^\nu \\
 &= -2ig_s^2 \int d^4k dz_i \delta(1 - z_1 - z_2 - z_3) \gamma_\nu (\not{p}' - \not{k} + m) \\
 &\quad \times \gamma_\mu (1 + \gamma_5) (\not{p} - \not{k} + m + \Delta m) \gamma^\nu \left[ k^2 - \lambda^2 z_1 \right. \\
 &\quad \left. + q^2 z_2 z_3 - m^2 (1 - z_1)^2 - 2m\Delta m z_3 (1 - z_1) \right. \\
 &\quad \left. - (\Delta m)^2 z_3 (1 - z_1) + i\epsilon \right]^{-3} .
 \end{aligned} \tag{A1}$$

$\lambda$  is the gluon mass to handle the infrared divergence. Separating off the logarithmic infinity, we have

$$\begin{aligned}
 \Lambda_\mu(p', p) &= \frac{\alpha_s}{2\pi} \int dz_i \delta(1 - z_1 - z_2 - z_3) \left[ \gamma_\mu (1 + \gamma_5) \right. \\
 &\quad \left. \times \left( -\frac{3}{2} + \ln \frac{D_0}{D} + \ln \frac{D_0}{D} \right)^{-\frac{1}{2}} \frac{N_\mu}{D} \right]
 \end{aligned} \tag{A2}$$

where

$$\begin{aligned}
 D_0 &= m^2 (1 - z_1)^2 + \lambda^2 z_1 + \Delta m (2m + \Delta m) z_3 (1 - z_1) , \\
 D &= D_0 - q^2 z_2 z_3 ,
 \end{aligned} \tag{A4}$$

$$N_{\mu} = \gamma_{\nu} \left[ \not{\epsilon} (1-z_2) - \not{\epsilon} z_3 + m \right] \gamma_{\mu} (1+\gamma_5) \left[ \not{\epsilon} (1-z_3) - \not{\epsilon} z_2 + m + \Delta m \right] \gamma^{\nu} \quad (A4)$$

and  $\mathcal{M}$  is a high energy cut-off on  $k^2$ . Since  $\Lambda_{\mu}$  is to be multiplied by free quark spinors,  $N_{\mu}$  can be written as

$$\begin{aligned} N_{\mu} = & \gamma_{\mu} \gamma_5 \left[ 2(1-z_2)(1-z_3)(-q^2 + m^2 + m\Delta m + (\Delta m)^2) \right. \\ & - 2(m+\Delta m)^2(1-z_3)z_3 - 2(1-z_2)z_2m^2 \\ & \left. + 2z_2z_3m(m+\Delta m) + 2m(m+\Delta m) \right] \\ & + 4p_{\mu} \gamma_5 \left[ m z_2(1-z_3) + (m+\Delta m)z_3^2 \right] \\ & - 4p_{\mu} \not{\epsilon} \gamma_5 \left[ (m+\Delta m)z_3(1-z_2) + m z_2^2 \right] \\ & + \gamma_{\mu} \left[ 2(1-z_2)(1-z_3)(-q^2 + 3m^2 + 3m\Delta m + (\Delta m)^2) \right. \\ & \left. - 2(m+\Delta m)^2(1-z_3)z_3 - 2z_2(1-z_2)m^2 - 2z_2z_3m(m+\Delta m) - 2m(m+\Delta m) \right] \\ & + 4p_{\mu} \left[ z_2(1-z_3)m - z_3^2(m+\Delta m) \right] \\ & + 4p_{\mu} \not{\epsilon} \left[ z_3(1-z_2)(m+\Delta m) - z_2^2m \right]. \quad (A5) \end{aligned}$$

The Gordon identity and its axial counter part are (free quark spin factors understood).

$$\frac{1}{2} \left[ (\not{\epsilon}' - \not{\epsilon}), \gamma_{\mu} \right] = -(p_{\mu}' + p_{\mu}) + (2m + \Delta m) \gamma_{\mu} \quad (A6)$$

$$\frac{1}{2} \left[ (\not{\epsilon}' - \not{\epsilon}), \gamma_{\mu} \right] \gamma_5 = -(p_{\mu}' + p_{\mu}) \gamma_5 - \Delta m \gamma_{\mu} \gamma_5 \quad (A7)$$

$N_{\mu}$  can therefore be rewritten as

$$\begin{aligned}
N_\mu = & \gamma_\mu \left[ -2(1-z_2)(1-z_3)q^2 + 2(2-(1-z_1))^2 \right. \\
& \left. - 2(1-z_1)(m+\Delta m)m + 2z_1\Delta m^2 \right] \\
& + \gamma_\mu \gamma_5 \left[ -2(1-z_2)(1-z_3)q^2 + \right. \\
& + 2((1-z_3)z_1 + 1 + z_2z_3 - z_2(1-z_2))(m+\Delta m)m \\
& \left. + 2((1-z_3)z_1 + z_1z_2)\Delta m^2 \right] \\
& + \left[ \not{q}, \gamma_\mu \right] \left[ -z_1(1-z_1)m - z_1(1-z_1-z_2)\Delta m \right] \\
& - 2q_\mu \gamma_5 \left[ (4z_2^2 - 4z_2(1-z_1) + (1-z_1)^2 + (1-z_1))m \right. \\
& \left. + (2z_2^2 - z_2(4-3z_1) + (1-z_1)^2 + (1-z_1))\Delta m \right] \\
& + 2q_\mu \left[ (2z_2(z_1-2) + (1-z_1)^2 + (1-z_1))m \right. \\
& \left. + (2z_2^2 + z_2(3z_1-4) + (1-z_1)^2 + (1-z_1))\Delta m \right] \\
& + \left[ \not{q}, \gamma_\mu \right] \gamma_5 \left[ z_1(1-z_1-z_2)\Delta m + z_1(1-2z_2-z_1)m \right] . \tag{A8}
\end{aligned}$$

The renormalized vertex is obtained by performing one subtraction at  $q^2=0$ . To first order in  $q$ , we therefore, obtain after integrating over the  $z_i$

$$\begin{aligned}
\Lambda_\mu^{\text{Ren}} = & q_\mu \gamma_5 \left( -\frac{\alpha_s}{4\pi} \right) \left[ \frac{1}{\Delta m} - \frac{1}{(2m+\Delta m)} \left( 3 - 2\left(\frac{m}{\Delta m}\right)^2 \right) \right. \\
& \left. - \frac{2(m+\Delta m)}{(2m+\Delta m)^2 (\Delta m)^3} \left( 3m(\Delta m)^2 + 2m^2(m+\Delta m) \right) \right. \\
& \left. \times \ln \left( 1 + \frac{\Delta m}{m} \right) \right]
\end{aligned}$$

$$\begin{aligned}
 & + \left[ \not{\epsilon}, \gamma_{\mu} \right] \frac{\alpha_s}{4\pi} \left[ \frac{1}{2(2m+\Delta m)} + \frac{m(m+\Delta m)}{\Delta m(2m+\Delta m)^2} \ln\left(1 + \frac{\Delta m}{m}\right) \right] \\
 & + q_{\mu} \frac{\alpha_s}{4\pi} \left[ \frac{1}{2m+\Delta m} - \frac{2}{(2m+\Delta m)^2 \Delta m} (5m^2 + 6m\Delta m + \frac{3}{2} (\Delta m)^2) \right. \\
 & + \frac{4(m+\Delta m)m}{(2m+\Delta m)^3 (\Delta m)^2} (5m^2 + 5m\Delta m + \frac{3}{2} (\Delta m)^2) \\
 & \left. \ln\left(1 + \frac{\Delta m}{m}\right) \right] \\
 & + \left[ \not{\epsilon}, \gamma_{\mu} \right] \gamma_5 \left( -\frac{\alpha}{4\pi} \right) \left[ \frac{1}{2\Delta m} - \frac{m}{(\Delta m)^2} \frac{1 + \frac{\Delta m}{m}}{2 + \frac{\Delta m}{m}} \ln\left(1 + \frac{\Delta m}{m}\right) \right]. \tag{A9}
 \end{aligned}$$

To first order in  $\Delta m/m$ ,

$$\begin{aligned}
 \Lambda_{\mu}^{\text{Ren}} = & \frac{\alpha_s}{4\pi m} \left\{ \frac{7}{3} \left(1 - \frac{\Delta m}{2m}\right) q_{\mu} \gamma_5 + \frac{1}{2} \left(1 - \frac{\Delta m}{2m}\right) \left[ \not{\epsilon}, \gamma_{\mu} \right] \right. \\
 & \left. - \frac{1}{3} \frac{\Delta m}{2m} q_{\mu} - \frac{1}{2} \frac{\Delta m}{2m} \left[ \not{\epsilon}, \gamma_{\mu} \right] \gamma_5 \right\}. \tag{A10}
 \end{aligned}$$

REFERENCES AND FOOTNOTES

- <sup>1</sup>K. Sugamoto, I. Tanihata, and J. Göring, Phys. Rev. Lett. 34, 1533 (1975).
- <sup>2</sup>F.P. Calaprice, S.J. Freedman, W.C. Mead, and H.C. Vantine, Phys. Rev. Lett. 35, 1566 (1975).
- <sup>3</sup>A.M. Nathan, G.T. Garvey, P. Paul, and E.K. Warburton, Phys. Rev. Lett. 35, 1137 (1975).
- <sup>4</sup>D. Wilkinson and D. Alburger, Phys. Rev. Lett. 26, 1127 (1971).
- <sup>5</sup>R. Tribble and G.T. Garvey, Phys. Rev. Lett. 32, 314 (1974); Phys. Rev. C12, 967 (1975).
- <sup>6</sup>A further complication may come from reliance upon the validity of CVC for the first-class contribution, the basis for which we understand has been weakened by the discovery of an uncertainty in the weak magnetism experiment of Y.K. Lee, L.W. Mo, and C.S. Wu, Phys. Rev. Lett. 10, 253 (1963).
- <sup>7</sup>B. Holstein and S.B. Treiman, "Second Class Currents", Princeton University Preprint (1975). In this paper, an interesting possibility of detecting second-class current effects in weak pion production is studied. See also M.A.B. Beg and J. Bernstein, Phys. Rev. D5, 714 (1972).
- <sup>8</sup>J. Gasser and H. Leutwyler N.P. B94, 269 (1975). H. Leutwyler, N.P. B76, 413 (1974). Note that the values of  $m_u$  and  $m_d$  are in perfect agreement with the relation:

$$\frac{m_{\bar{d}} - m_u}{m_{\bar{d}} + m_u} = \frac{m_{k^0}^2 - m_{k^+}^2}{m_{\pi}^2} \approx 0.2 \quad ,$$

which can be deduced from M. Gell-Mann, R.J. Oakes and B. Renner, Phys. Rev. 175, 2195 (1968).

- <sup>9</sup>T. de Grand, R.L. Jaffe, K. Johnson, and J. Kiskis, Phys. Rev. D, 2060 (1975).
- <sup>10</sup>H.D. Politzer, Phys. Rev. Lett. 30, 1346 (1973); D.G. Gross and F. Wilczek, Phys. Rev. Lett. 30, 1343 (1973); T. Appelquist and H.D. Politzer, Phys. Rev. Lett. 34, 43 (1975).
- <sup>11</sup>A. Chodos, R.L. Jaffe, K. Johnson and C.B. Thorn, Phys. Rev. D10, 2599 (1974).
- <sup>12</sup>Drell and Bjorken, "Relativistic Quantum Fields", McGraw-Hill, New York, (1965). For the analogous calculation of the electron magnetic moment in QED, see J.M. Luttinger, Phys. Rev. 74, 893 (1948).
- <sup>13</sup>The gluon wavefunction is the TM mode with the boundary condition  $\frac{d}{dR} [Rj_{\ell}(kR)] = 0$ . See Morse and Feschbach, Methods of Mathematical Physics, p. 1870.
- <sup>14</sup>We follow the procedures of Drell and Bjorken, "Relativistic Quantum Mechanics" McGraw-Hill, New York, (1964), p.166.

FIGURE CAPTIONS

- Fig. 1 Feynman diagram for the gluon exchange correction to  $d \rightarrow u + w^-$ . The wavy line is a gluon.
- Fig. 2: Single quark modifications to order  $\alpha_s$ .
- Fig. 3: Modifications involving quark-quark correlations to order  $\alpha_s$ . The blob in 3b is the instantaneous Coulomb interaction,  $H_S^{(2)}$ .




Fig. 1




Fig. 2




Fig. 3

$$\frac{m_{\bar{d}} - m_u}{m_{\bar{d}} + m_u} = \frac{m_{k^0}^2 - m_{k^+}^2}{m_{\pi}^2} \approx 0.2 \quad ,$$

which can be deduced from M. Gell-Mann, R.J. Oakes and B. Renner, Phys. Rev. 175, 2195 (1968).

- <sup>9</sup>T. de Grand, R.L. Jaffe, K. Johnson, and J. Kiskis, Phys. Rev. D, 2060 (1975).
- <sup>10</sup>H.D. Politzer, Phys. Rev. Lett. 30, 1346 (1973); D.G. Gross and F. Wilczek, Phys. Rev. Lett. 30, 1343 (1973); T. Appelquist and H.D. Politzer, Phys. Rev. Lett. 34, 43 (1975).
- <sup>11</sup>A. Chodos, R.L. Jaffe, K. Johnson and C.B. Thorn, Phys. Rev. D10, 2599 (1974).
- <sup>12</sup>Drell and Bjorken, "Relativistic Quantum Fields", McGraw-Hill, New York, (1965). For the analogous calculation of the electron magnetic moment in QED, see J.M. Luttinger, Phys. Rev. 74, 893 (1948).
- <sup>13</sup>The gluon wavefunction is the TM mode with the boundary condition  $\frac{d}{dR} [Rj_{\ell}(kR)] = 0$ . See Morse and Feschbach, Methods of Mathematical Physics, p. 1870.
- <sup>14</sup>We follow the procedures of Drell and Bjorken, "Relativistic Quantum Mechanics" McGraw-Hill, New York, (1964), p.166.

FIGURE CAPTIONS

- Fig. 1 Feynman diagram for the gluon exchange correction to  $d \rightarrow u + w^-$ . The wavy line is a gluon.
- Fig. 2: Single quark modifications to order  $\alpha_s$ .
- Fig. 3: Modifications involving quark-quark correlations to order  $\alpha_s$ . The blob in 3b is the instantaneous Coulomb interaction,  $H_S^{(2)}$ .




Fig. 1




Fig. 2




Fig. 3