FERMILAB-PUB-85-190-E
COLO-HEP-104
September 1985
E516

A Study of the Inclusive $K_S^0, \Lambda, \bar{\Lambda}$ Production in Diffractive γP Interactions

S. Bhadra, J.A. Appel, D.F. Bartlett, B.H. Denby, J. Biel, D. Bintinger, V.K. Bharadwaj, D.E. Blodgett, S.B. Bracker, J. Bronstein, C. Daum, A.L. Duncan, A.M. Eisner, J.R. Elliott, P. Estabrooks, G.F. Hartner, G.R. Kalbfleisch, R.G. Kennett, B.R. Kumar, M.J. Losty, A. Lu, G.J. Luste, P.M. Mantsch, J.F. Martin, R.J. Morrison, T. Nash, U. Nauenberg, J. Pinfold, M. Robertson, W. Schmidke, K.K. Shahbazian, R.A. Sheperd, K. Sliwa, M.D. Sokoloff, W.J. Spalding, J. Stacey, K.C. Stanfield, M. Streetman, D.J. Summers, S.E. Willis, M.S. Witherell, S.J. Yellin, C.J. Zorn

The University of Colorado, Boulder, Colorado, 80309, the University of California at Santa Barbara, Santa Barbara, California, 93106, Carleton University, Ottawa, K1S 5B6, Canada, the Fermi National Accelerator Laboratory, Batavia, Illinois, 60510, the National Research Council of Canada, Ottawa, Ontario, K1A 0R6, Canada, the University of Oklahoma, Norman, Oklahoma, 73019, and The University of Toronto, Toronto, M5S 1A7, Canada

To be Submitted to Physical Review Letters.

ABSTRACT

We have studied inclusive K_s,Λ, and $\bar{\Lambda}$ photoproduction over the ranges 40 $< E_\gamma <$ 170 GeV and forward produced mass $2 < M_F < 10$ GeV. We observe equal Λ and $\bar{\Lambda}$ production rates and spectra as expected in a diffractive process where the target proton remains unaltered. We show that the fraction of hadronic events with a strange particle produced in events with forward mass M_F agrees well with the same measurement in e^+e^- annihilation at a center of mass energy $E_{cm} = M_F$. The $X = 2P/M_F$ distributions of these three particles in the forward mass rest frame are compared with theoretical predictions.

PACS 13.60-r, 13.60.Hb, 13.60.Rj

The inclusive production of K_s, Λ and $\bar{\Lambda}$ in photon initiated reactions and in e^+e^- annihilation can be used to compare the dynamics of strange particle production in the two reactions. Any observed differences may indicate variations in the structure of the initial states or in the hadronization process in each reaction. In this experiment we show that the fraction of K_s, Λ or $\bar{\Lambda}$ produced in diffractive photoproduction is very similar to that in e^+e^- annihilation. This comparison, which has never been possible before, is made by comparing our measured photoproduction rates at a given forward mass, M_F, with the existing e^+e^- data at a center of mass energy $E_{cm} = M_F$.

Our measurement of $\gamma + p \rightarrow K_s,(\Lambda),(\bar{\Lambda}) + X + p_{recoil}$ used the Fermilab Tagged Photon Spectrometer (TPS). This detector consisted of a large acceptance forward spectrometer and a highly segmented recoil detector with both tracking and dE/dx capabilities.[1,2] In addition, a tagging system gave a good measurement of the incident photon energy. The photon beam was generated by a 170 GeV electron beam impinging on a 0.2 radiation length thick Cu radiator. The photons then interacted in a 1.5-m liquid hydrogen target. The error in the photon energy was ~5%.

In this experiment the forward mass was calculated as a missing mass using the measurements of the recoil proton. A trigger processor[3] was used to select rapidly events with single recoiling protons, was able to calculate the produced forward mass, and allowed us to record only events with a forward mass greater than 2 GeV. The error in the forward mass averaged 2.5%. In addition, the recoil detector determined not only that the recoil was a proton, but also that it was not associated with other particles present in the recoil detector. This allowed us to know whether the recoiling proton was the original target or whether it resulted from a recoiling N* decay.

The latter protons would give rise to a calculated forward (or missing) mass larger than the actual value.

The data presented in this report consists of a sample in which there was only one charged particle in the recoil detector and it was identified as a proton. Hence, we are studying particle production via the process generally described as "diffractive."

We made a detailed study of how often the forward mass was miscalculated because the proton was really a byproduct of a N* decay where the associated pions were not observed in the recoil detector. This study was done with events in which the forward system of particles was fully reconstructed by the spectrometer.[4] The conclusion was that the average fraction of incorrectly determined mass values was about 8%. The maximum fraction, which occurred for $6 < M_F < 8$ GeV was 12%.

The numbers of $K_S \rightarrow \pi^+\pi^-$, $\Lambda \rightarrow p\pi^-$, and $\bar{\Lambda} \rightarrow \bar{p}\pi^+$ events were determined from the size of the clear peak above background seen in the diparticle mass spectra. The background was effectively reduced by cuts in the distance of closest approach (<1 cm) between the charged tracks and by requiring that the Cerenkov counter within the spectrometer identify the tracks as pions or protons in the appropriate cases. While the background was reduced by a factor of 3 for K_S and 9 for $\Lambda,\bar{\Lambda}$ decays, the loss of signal was no more than 15%. These factors were reproduced by our Monte Carlo program which included a detector simulation. The number of events observed is shown in Table 1. These totals were corrected for decay branching ratios, losses due to detector inefficiencies, and decays outside the active region of the detector.

The photoproduction data was analyzed as a function of both the center of mass energy and the forward mass. In fig. 1 we show the number of $K_S,\Lambda,\bar{\Lambda}$ per hadronic event as a function of the forward mass for various incident photon

energies. The data indicate that the production of these particles does not depend on the incident energy for a given forward mass, while it does depend on the forward mass for a given photon energy. Hence we conclude that the forward mass is the more relevant variable in this problem. This is similar to the behavior in hadronic spectra[5].

Hadronic reactions have not generally been analyzed in terms of the forward mass, but in terms of the available energy ($E_{avail} = E_{c.m.} - \sum M_{initial}$). We show, for completeness, in fig. 2, a comparison of K_s production for various comparable photon[6] and hadronic reactions[7] in terms of this variable. Our data points, because we only triggered on $M_F > 2$ GeV, have been corrected for that part of the K_s and hadronic cross section with $M_F < 2$ GeV.[8] This leads to a reduction in the fraction of K_s by the factor .70 ± .07. The similarity between photon induced and pion induced reactions is clear, as is the discrepancy between these and e^+e^- annihilation[9] when plotted in this variable. A similar comparison in the case of Λ and $\bar{\Lambda}$ production yields the same conclusions.

In. fig. 3 we compare our results on K_s, Λ and $\bar{\Lambda}$ production as a function of the forward mass M_F with those in e^+e^- annihilation[9] at the center of mass energy $E_{cm} = M_F$. The Λ, $\bar{\Lambda}$ data is averaged since they are equal. The agreement between these two reactions is striking. The only discrepancy is at high mass where the production in e^+e^- annihilation may be slightly larger than in photoproduction. Hence we are lead to the main conclusion of this study: the photon initiated diffractive production of strange particles in mass systems > 2 GeV cannot be distinguished from that of an e^+e^- annihilation state with the same mass.

We have also measured the distribution in Feynman $x = 2P/M_F$ of the K_s, Λ and $\bar{\Lambda}$. The data for $x_F d\sigma/dx_F$ was fitted to the form $A\,(1 - x)^{\alpha}$. The

values of α which we obtained were $\alpha = 1.8 \pm .4$ for K_s and $3.04 \pm .87$ for $\Lambda + \bar{\Lambda}$. These results are in marginal agreement with predictions of photon fragmentation models ($\alpha = 1$ for K_s, $\alpha = 2$ for $\Lambda,\bar{\Lambda}$).[10] Finally, we have also observed Ξ^-, $\bar{\Xi}^-$ production. The ratio of Ξ to Λ production is $.07 \pm 0.015$. This result is in good agreement with the same ratio in e^+e^- annihilation.[11]

We would like to thank the Fermilab staff for their support of the TPS effort. This work was supported by grants from the U.S. Department of Energy, the Canadian National Research Council, and the Canadian National Science and Engineering Research Council.

References

1. The spectrometer is well described in the theses of A. Duncan (U. Colorado), B. Denby (UC. Santa Barbara), D. Summers (UC. Santa Barbara), S. Bhadra (U. Colorado) (unpublished).
2. G. Hartner et al., NIM 216, 13, (1983)
3. J. F. Martin et al., Proceedings of the Conference on Applications of Microprocessors to H.E.P. CERN 81-07, 164, (1981). E. Barsotti et al., IEEE Trans. Nuc. Sci., N5-26, 686, (1979)
4. S. Bhadra thesis (U. Colorado) (unpublished)
5. Rajendran Raja Phys. Rev. D18, 204, (1978)
6. K. Abe et al., Phys. Rev. D29, 1877, (1984)
7. M. Alston-Garnjost et al. Phys. Rev. Letters 35, 142, (1975)
   P. H. Stuntebeck et al., Phys. Rev. D9, 608, (1974)
   P. Bosetti et al., Nuclear Phys. B94, 21, (1975)
   D. Ljung et al., Phys. Rev. D15, 3163, (1977)
   D. Bogert et al., Phys. Rev. D16, 2098, (1977)
   F. Barreiro et al., Phys. Rev. D17, 669, (1978)
   D. Brick et al., Nuclear Phys. B164, 1, (1980), Phys. Rev. D20, 2123, (1979).
   I.V. Aijenko, et al. Nuclear Phys. B165, 1, (1980)
   N. N. Biswas et al., Nuclear Phys. B167, 41, (1980)
8. T. Chapin et al., Phys. Rev. D31, 17, (1985)
9. V. Luth et al. (MARK I Coll.), Phys. Letters 70B, 120, (1977)
   G. S. Abrams et al. (MARK I Coll.), Phys. Rev. Letters 44, 10, (1980)
   R. Brandelik et al. (TASSO Coll.), Phys. Letters 94B, 91, (1980)
   C. H. Berger et. al. (PLUTO Coll.), Phys. Letters 104B, 79, (1981)
   R. Brandelik et al. (TASSO Coll.), Phys. Letters 105B, 75, (1981)

M. S. Alam et al. (CLEO Coll.), CLNS/82/547, CLEO 82-06 (1982)

W. Bartel et al. (JADE Coll.), Zeitschrift fur Physik C20, 187, (1983)

H. Aihara et al. (TPC Coll.), Phys. Rev. Letters 53, 2378, 1984; 54, 274, (1985)

10. R. Blankenbecler and S. Brodsky, Phys. Rev. D10, 2973, (1974)

T. A. DeGrand and J. Randa, Phys. Letters 110B, 484, (1982)

11. M. Althoff et al. (TASSO Coll.) Phys. Letters 130B, 340, (1983). Note that this result is at higher $E_{c.m.}$ than our range in forward mass.

TABLE 1

Mass (GeV)→	2 - 4	4 - 6	6 - 8	8 - 10
Hadronic Events	371198	388683	368202	209248
$K_S \to \pi^+\pi^-$	8192 ± 229	10814 ± 344	10309 ± 337	5736 ± 269
$\Lambda \to p\ \pi^-$	584 ± 56	1399 ± 115	1492 ± 118	1032 ± 97
$\bar{\Lambda} \to \bar{p}\ \pi^+$	505 ± 154	1279 ± 124	1509 ± 132	923 ± 111

FIGURE CAPTIONS

Fig. 1. Fraction of Hadronic Events with a $K_s, \Lambda, \bar{\Lambda}$ in the final state as a function of the forward mass for various incident photon energies.

Fig. 2. Fraction of Hadronic Events with a K_s in the final state as a function of the available energy ($E_{avail} = E_{c.m.} - \sum M_{initial}$). The comparison data is from ref. 6,7 and 9. The errors on the data points of this experiment do not include an overall normalization error of ± 10% due to the correction for low mass events.

Fig. 3. Fraction of Hadronic Events with a $K_s, \Lambda, \bar{\Lambda}$ in the final state as a function of the forward mass for this experiment and as a function of $E_{c.m.}$ for e^+e^- annihilation. The comparison data is from ref. 9.


$\bar{\Lambda}$ Λ, K_S
40 < E < 70 GeV
70 < E < 100 GeV
100 < E_γ < 170 GeV
K_S
Λ, $\bar{\Lambda}$
NUMBER OF EVENTS / NUMBER OF HADRONIC EVENTS
0.1
0.01
0.001
2
4
6
8
10
12
M_F (GeV)


○ γp → K°s X Ref. 7
□ π± p → Ks X Ref. 7
× πd → Ks X Ref. 6
● γp → Ks X (p) THIS EXP
■ e+e− → Ks X Ref. 9
NUMBER OF Ks EVENTS / NUMBER OF HADRONIC EVENTS
10
1.0
0.1
0.01
4
8
12
16
20
24
28
32
36
E AVAIL (GeV)


□ e+ e− → KS X
● γp → KS X(p) THIS EXP.
□ e+ e− → Λ, Λ̄ X
● γp → Λ, Λ̄ X(p) THIS EXP. (AVE.)
NUMBER OF EVENTS / NUMBER OF HADRONIC EVENTS
0.1
0.01
2
4
6
8
10
12
14
16
MF, √S (GeV)