

Fermi National Accelerator Laboratory

FERMILAB-Pub-87/155-A
October 1987

On the Magnitude of Baryon-to-Photon Ratio Inhomogeneities Resulting From a First Order Quark/Hadron Transition

Michael S. Turner

NASA/Fermilab Astrophysics Center
Fermi National Accelerator Laboratory
Batavia, IL 60510-0500

and

Departments of Physics and Astronomy and Astrophysics
Enrico Fermi Institute
The University of Chicago
Chicago, IL 60637

Abstract. We show that consideration of the low lying baryon states in addition to the neutron and proton (specifically, the Λ , Σ , and Δ states) reduces the ratio of baryon number density in the quark phase to that in the hadronic phase by more than a factor of 2. This implies that inhomogeneities in the local baryon number to photon ratio produced during the quark/hadron transition are likely to be smaller than previous estimates, and therefore unless the transition temperature is less than ~ 150 MeV, the effects upon primordial nucleosynthesis will not be significant.

Submitted to The Physical Review D.



It has been pointed out that if the quark/hadron transition¹ is strongly first order and if the transition temperature is low enough, then large local fluctuations in the baryon-to-photon ratio can arise², and might significantly modify the predictions of the standard scenario for primordial nucleosynthesis³⁻⁵. [For a review of the standard scenario of nucleosynthesis see Refs. 6.] Among other things, the potential effect of the quark/hadron transition upon primordial nucleosynthesis depends upon the magnitude of the fluctuations in the baryon-to-photon ratio, η . The size of the fluctuations is estimated by computing the ratio ($\equiv R$) of the net baryon number density in the quark phase to that in the hadron phase at the critical temperature T_c , assuming thermal and chemical equilibrium between the quark and hadron phases⁷. In so doing, the only baryonic states in the hadronic phase that have been taken into account are the neutron and proton. Here we point out that the predicted value of this ratio R decreases significantly when other low lying baryon states are included: specifically, the $J^P = 3/2^+$ Δ^{++} , Δ^+ , Δ^0 , Δ^- states and the $J^P = 1/2^+$, strangeness - 1 states, Λ , Σ^+ , Σ^0 , Σ^- .

The net baryon number density in a species i with internal degrees of freedom g_i and baryon number ± 1 which is very non-relativistic and in thermal equilibrium with chemical potential μ_i is:

$$n_{NR}^B = 2g_i(m_i T/2\pi)^{3/2} \exp(-m_i/T) \sinh(\mu_i/T) \quad (1a)$$

where T is the temperature and m_i is the mass of species i .

The net baryon number density associated with a highly-relativistic species with internal degrees of freedom g_i and baryon number $\pm \frac{1}{3}$ in thermal equilibrium is:

$$n_R^B = (g_i T^3/6\pi^2)[4a(\mu_i/T) + \frac{1}{3}(\mu_i/T)^3], \quad (1b)$$

where $a = 1 - \frac{1}{2^3} + \frac{1}{3^3} - \frac{1}{4^3} + \dots \simeq 0.82246$.

Numerical studies indicate that the transition temperature for the quark/hadron (i.e., deconfinement/confinement) transition is probably in the range¹: $T_c \sim 2\Lambda_{\overline{MS}} \simeq 100 - 400$ MeV. [The quantity $\Lambda_{\overline{MS}}$ is the QCD renormalization scale computed in the modified, minimal subtraction scheme⁸; recent determinations indicate that $\Lambda_{\overline{MS}} \simeq 100$ MeV-400 MeV (Ref. 9).] In the quark phase, the u and d quarks (masses $\lesssim 10$ MeV) are most certainly relativistic, while it is uncertain whether or not the s quark (mass ~ 150 -300 MeV) is. In the hadronic phase, all baryonic states are more massive than ~ 940 MeV and so are non-relativistic. Making the assumption of thermal and chemical equilibrium between the quark and hadronic phases,

and assuming that the quark chemical potential is flavor-independent ($\equiv \mu_q$) and that the hadron chemical potential is state-independent ($\equiv \mu_h$) it follows that:

$$3\mu_q = \mu_h. \quad (2)$$

The net baryon number density in the quark and hadronic phases is respectively:

$$n_{Bq} = (2 \text{ or } 3)(4a/3\pi^2)\mu_h T^2 \quad (3a)$$

$$n_{Bh} = 2\mu_h T^2 (2\pi)^{-3/2} \{4x_N^{3/2} e^{-x_N} + 2x_\Lambda^{3/2} e^{-x_\Lambda} + 6x_\Sigma^{3/2} e^{-x_\Sigma} + 16x_\Delta^{3/2} e^{-x_\Delta}\} \quad (3b)$$

where $x_N = m_N/T$ ($g_N = 4$, $m_N \simeq 940 \text{ MeV}$), $x_\Lambda = m_\Lambda/T$ ($g_\Lambda = 2$, $m_\Lambda \simeq 1120 \text{ MeV}$), $x_\Sigma = m_\Sigma/T$ ($g_\Sigma = 6$, $m_\Sigma \simeq 1190 \text{ MeV}$), $x_\Delta = m_\Delta/T$ ($g_\Delta = 16$, $m_\Delta \simeq 1230 \text{ MeV}$), and $g_{quark} = 6$ (for each flavor). Only terms of $O(\mu_h/T)$ have been retained in Eqns(3) since $\mu_h/T \sim 10^{-10} \ll 1$. In the expression for n_{Bq} the 2(3) pertains if u and d (u,d, and s) quarks are considered. The baryon number density contrast R between the two phases at the critical temperature is:

$$\begin{aligned} R &= (n_{Bq}/n_{Bh})|_{T_c} \\ &\simeq (2 \text{ or } 3)(2/\pi)^{1/2}(2a/3)[2x_N^{3/2} e^{-x_N} + x_\Lambda^{3/2} e^{-x_\Lambda} \\ &\quad + 3x_\Sigma^{3/2} e^{-x_\Sigma} + 8x_\Delta^{3/2} e^{-x_\Delta}]^{-1} \end{aligned}$$

where $x_i = m_i/T_c$. The baryon number contrast R as a function of T_c is shown in Fig. 1.

From Fig. 1 the importance of including the $I = 3/2$, $J = 3/2$ Δ resonance is very clear: it leads to a reduction in R by a factor of more than 2 for $T_c \gtrsim 150 \text{ MeV}$. The inclusion of the strangeness -1 states is much less important. In addition, there is some question whether or not the strangeness -1 states would be in thermal equilibrium in the hadronic phase. [We have also considered the effect on R of including the next lowest mass baryon state, the Ξ^- , Ξ^0 (strangeness = -2) states. For $T_c \gtrsim 100 \text{ MeV}$, their effect upon R is less than $\sim 10\%$.] Note that for a sufficiently high value of T_c it becomes thermodynamically favorable for the baryon number to reside predominantly in the hadronic phase and $R < 1$.

In sum, for $T_c \gtrsim 150 \text{ MeV}$ the inclusion of the low mass states beyond the neutron and proton ensures that R is less than 10, and for $T_c \geq 200 \text{ MeV}$ that R is less than 2. Thus it is very unlikely that sufficiently large inhomogeneities arise to affect primordial nucleosynthesis, unless T_c is very low.

[The importance of the low lying baryon resonances has also been pointed out recently by Alcock et al⁴, who arrive at essentially the conclusion as ours.]

This work was supported in part by the NASA and DOE (at Fermilab) and by an Alfred P. Sloan Fellowship, and was completed at the Aspen Center for Physics.


References

1. There are actually two transitions associated with the SU(3) color gauge theory (or QCD): the chiral symmetry breaking transition and the deconfinement/confinement (or quark/hadron) transition. In the pure SU(3) theory (no dynamical) quarks, both are strongly first order phase transitions and occur at the same temperature. When the dynamical effects (e.g., screening due to quark loops) of colored fermions (i.e., quarks) are included, the situation is far from being clear. In the very massive quark limit ($m_q \gg \Lambda_{\overline{MS}}$), the quark/hadron transition is strongly first order (equivalent to the pure gauge theory case), but there is no chiral symmetry breaking transition. In the opposite limit, very light quarks ($m_q \ll \Lambda_{\overline{MS}}$), with 3 or more flavors of quarks the chiral transition is first order. The effect of 'light' quarks is to soften the quark/hadron transition, apparently making it second order (or perhaps not even a phase transition at all). The effect of 'sufficiently heavy' quarks is to eliminate the chiral phase transition. In fact it is possible that there exists a range of quark masses for which neither transition is first order (or a phase transition at all). The nature of these transitions for realistic quark masses (i.e., two very light quarks and at least one intermediate mass quark) is still uncertain because of the difficulties of including the effects of dynamical quarks, especially quarks of differing masses. However, recent numerical work suggests that the deconfinement/confinement transition is either weakly first order or second order, with a transition temperature of $\sim 2\Lambda_{\overline{MS}} \sim 100 - 400$ MeV. The following are recent numerical studies of the SU(3) theory and some relevant theoretical papers (note, no attempt was made at completeness): J. Kogut et al., *Phys. Rev. Lett.* **50**, 393 (1983); J. Engels, F. Karsch, and H. Satz, *Phys. Lett.* **113B**, 398 (1982); J. Engels, F. Karsch, H. Satz, and I. Montvay, *Nucl. Phys. B* **B205**[FS5], 545 (1982); B. Svetitsky and F. Fucito, *Phys. Lett.* **131B**, 165 (1983); T. Celik, J. Engels, and H. Satz, *Phys. Lett.* **129B**, 323 (1983); S. A. Gottlieb et al., *Phys. Rev. Lett.* **55**, 1958 (1985); N.

- H. Christ and A. E. Terrano, *Phys. Rev. Lett.* **56**, 111 (1986); J. B. Kogut et al., *Phys. Rev. Lett.* **53**, 644 (1984); F. Fucito and S. Solomon, *Phys. Rev. Lett.* **55**, 2641 (1985); M. Fukugita and A. Ukawa, *Phys. Rev. Lett.* **57**, 503 (1986); E. V. E. Kovacs, D. K. Sinclair, and J. B. Kogut, *Phys. Rev. Lett.* **58**, 751 (1987); J. Kogut and D.K. Sinclair, *Nucl. Phys.* **B280**[FS18], 625 (1987); J. Kogut, *Phys. Rev. Lett.* **56**, 2557 (1986); J. Polonyi, H.W. Wyld, J. Kogut, J. Shigemitsu, and D.K. Sinclair, *Phys. Rev. Lett.* **53**, 644 (1984); T. Banks and A. Ukawa, *Nucl. Phys.* **B225**[FS9], 145 (1983); R.D. Pisarski and F. Wilczek, *Phys. Rev.* **D29**, 338 (1984); J. Kogut, J. Polonyi, H. Wyld, J. Shigemitsu, and D.K. Sinclair, *Nucl. Phys.* **B251**[FS13] (1985); M. Fukugita, S. Ohta, Y. Oyanagi, and A. Ukawa, *Phys. Rev. Lett.* **58**, 2515 (1987).
2. E. Witten, *Phys. Rev.* **D30**, 272 (1984).
 3. J. H. Applegate, C. J. Hogan, and R. J. Scherrer, *Phys. Rev.* **D35**, 1151 (1987).
 4. C. R. Alcock, G. M. Fuller, and G. J. Mathews, *Astrophys. J.*, in press (1988) (UCRL preprint 95896), and work in preparation (1987).
 5. H. Reeves, 'The quark-lithium connection', in *Proc. of the Int'l. School of Physics, Enrico Fermi: Confrontation Between Theories and Observations in Cosmology* (Varenna, July 1987); J. Audouze, *ibid.*
 6. J. Yang, M.S. Turner, G. Steigman, D.N. Schramm, and K.A. Olive, *Astrophys. J.* **281**, 493 (1984); A. M. Boesgaard and G. Steigman, *Ann. Rev. Astron. Astrophys.* **23**, 319 (1985).
 7. Applegate and Hogan have emphasized that the value of R may not be determined so much by equilibrium thermodynamics, but rather more by nonequilibrium processes, such as, shock dissipation, neutrino heat conduction, and expansion of an inhomogeneous, two-phase mixture. See, J. Applegate and C.J. Hogan, *Phys. Rev.* **D31**, 3037 (1985); **D34**, 1938(E) (1986).
 8. W.A. Bardeen, A.J. Buras, D.W. Duke, and T. Muta, *Phys. Rev.* **D18**, 3998 (1978).
 9. E.g., see the recent review of experimental determinations of $\Lambda_{\overline{MS}}$ by D.W. Duke and R.G. Roberts, *Phys. Rep.* **120**, 275 (1985).

Figure Caption

Figure 1 - Equilibrium ratio of net baryon number density in the quark phase to that in the hadron phase as a function of the critical temperature T_c . In the quark phase only 2 quark flavors (u and d) have been used; if the s quark is also in thermal equilibrium and relativistic ($m_s \ll T_c$), then these results must be scaled upward by a factor of 3/2.



- FIGURE 1 -