

Fermi National Accelerator Laboratory

FERMILAB-Pub-93/102-E

E760

Production of the $f_2(1520)$ Resonance in Antiproton-Proton
Annihilations at $\sqrt{s} = 2980$ and 3526 MeV

T.A. Armstrong et al
the E760 Collaboration

*Fermi National Accelerator Laboratory
P.O. Box 500, Batavia, Illinois 60510*

May 1993

Submitted to *Physics Letters*

Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

**PRODUCTION OF THE $f_2(1520)$ RESONANCE IN ANTIPROTON-
PROTON ANNIHILATIONS AT $\sqrt{s} = 2980$ AND 3526 MeV**

T.A. Armstrong⁶, D. Bettoni², V. Bharadwaj¹, C. Biino⁷, G. Borreani²,
D. Broemmelsiek⁴, A. Buzzo³, R. Calabrese², A. Ceccucci⁷, R. Cester⁷, M.
Church¹, P. Dalpiaz², P.F. Dalpiaz², R. Dibenedetto⁷, D. Dimitroyannis⁵,
M.G. Fabbri², J. Fast⁴, A. Gianoli², C.M. Ginsburg⁵, K. Gollwitzer⁴, A.
Hahn¹, M. Hasan⁶, S. Hsueh¹, R. Lewis⁶, E. Luppi², M. Macri³, A.M.
Majewska⁶, M. Mandelkern⁴, F. Marchetto⁷, M. Marinelli³, J. Marques⁴,
W. Marsh¹, M. Martini², M. Masuzawa⁵, E. Menichetti⁷, A. Migliori⁷, R.
Mussa⁷, M. Pallavicini³, S. Palestini⁷, N. Pastrone⁷, C. Patrignani³, J. Peo-
ples Jr.¹, L. Pesando⁷, F. Petrucci², M.G. Pia³, S. Pordes¹, P. Rapidis¹, R.
Ray^{5,1}, J. Reid⁶, G. Rinaudo⁷, B. Roccauzzo⁷, J. Rosen⁵, A. Santroni³, M.
Sarmiento⁵, M. Savrie², A. Scalisi³, J. Schultz⁴, K.K. Seth⁵, A. Smith⁴, G.A.
Smith⁶, M. Sozzi⁷, S. Trokenheim⁵, M.F. Weber⁴, S. Werkema¹, Y. Zhang⁶,
J. Zhao⁵, G. Zioulas⁴.

¹*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, U.S.A.*

²*I.N.F.N. and University of Ferrara, 44100 Ferrara, Italy*

³*I.N.F.N. and University of Genoa, 16146 Genoa, Italy*

⁴*University of California at Irvine, California 92717, U.S.A.*

⁵*Northwestern University, Evanston, Illinois 60208, U.S.A.*

⁶*Pennsylvania State University, University Park, Pennsylvania 16802,
U.S.A.*

⁷*I.N.F.N. and University of Turin, 10125 Turin, Italy.*

Abstract

We present measurements of the $2\pi^0$ mass spectrum in proton anti-proton annihilations. The $f_2(1520)$ resonance is strongly produced in the $3\pi^0$ channel. Its production along with an η is observed, but strongly suppressed. Other features are seen in the data, including the $f_2(1270)$ and structures at $2000 \text{ MeV}/c^2$ and above.

In a separate paper ⁽¹⁾ we reported evidence for several resonances in the 1.5 - 2.1 GeV/c² mass range decaying to $\eta\eta$. This investigation of light quark mesons was carried out during an experiment designed to study charmonium decays. We have also studied the $2\pi^0$ mass spectrum in the same data in the reactions:-

$$\bar{p} + p \rightarrow \pi^0 + \pi^0 + \pi^0 \quad (1)$$

and

$$\bar{p} + p \rightarrow \pi^0 + \pi^0 + \eta. \quad (2)$$

The data reported in this paper were taken in conjunction with scans of the $\eta_c(2980)$ and $^1P_1(3526)$ charmonium resonances. The trigger and reconstruction process for these 6-photon final states are described in Reference (1). After identification of events with six clusters in the central electromagnetic calorimeter, a kinematic fit was made to a 6-photon hypothesis. As discussed in Reference (1), two-photon pairs with an invariant mass in the range 70-200 MeV/c² (449-649 MeV/c²) were identified as coming from a $\pi^0(\eta)$, and the event was then fitted to (1) and (2). After applying cuts, we are left with a total of $1.87 \times 10^6(9.92 \times 10^5)$ events from reaction (1) and $6.07 \times 10^6(3.06 \times 10^5)$ from reaction (2) at 2980(3526) MeV.

We show in Figure 1 Dalitz plots for reaction (1) at (a) 2980 MeV and (b) 3526 MeV. In both plots one can see clearly the $f_2(1270)$, a narrow band near 1500 MeV/c² and a broad structure at around 2000 MeV/c². These features can also be seen in the mass projections, Figures 2a (2980 MeV) and 2b (3526 MeV), superimposed on a background, the magnitude of which suggests that it is due mainly to reflections in the Dalitz plot, rather than non-resonant background. As the 1500 MeV/c² object is well-resolved in the low energy data, we have performed a fit to the mass projections for the region 900-1700 MeV/c² using two Breit-Wigner resonant structures and a polynomial to describe the background, and we find a mass of 1508 ± 10 MeV/c² and a width of 103 ± 15 MeV/c² for the 1500 MeV object.

In a recent experiment at LEAR ⁽²⁾, the Crystal Barrel Collaboration (CB) observed the $f_2(1520)$ as a $2\pi^0$ meson state with $M = 1515 \pm 10$ MeV/c², $\Gamma = 120 \pm 10$ MeV/c², and $J^{PC} = 2^{++}$ in reaction (1) at rest. Possible earlier evidence for the $f_2(1520)$ comes from Gray et al ⁽³⁾, Bridges et al ⁽⁴⁾ and May et al ⁽⁵⁾, in at-rest annihilations. Because of the good agreement of our measurement of the mass and width of the 1500 MeV object with the CB result, we identify it with the $f_2(1520)$. If this is correct, it would add support

to the argument of Dover et al ⁽⁶⁾ that the $f_2(1520)$ is produced uniquely in antiproton-proton annihilation reactions. The broad (250 MeV) structure at 1790 ± 50 MeV/c² required by the CB analysis is not immediately evident in our data.

Figure 3 shows the $\pi\pi$ mass spectrum for reaction (1) at 2980 MeV for 6 intervals in the c.m. production angle, starting with $0 < \cos\theta^* < 0.1$ in (a) and continuing in uniform intervals of 0.1 thereafter. From these data we conclude that, in contrast to the $f_2(1270)$, the $f_2(1520)$ signal is strongest near 90° in the c.m., and decreases thereafter.

For reaction (2), while the $2\pi^0$ combination is unique, the presence of resonances decaying to $\eta\pi$ adds complexity to the Dalitz plots. Bands due to the $a_0(980)$ and $a_2(1320)$, as well as the $f_2(1270)$, are present. The 2980 (3526) MeV data also show possible $\eta\pi$ ($\pi\pi$) structures near 2000 MeV/c².

Figure 4 shows the same distributions as in Figure 3 for reaction (2) at 2980 MeV. Qualitatively, the same features of the $f_2(1270)$ are seen here as in Figure 3. In addition, a small but significant structure near 1500 MeV/c² is seen in four of these plots. This we attribute to the decay of the $f_2(1520)$. Fits to the mass and width in these distributions give an average mass and width of 1525 ± 10 MeV/c² and 111 ± 10 MeV/c². Taking into account the differing acceptances for reactions (1) and (2), we conclude that the ratio of the production of the $f_2(1520)$ in reaction (2) to that in reaction (1) is $17.0 \pm 7\%$, averaging over all c.m. production angles. The error quoted is statistical only. Systematic errors arising from the acceptance calculations are estimated to be of order 50%.

We have also fitted the mass projections of Figures 2a and 2b in the region 1600-2150 MeV/c² to a single Breit-Wigner plus background. We find a mass and width of 1964 ± 35 and 225 ± 50 MeV/c² respectively.

In summary, we have observed the $f_2(1520)$ for the first time in two final states from in-flight antiproton-proton annihilation.

We gratefully acknowledge technical support from our collaborating institutions and the outstanding contribution of the Fermilab Accelerator Division Antiproton Department. This work was funded by the U.S. Department of Energy, the National Science Foundation, and the Italian Istituto Nazionale di Fisica Nucleare.

REFERENCES

1. T.A. Armstrong et al, " Evidence for $\eta\eta$ Resonances in Antiproton-Proton Annihilations at $2950 < \sqrt{s} < 3620$ MeV ", submitted to Phys. Lett. B.
2. E. Aker et al, Phys. Lett. B260 (1991) 249.
3. L. Gray et al, Phys. Rev. D27 (1983) 307.
4. D. Bridges et al, Phys. Rev. Lett. 57 (1986) 1534.
5. B. May et al, Z. Phys. C46 (1990) 191, 203.
6. C.B. Dover et al, Phys. Rev. C43 (1991) 379.

FIGURE CAPTIONS

Fig.1. Dalitz plots for reaction (1) at (a) 2980 and (b) 3526 MeV c.m. energy.

Fig.2. Mass projections for reaction (1) at (a) 2980 and (b) 3526 MeV

Fig.3. Mass projections for reaction (1) at 2980 MeV: (a-f) correspond respectively to the intervals: $0 - 0.1$, $0.1 - 0.2$, $0.2 - 0.3$, $0.3 - 0.4$, $0.4 - 0.5$ and $0.5 - 0.6$ in the cosine of the c.m. di-pion production angle.

Fig.4. Mass projections for reaction (2) at 2980 MeV for the same intervals of the c.m. production angle as in Fig. 3.




FIGURE 1A




FIGURE 1B





FIGURE 2




FIGURE 3




FIGURE 4