



Fermi National Accelerator Laboratory

FERMILAB-Pub-99/040-E

E799-II

**Measurement of the Branching Ratio of $\pi^0 \rightarrow e^+e^-$
using $K_L \rightarrow 3\pi^0$ Decays in Flight**

A. Alavi-Harati et al.
The E799-II Collaboration

*Fermi National Accelerator Laboratory
P.O. Box 500, Batavia, Illinois 60510*

March 1999

Submitted to *Physical Review Letters*

Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Distribution

Approved for public release; further dissemination unlimited.

Copyright Notification

This manuscript has been authored by Universities Research Association, Inc. under contract No. DE-AC02-76CHO3000 with the U.S. Department of Energy. The United States Government and the publisher, by accepting the article for publication, acknowledges that the United States Government retains a nonexclusive, paid-up, irrevocable, worldwide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government Purposes.

Measurement of the branching ratio of $\pi^0 \rightarrow e^+e^-$ using $K_L \rightarrow 3\pi^0$ decays in flight

A. Alavi-Harati^{1,2}, I.F. Albuquerque¹⁰, T. Alexopoulos¹², M. Arenton¹¹, K. Arisaka², S. Averitte¹⁰, A.R. Barker⁵, L. Bellantoni⁷, A. Bellavance⁹, J. Belz¹⁰, R. Ben-David⁷, D.R. Bergman¹⁰, E. Blucher⁴, G.J. Bock⁷, C. Bown⁴, S. Bright⁴, E. Cheu¹, S. Childress⁷, R. Coleman⁷, M.D. Corcoran⁹, G. Corti¹¹, B. Cox¹¹, M.B. Crisler⁷, A.R. Erwin¹², R. Ford⁷, A. Golossanov¹¹, G. Graham⁴, J. Graham⁴, K. Hagan¹¹, E. Halkiadakis¹⁰, K. Hanagaki⁸, S. Hidaka⁸, Y.B. Hsiung⁷, V. Jejer¹¹, J. Jennings², D.A. Jensen⁷, R. Kessler⁴, H.G.E. Kobrak³, J. LaDue⁵, A. Lath¹⁰, A. Ledovskoy¹¹, P.L. McBride⁷, A.P. McManus¹¹, P. Mikelsons⁵, E. Monnier^{4,*}, T. Nakaya⁷, U. Nauenberg⁵, K.S. Nelson¹¹, H. Nguyen⁷, V. O'Dell⁷, M. Pang⁷, R. Pordes⁷, V. Prasad⁴, C. Qiao⁴, B. Quinn⁴, E.J. Ramberg⁷, R.E. Ray⁷, A. Roodman⁴, M. Sadamoto⁸, S. Schnetzer¹⁰, K. Senyo⁸, P. Shanahan⁷, P.S. Shawhan⁴, W. Slater², N. Solomey⁴, S.V. Somalwar¹⁰, R.L. Stone¹⁰, I. Suzuki⁸, E.C. Swallow^{4,6}, R.A. Swanson³, S.A. Taegar¹, R.J. Tesarek¹⁰, G.B. Thomson¹⁰, P.A. Toale⁵, A. Tripathi², R. Tschirhart⁷, Y.W. Wah⁴, J. Wang¹, H.B. White⁷, J. Whitmore⁷, B. Winstein⁴, R. Winston⁴, J.-Y. Wu⁵, T. Yamanaka⁸, E.D. Zimmerman^{4,†,‡}

¹ University of Arizona, Tucson, Arizona 85721

² University of California at Los Angeles, Los Angeles, California 90095

³ University of California at San Diego, La Jolla, California 92093

⁴ The Enrico Fermi Institute, The University of Chicago, Chicago, Illinois 60637

⁵ University of Colorado, Boulder, Colorado 80309

⁶ Elmhurst College, Elmhurst, Illinois 60126

⁷ Fermi National Accelerator Laboratory, Batavia, Illinois 60510

⁸ Osaka University, Toyonaka, Osaka 560 Japan

⁹ Rice University, Houston, Texas 77005

¹⁰ Rutgers University, Piscataway, New Jersey 08855

¹¹ The Department of Physics and Institute of Nuclear and Particle Physics, University of Virginia, Charlottesville, Virginia 22901

¹² University of Wisconsin, Madison, Wisconsin 53706

(March 9, 1999)

The branching ratio of the rare decay $\pi^0 \rightarrow e^+e^-$ has been measured in E799-II, a rare kaon decay experiment using the KTeV detector at Fermilab. The π^0 's were produced in fully-reconstructed $K_L \rightarrow 3\pi^0$ decays in flight. We observed 275 candidate $\pi^0 \rightarrow e^+e^-$ events, with an expected background of 21.4 ± 6.2 events which includes the contribution from Dalitz decays. We measured $\text{BR}(\pi^0 \rightarrow e^+e^-, (m_{e^+e^-}/m_{\pi^0})^2 > 0.95) = (6.09 \pm 0.40 \pm 0.24) \times 10^{-8}$, where the first error is statistical and the second systematic. This result is the first significant observation of the excess rate for this decay above the unitarity lower bound. It is in agreement with expectations from $\pi^0\gamma^*\gamma^*$ form factor models and recent chiral perturbation theory calculations.

PACS numbers: 13.20.Cz, 13.40.Gp, 13.40.Hq

* On leave from C.P.P. Marseille/C.N.R.S., France

† To whom correspondence should be addressed. Electronic address: edz@fnal.gov.

‡ Current address: Columbia University, New York, New York 10027.

The decay $\pi^0 \rightarrow e^+e^-$ has received much experimental and theoretical attention since its branching ratio was first calculated by Drell in 1959 [1]. Within the Standard Model, this decay proceeds predominantly through a two-photon intermediate state, at a rate less than 10^{-7} that of $\pi^0 \rightarrow \gamma\gamma$. Relative to $\pi^0 \rightarrow \gamma\gamma$, $\pi^0 \rightarrow e^+e^-$ is suppressed by a helicity factor $(2m_e/m_{\pi^0})^2$ as well as by two orders of α_{EM} . The contribution to the decay from on-shell internal photons has been calculated exactly in QED [2], and forms the “unitarity bound,” a lower limit on the branching ratio which is $\text{BR}(\pi^0 \rightarrow e^+e^-) \geq 4.69 \times 10^{-8}$, ignoring final-state radiative effects. The contribution from off-shell photons depends on the $\pi^0\gamma^*\gamma^*$ form factor, and is model-dependent. Recent vector meson dominance [3] and chiral perturbation theory [4,5] calculations predict this contribution to be somewhat smaller than that from on-shell photons, giving a total branching ratio of $(6 - 9) \times 10^{-8}$, ignoring radiative corrections.

Earlier experiments have produced conflicting measurements of the branching ratio for this mode. The earliest measurements of $\pi^0 \rightarrow e^+e^-$ were performed by a Geneva-Saclay group in 1978 [6] and by a LAMPF group in 1983 [7]. The Geneva-Saclay group used π^0 's produced by the decay $K^+ \rightarrow \pi^+\pi^0$ in flight; the LAMPF experiment used the charge-exchange process $\pi^-p \rightarrow \pi^0n$ from a 300 MeV/c pion beam. Both experiments favored a branching ratio of $\sim 2 \times 10^{-7}$, which would be very hard to accommodate within the Standard Model. A 1989 search by the SINDRUM collaboration [8], using the charge-exchange reaction with stopped pions, produced a 90% confidence level upper limit of 1.3×10^{-7} , which excluded the central values of both previous measurements. More recently, measurements by BNL E851 [9] and FNAL E799-I [10] have observed the decay at the $(5 - 10) \times 10^{-8}$ level, near the Standard Model expectation. The BNL measurement used the $K^+ \rightarrow \pi^+\pi^0$ decay in flight technique, while the FNAL experiment used $K_L \rightarrow 3\pi^0$ decays in flight as a source of neutral pions.

In this Letter we present a new, precision measurement of $\text{BR}(\pi^0 \rightarrow e^+e^-)$ from E799-II, a rare K_L decay experiment which took data in 1997 at Fermilab. The π^0 's were produced using $K_L \rightarrow 3\pi^0$ decays in flight, where the other two π^0 's in the event decayed to $\gamma\gamma$. The $\pi^0 \rightarrow e^+e^-$ events were normalized to Dalitz decays ($\pi^0 \rightarrow e^+e^-\gamma$) with $m_{e^+e^-} > 65$ MeV/c², which were collected and analyzed simultaneously. High- $m_{e^+e^-}$ events were used in order to keep the charged track kinematic variables as similar as possible for the signal and normalization modes, and thus cancel many detector-related systematic errors.

This technique, which was adapted from E799-I, has significant advantages over those used in other measurements. The π^0 's were produced and decayed in vacuum, eliminating backgrounds and resolution smearing from decay products scattering or converting in charge-exchange targets. The continuum process $K_L \rightarrow \pi^0\pi^0e^+e^-$ has never been observed, and does not pose a significant background. By contrast, the analogous processes in K^+ experiments ($K^+ \rightarrow \pi^+e^+e^-$) and charge-exchange experiments ($\pi^-p \rightarrow e^+e^-X$) both produce large

backgrounds to $\pi^0 \rightarrow e^+e^-$. Reconstruction of the full kaon decay provides redundant kinematic constraints, eliminating all non- $K_L \rightarrow 3\pi^0$ backgrounds.

Fig. 1 shows the E799-II detector. The elements of the spectrometer relevant to this measurement are described below. Two nearly-parallel neutral kaon beams were produced by 800 GeV protons striking a 30 cm BeO target at a targeting angle of 4.8 mrad. Two neutral beams, each one up to $0.35 \mu\text{sr}$, were defined by collimators. A Pb absorber converted photons in the beam, and charged particles were removed by a series of sweeping magnets. After the last collimator and sweeping magnet, 94 m from the target, a 65 m long vacuum decay region began. The decay region ended at a Mylar-Kevlar vacuum window 159 m from the target. The beams in the decay region were composed mostly of neutrons and K_L , with small numbers of K_S , Λ^0 , $\bar{\Lambda}^0$, Ξ^0 , $\bar{\Xi}^0$. These short-lived particles tended to decay upstream. The K_L momentum ranged from ~ 20 to ~ 200 GeV/c; approximately 5% of the K_L decayed before reaching the end of the decay region.

Immediately downstream of the vacuum window was a charged particle spectrometer consisting of four drift chambers with two orthogonal views. Each view comprised two overlapping planes, offset by a half cell. A dipole magnet provided a transverse momentum kick of 205 MeV/c in the horizontal direction. Charged particle resolution in the chambers was $\sim 100 \mu\text{m}$. Helium bags were placed between the chambers to reduce multiple scattering and photon conversions.

Photon energy measurement and electron identification were performed using a 3100-crystal pure CsI electromagnetic calorimeter [11]. The photon energy resolution was $\sim 1\%$, averaged over the energy range typical of $\pi^0 \rightarrow e^+e^-$ events (2–60 GeV). Immediately upstream of the CsI, two overlapping banks of scintillation counters (the “trigger hodoscopes”) provided fast signals for triggering on charged particles. Downstream of the calorimeter, a 10 cm lead wall followed by a scintillator plane formed a hadron veto which rejected at trigger level events with charged pions in the final state. Finally, an eleven-plane photon veto system consisting of lead-scintillator counters throughout the decay region and spectrometer detected particles which left the fiducial volume.

A single trigger was used to collect both $\pi^0 \rightarrow e^+e^-$ candidates and $\pi^0 \rightarrow e^+e^-\gamma$ events for normalization. The trigger required at least 24 GeV of in-time energy in the CsI, and hits in the drift chambers and trigger hodoscopes consistent with at least two tracks. Events were rejected when more than 0.5 GeV was deposited in any photon veto counter, or more than the equivalent of 2.5 minimum ionizing particles were detected in the hadron veto. A hardware processor [12] required at least four energy clusters in the CsI, where a “cluster” was a set of contiguous crystals with at least 1 GeV deposited in each crystal.

Offline, the two tracks were required to form a common vertex inside the decay region. They also had to be electron candidates, defined as tracks which pointed to a CsI cluster whose energy E was within $\pm 8\%$ of the track momentum p .

The reconstructed kaon energy in the event was required to be at least 40 GeV, and each cluster in the CsI had to be at least 1.5 GeV. These cuts reduced the dependence of the result on the CsI trigger thresholds. Events with more than two charged tracks were rejected. Clusters without tracks pointing to them were assumed to be photons. $\pi^0 \rightarrow e^+e^-$ candidates were required to have four photons; $\pi^0 \rightarrow e^+e^-\gamma$ candidates were required to have five.

The photons were reconstructed as $\pi^0 \rightarrow \gamma\gamma$ decays. Assuming that two photons came from a $\pi^0 \rightarrow \gamma\gamma$ decay, we calculated the distance Z of the decay from the CsI (using the assumption that the angle between the photons was small):

$$Z_{12} = (r_{12}/m_{\pi^0})\sqrt{E_1 E_2}$$

where E_i is the energy of photon i and r_{ij} is the transverse separation of photons i and j at the CsI. For each combination of the photons into two π^0 's, we calculated the ‘‘pairing χ^2 ’’ for the hypothesis that the two π^0 decays occurred at the same position. For four-photon $K_L \rightarrow 3\pi^0, \pi^0 \rightarrow e^+e^-$ candidates, there were three pairing combinations, and the one with the lowest χ^2 was selected. For the five-photon $K_L \rightarrow 3\pi^0, \pi^0 \rightarrow e^+e^-\gamma$ candidates, there were fifteen combinations; the best pairing was selected and the unpaired photon was assumed to have come from the $\pi^0 \rightarrow e^+e^-\gamma$ decay.

After selecting a photon pairing, a weighted average was taken of the Z positions of the two π^0 's. The photon four-momenta were calculated assuming the photons originated at that Z position and the transverse position of the reconstructed two-electron vertex. This reconstruction method, which took advantage of the high resolution of the CsI, was used because it allowed kinematic quantities to be calculated in nearly the same way for the signal and normalization modes, thereby canceling certain systematic errors.

The basic reconstruction cuts described below were applied, and the samples obtained were used to study acceptance and backgrounds. The total invariant mass $m_{\pi^0\pi^0 e^+e^-}$ was required to be within 50 MeV/ c^2 of the K_L mass. The total momentum transverse to the kaon direction was required to be less than 30 MeV/ c . For the normalization sample, the Dalitz decay mass $m_{e^+e^-\gamma}$ was required to be within 30 MeV/ c^2 of the π^0 mass, and the reconstructed electron pair mass $m_{e^+e^-} > 70$ MeV/ c^2 in order to avoid systematic errors from mass resolution smearing near the 65 MeV/ c^2 cutoff.

A detailed Monte Carlo (MC) simulation was used to estimate acceptance for the signal and normalization modes, as well as the level of backgrounds in the samples. Both the signal and normalization MC were implemented with radiative corrections based on analytic calculations. The $\pi^0 \rightarrow e^+e^-$ MC used the $\mathcal{O}(\alpha_{EM})$ radiation model of Bergström [13], and the $\pi^0 \rightarrow e^+e^-\gamma$ MC used an $\mathcal{O}(\alpha_{EM}^2)$ calculation based on the work of Mikaelian and Smith [14].

At this stage, the sample in the $\pi^0 \rightarrow e^+e^-$ signal region ($0.132 < m_{e^+e^-} < 0.138$ GeV/ c^2) was background-dominated. Fig. 2 shows the distribution of $m_{e^+e^-}$ for data and for the MC background predictions. The backgrounds,

which all came from $K_L \rightarrow 3\pi^0$ decays, were as follows. Very high- $m_{e^+e^-}$ Dalitz decays ($\pi^0 \rightarrow e^+e^-\gamma$) could be misreconstructed as $\pi^0 \rightarrow e^+e^-$ if the photon was not detected. Another type of background came from decays with four electrons in the final state. When one electron of each sign was soft, the spectrometer magnet could sweep them out of the fiducial volume, leaving only two reconstructible tracks. The four electrons could come from $K_L \rightarrow 3\pi^0$ with multiple $\pi^0 \rightarrow e^+e^-\gamma$ decays, from a $\pi^0 \rightarrow e^+e^-e^+e^-$ decay, or from photon conversions in the $(3.55 \pm 0.17) \times 10^{-3}$ radiation length vacuum window assembly [15].

The four-electron backgrounds fell into two categories. 1) ‘‘Correctly paired’’ four-track backgrounds, where all four electrons came from the same π^0 : these included i) $\pi^0 \rightarrow e^+e^-e^+e^-$ decays, ii) $\pi^0 \rightarrow e^+e^-\gamma$ where the photon from the Dalitz decay converted, iii) $3\pi^0 \rightarrow 6\gamma$ where two photons from the same π^0 converted. In correctly paired four-track events, the reconstructed $m_{e^+e^-}$ was generally below m_{π^0} , and $m_{\pi^0\pi^0 e^+e^-}$ was slightly below m_{K_L} . 2) ‘‘Mispaired’’ four-track backgrounds, where the four electrons came from *different* π^0 's: These included i) events where two π^0 's decayed to $e^+e^-\gamma$, ii) $\pi^0 \rightarrow e^+e^-\gamma$ events where a photon from a different π^0 converted, iii) $3\pi^0 \rightarrow 6\gamma$ events where two photons from different π^0 decays converted. In these cases, because the two observed electrons did not come from the same π^0 , the $m_{e^+e^-}$ distribution was nearly flat. Because the four photons were not from two $\pi^0 \rightarrow \gamma\gamma$ decays, the Z position and the photon four-momenta were misreconstructed, giving a flat $m_{\pi^0\pi^0 e^+e^-}$ distribution as well.

Requiring the pairing χ^2 to be below 4.5 removed 88% of the mispaired four-track background at the cost of 10% of the signal. Tightening the total mass cut to $|m_{\pi^0\pi^0 e^+e^-} - m_{K_L}| < 10$ MeV/ c^2 removed a further 80% of the mispaired four-track background with negligible signal loss.

The correctly paired four-track background could not be reduced significantly with pairing or kinematic cuts. About 99% of these events, as well as 98% of the remaining mispaired background and 8% of signal, were removed by cutting events with evidence of extra in-time tracks in the second drift chamber. The last three cuts were applied to both the signal and normalization samples.

After all cuts, the total background was dominated by high-mass Dalitz decays (18.1 ± 4.7 events). Smaller backgrounds came from correctly paired (2.8 ± 1.1 events) and mispaired (0.5 ± 0.5 events) four-track modes. The errors on the four-track backgrounds were from MC statistics; the error on the Dalitz background reflects MC statistics and a 20% systematic uncertainty in the MC prediction of the misreconstructed $m_{e^+e^-}$ tail. An (18 ± 5)% discrepancy was seen between the data and the MC prediction in the level of the low- $m_{e^+e^-}$ Dalitz background between $0.110 < m_{e^+e^-} < 0.125$ GeV/ c^2 (Fig. 3). We have treated this conservatively as an additional systematic error on the background, added in quadrature. The final background estimate was therefore 21.4 ± 6.2 events.

Radiative corrections to $\pi^0 \rightarrow e^+e^-$ had a significant effect on the acceptance. Internal bremsstrahlung can produce a

$e^+e^-\gamma$ final state with $m_{e^+e^-} < m_{\pi^0}$, indistinguishable from the tree-level Dalitz decay. Following the convention of Ref. [10], we imposed a cutoff $(m_{e^+e^-}/m_{\pi^0})^2 > 0.95$. We thus quote the branching ratio for this range only, after subtracting the small contribution from the Dalitz diagram. In this kinematic region, interference between the two processes is negligible [13].

The normalization data set contained 650 264 events, with negligible backgrounds. The acceptance for Dalitz decays with $m_{e^+e^-} > 65 \text{ MeV}/c^2$ was 1.03%, for kaons which decayed between 90 and 160 m from the target and had momentum between 20 and 200 GeV. The acceptance for the signal mode was 2.52%. Scaling by the ratio of the acceptances, and the Dalitz decay branching ratio, the single-event sensitivity for $\pi^0 \rightarrow e^+e^-\gamma$, $(m_{e^+e^-}/m_{\pi^0})^2 > 0.95$ was 2.40×10^{-10} .

In the data, 275 $\pi^0 \rightarrow e^+e^-$ candidate events were observed (Fig. 3). Subtracting the estimated background yielded the total sample size of 253.6 ± 16.6 events (the error is statistical only).

Significant systematic errors came from parameters of the Dalitz decays used in the normalization, primarily the 2.7% uncertainty in the $\pi^0 \rightarrow e^+e^-\gamma$ branching ratio (1.198 ± 0.032) $\times 10^{-2}$ [16]. In addition, the $m_{e^+e^-}$ cutoff in the normalization Dalitz decays introduced a dependence of the acceptance on the Dalitz decay form factor. The MC used the form factor slope of 0.033 ± 0.003 measured by the CELLO collaboration [17], which gives the result that the $m_{e^+e^-} > 65 \text{ MeV}/c^2$ region contains 3.19% of all Dalitz decays. This form factor measurement used the reaction $e^+e^- \rightarrow \pi^0 e^+e^-$ in a region of spacelike momentum transfer, extrapolating the slope to the kinematic region of the Dalitz decay assuming vector meson dominance. The most recent direct measurement from the Dalitz decay is consistent but less precise [18]. Using the CELLO form factor, the observed $m_{e^+e^-}$ distribution (Fig. 4) was consistent with the MC. The statistical precision of our fit was 0.007, which we have taken to be the uncertainty in the form factor; this translates into an 0.5% systematic error on our measurement of $\pi^0 \rightarrow e^+e^-$.

Although the Dalitz decay branching ratio and the background uncertainty dominated the systematic error, there were some smaller acceptance uncertainties as well. A 1.0% uncertainty was assigned to the efficiency of the pairing χ^2 cut, and a 1.2% systematic error was assigned to the efficiency of the $m_{e^+e^-}$ cut in the normalization sample. These errors were determined from resolution studies using fully-reconstructed Dalitz decays.

Adding all the systematic errors in quadrature, we obtained a total systematic error of 4.0%. Our result for the branching ratio is $\text{BR}(\pi^0 \rightarrow e^+e^-\gamma, (m_{e^+e^-}/m_{\pi^0})^2 > 0.95) = (6.09 \pm 0.40 \pm 0.24) \times 10^{-8}$, where the first error is statistical and the second systematic.

For comparison with the unitarity bound and with theoretical models which neglect final-state radiation, we can invert the radiative corrections and extrapolate our result to the “lowest-order” rate (what the branching ratio would be in the

absence of final-state radiation). This yields $\frac{\Gamma_{e^+e^-}^{\text{lowest order}}}{\Gamma_{\text{all}}} = (7.04 \pm 0.46(\text{stat}) \pm 0.28(\text{syst})) \times 10^{-8}$, which is over four standard deviations above the unitarity bound. This result, which is in agreement with recent Standard Model predictions, represents the first statistically significant observation of an excess above unitarity.

E799-II expects to accumulate 2 to 4 times more data in a 1999 run, which will allow a further refinement of this measurement. This result should provide constraints for predictions of related decay modes such as $\eta \rightarrow \mu^+\mu^-$ and $K_L \rightarrow \mu^+\mu^-$, and we hope that future experiments will be able to test these predictions.

We gratefully acknowledge the support and effort of the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported in part by the U.S. Department of Energy, The National Science Foundation and The Ministry of Education and Science of Japan. In addition, A.R.B., E.B. and S.V.S. acknowledge support from the NYI program of the NSF; A.R.B. and E.B. from the Alfred P. Sloan Foundation; E.B. from the OJI program of the DOE; K.H., T.N. and M.S. from the Japan Society for the Promotion of Science. P.S.S. acknowledges receipt of a Grainger Fellowship.

-
- [1] S. Drell, *Nuovo Cim.* **XI**, 693 (1959).
 - [2] L. Bergström, *Z. Phys.* **C14**, 129 (1982).
 - [3] Ll. Ametller, A. Bramon, and E. Massó *Phys. Rev.* **D48**, 3388 (1993).
 - [4] M. J. Savage, M. Luke, and M. B. Wise *Phys. Lett.* **B291**, 481 (1992).
 - [5] D. Gomez Dumm and A. Pich, *Phys. Rev. Lett.* **80**, 4633 (1998).
 - [6] J. Fischer *et al.*, *Phys. Lett.* **73B**, 364 (1978).
 - [7] J. S. Frank *et al.*, *Phys. Rev.* **D28**, 423 (1983).
 - [8] C. Niebuhr *et al.*, *Phys. Rev.* **D40**, 2976 (1989).
 - [9] A. Deshpande *et al.*, *Phys. Rev. Lett.* **71**, 27 (1993).
 - [10] K. S. McFarland *et al.*, *Phys. Rev. Lett.* **71**, 31 (1993).
 - [11] A. J. Roodman, “The KTeV Pure CsI Calorimeter,” *Proceedings of the VII International Conference on Calorimetry*, World Scientific (1998).
 - [12] C. Bown *et al.*, *Nucl. Inst. Meth.* **A369**, 248 (1996).
 - [13] L. Bergström, *Z. Phys.* **C20**, 135 (1983).
 - [14] K. O. Mikaelian and J. S. Smith, *Phys. Rev.* **D5**, 1763 (1972) and *Phys. Rev.* **D5**, 2891 (1972).
 - [15] E. D. Zimmerman, Report EFI-98-43, The University of Chicago, (1998)
 - [16] Particle Data Group, *Eur. Phys. J.* **C3**, 1 (1998).
 - [17] H. Behrend *et al.*, *Z. Phys.* **C49**, 401 (1991).
 - [18] R. Meijer Drees *et al.*, *Phys. Rev.* **D45**, 1439 (1992).




FIG. 1. Plan view of the KTeV spectrometer as configured for E799-II. The horizontal scale is compressed.




FIG. 4. Reconstructed $m_{e^+e^-}$ for normalization Dalitz decays.




FIG. 2. Reconstructed $m_{e^+e^-}$ after basic reconstruction cuts. Left: background MC predictions. Right: E799-II data overlaid on sum of background predictions. (Backgrounds are normalized to the observed number of fully-reconstructed $\pi^0 \rightarrow e^+e^-\gamma$ decays.)




FIG. 3. Reconstructed $m_{e^+e^-}$ after all cuts. Background and signal MC are shown along with data (points with errors). The vertical dotted lines indicate the defined signal region ($0.132 < m_{e^+e^-} < 0.138 \text{ GeV}/c^2$).