Fermilab FERMILAB-CONF-05-218-E CDF JUNE 2005

ELECTROWEAK MEASUREMENTS AT CDF *

A. SIDOTI †

Laboratoire de Physique Nuclèaire et de Hautes Energies
Universitè "Pierre et Marie Curie" (Paris VI)
4, Place Jussieu
75252 Paris Cedex 05, France

We present some recent measurements on electroweak physics using data collected by the CDF experiment at the Tevatron proton anti-proton collider ($\sqrt{s}$ = 1.96TeV) at Fermilab (Batavia, Ill, USA).

1. Introduction

The CDF electroweak physics program is one of the key components of the RunII. Electroweak measurements are complementary to those performed at e^+e^- machines (LEP and SLD). The former can produce a larger number of W bosons and can produce $Z/\gamma^\star$ at higher invariant mass with respect of the latters.

We will review CDF electroweak physics measurements using data collected from February 2002. The integrated luminosity of data ranges from 64 pb^{-1} to ∼200 pb^{-1} depending on the measurement.

2. W and Z inclusive cross section measurements

Due to the high branching ratios and clean signature W and Z bosons are identified through their leptonic decays. Inclusive cross sections of both W and Z have been measured using all the three leptons: electrons, muons and *taus* and using all the available subdetectors of CDF extending in particular the geometric acceptance at high pseudorapidity η[a]. All the measurements

*On behalf of the CDF collaboration

†Work supported by Research Training Network of E.U. "Probe of New Physics" HPRN-CT-2002-00292 Contract

[a] η is related to the azimuthal angle through the relation $\eta = -\log\tan(\theta/2)$

performed are shown in Fig.1 and are in agreement with the theoretical predictions [1].



Figure 1. $\sigma(p\bar{p} \to W) \times BF(W \to \ell\nu_\ell)$ (left) and $\sigma(p\bar{p} \to Z) \times BF(Z \to \ell\ell)$ (right) measured at CDF. The blue band indicates the theoretical (NNLO) predictions.

A stringent test of the Standard Model can be performed by evaluating the ratio R of the W and Z boson cross sections:

$$R = \frac{\sigma(p\bar{p} \to W) \times \mathrm{BF}(W \to \ell\nu_\ell)}{\sigma(p\bar{p} \to Z) \times \mathrm{BF}(Z \to \ell\ell)}$$

that can be written as

$$R = \frac{\sigma(p\bar{p} \to W)}{\sigma(p\bar{p} \to Z)} \times \frac{\Gamma(Z)}{\Gamma(Z \to \ell\ell)} \times \frac{\Gamma(W \to \nu\nu_\ell)}{\Gamma(W)}.$$

Inserting the theoretical values of the total cross sections and of the partial width $W \to \ell\nu_\ell$ and the experimental partial and total widths of the Z boson from LEP, it is possibile to extract indirectly the total width of the W boson. The measured value by CDF is:

$$\Gamma(W) = 2079 \pm 41 \text{ MeV} \quad \text{CDF } e + \mu \text{ channel } \int \mathcal{L}dt = 72 \text{ pb}^{-1}$$
$$\Gamma(W) = \quad 2056 \pm 44 \text{ MeV} \quad \text{CDF } \mu \text{ channel } \int \mathcal{L}dt = 194 \text{ pb}^{-1}$$

well in agreement with both the PDG world average[3] and the theoretical predictions:

$$\Gamma(W)_{\mathrm{PDG}} = 2118 \pm 41 \text{ MeV} \quad \Gamma(W)_{\mathrm{Th.}} = 2092.1 \pm 2.5 \text{ MeV}$$

3. W Boson Asymmetries

A precise charge asymmetry as a function of rapidity of the W boson provides constraints on the parton fluxes of the incoming protons and therefore provides a better determination of the parton distribution functions (pdf). The leptonic decay of the W boson makes difficult to measure directly the W rapidity itself. Instead, the electron asymmetry is measured that is the convolution of the W production charge asymmetry and the V-A asymmetry from the W decay. The lepton charge aymmetry is defined as:

$$A(\eta_\ell) = \frac{d\sigma_+/d\eta_\ell - d\sigma_-/d\eta_\ell}{d\sigma_+/d\eta_\ell + d\sigma_-/d\eta_\ell}$$

where η_ℓ is the pseudorapidity of the lepton. CDF has measured the W charge asymmetry in the electron channel. The result as a function of the electron rapidity is shown in Fig.2. The measurement has been corrected for the effect of charge mis-identification and background contributions both dependent on the pseudorapidity. An additional selection $35 < E_T < 45$ GeV is applied on the electron energy to increase the sensitivity to different pdf. In fact the direction of an electron with higher energies is closer to the direction of emission of the W boson enhancing therefore the W production asymmetry[4]



Figure 2. W charge asymmetry as a function of the electron pseudorapidity, with $35 < E_T < 45$ GeV. Comparison with expectations from different pdf are shown.

4. Di-Boson Production

Di-boson processes probe electroweak gauge bosons interactions as well as sources of physics beyond Standard Model.

4.1. *$W\gamma$ and $Z\gamma$ production*

When one of the bosons is a photon the main backgounds are π_0 and jets faking a photon. To reduce these, CDF exploits the high spatial resolution of the electromagnetic calorimeter provided by the *showermax* detector[5] and a track isolation criterium. The $W\gamma \to \ell\nu_\ell\gamma$ process has been studied by CDF in the electron and muon channel. Candidate events are selected requiring a high-P_T momentum lepton with large missing transverse energy and $M_T(\ell,\nu) < 120\,\mathrm{GeV}/c^2$. The presence of a photon with $E_T > 7$ GeV well separated from the lepton ($\Delta R(\ell,\gamma) > 0.7$[b]) is required. The cross section times branching fraction measured is:

$$\sigma \times \mathrm{BF}(W\gamma \to \ell\nu_\ell\gamma) = 18.1 \pm 1.6(\mathrm{stat.}) \pm 2.4(\mathrm{syst.}) \pm 1.2(\mathrm{lum.})\mathrm{pb}^{-1}$$

The predicted cross section including the photon acceptance is [7]:

$$\sigma \times \mathrm{BF}(W\gamma \to \ell\nu_\ell\gamma)_{\mathrm{Th.}} = 19.3 \pm 1.4 \text{ pb}$$

The photon transverse energy is shown in Fig.3.

$Z\gamma \to \ell\ell\gamma$ candidates are selected requiring two oppositely charged high momentum leptons with a dilepton invariant mass such that $M_{\ell,\ell} > 40$ GeV/c^2. The photon selection is the same of the $W\gamma \to \ell\nu_\ell\gamma$. The cross section times branching fraction measured is[6]:

$$\sigma \times \mathrm{BF}(Z\gamma \to \ell\ell\gamma) = 4.6 \pm 0.5(\mathrm{syst} + \mathrm{stat}) \pm 0.3(\mathrm{lum.})\mathrm{pb}^{-1}$$

The predicted cross section including the photon acceptance is [7]:

$$\sigma \times \mathrm{BF}(W\gamma \to \ell\nu_\ell\gamma)_{\mathrm{Th.}} = 4.5 \pm 0.3$$

4.2. *WW Production*

WW production has been studied at CDF in the $WW \to \ell\nu_\ell\ell'\nu'_\ell$ channel looking for two oppositely charged high P_T leptons and large missing transverse energy in the final state. Using and integrated luminosity of 184 pb^{-1} 17 candidate events have been found with an estimated background of $5.0^{+2.2}_{-0.8}$ events. The cross section measured is[8]:

$$\sigma(p\bar{p} \to WW) = 14.6^{+5.8}_{-5.1}(\mathrm{stat})^{+1.8}_{-3.0}(\mathrm{syst}) \pm 0.9(\mathrm{lum})\mathrm{pb}$$

well in agreement with the predicted cross section:

$$\sigma(p\bar{p} \to WW)_{\mathrm{Th.}} = 11.3 \pm 1.3\mathrm{pb}$$

[b] $R = \sqrt{\Delta\phi^2 + \Delta\eta^2}$



Figure 3. Photon E_T (left) and Cluster transverse mass (right) for $W\gamma \to \ell\nu\gamma$ shown for observed events together with signal and background expectations. Anomalous gauge coupling will increase the high energy tail of the photon E_T distribution.

5. Conclusions

CDF is producing interesting results in the electroweak sector. Most of them have a precision that is limited by statistics. Therefore we are approaching a period where, with the available datasets, CDF can produce electroweak physics measurements with the smallest single experiment uncertainties. We are also refining the analysis using new methods that will decrease the systematics uncertainties. So far, no SM deviations have been observed.

Acknowledgments

It is a pleasure to thank all those at CDF and at Fermilab that are working hard for the success of the RunII of the Tevatron. I would also like to thank the organizers of this wonderful Lake Louise Winter Institute.

References

1. P.J. Sutton *et al.*, Phys Rev **D45**, 2349 (1992); P.J. Rijken *et al.*, Phys Rev **D51** 44 (1995); R. Hamberg *et al.*, Nucl. Phys. **B359** 343, (1991), R.V. Harlander *et al.*, Phys. Rev. Lett. **88** 201801 (2002).
2. D. Acosta *et al.* [CDF II Collaboration], Phys. Rev. Lett. **94**, 091803 (2005).
3. S. Eidelman *et al*, Phys. Lett. **B 592**, 1 (2004).
4. D. Acosta *et al.* [CDF Collaboration], Phys. Rev. D **71**, 051104 (2005).
5. G. Apollinari, *et al.* Nucl. Instrum. Meth. A**412**,515-526, (1998).
6. D. Acosta *et al.* [CDF II Collaboration], Phys. Rev. Lett. **94**, 041803 (2005).
7. U. Baur, T. Han and J. Ohnemus, Phys. Rev. D **53** (1996) 1098.
8. D. Acosta *et al.* [CDF Collaboration], `hep-ex/0501050`.