FERMILAB-PUB-11-171-E

arXiv:1104.1585v1 [hep-ex] 8 Apr 2011

Search for the Rare Radiative Decay: $W \to \pi\gamma$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV

T. Aaltonen,[21] B. Álvarez González[w],[9] S. Amerio,[41] D. Amidei,[32] A. Anastassov,[36] A. Annovi,[17] J. Antos,[12] G. Apollinari,[15] J.A. Appel,[15] A. Apresyan,[46] T. Arisawa,[56] A. Artikov,[13] J. Asaadi,[51] W. Ashmanskas,[15] B. Auerbach,[59] A. Aurisano,[51] F. Azfar,[40] W. Badgett,[15] A. Barbaro-Galtieri,[26] V.E. Barnes,[46] B.A. Barnett,[23] P. Barria[dd],[44] P. Bartos,[12] M. Bauce[bb],[41] G. Bauer,[30] F. Bedeschi,[60] D. Beecher,[28] S. Behari,[23] G. Bellettini[bb],[60] J. Bellinger,[58] D. Benjamin,[14] A. Beretvas,[15] A. Bhatti,[48] M. Binkley*,[15] D. Bisello[bb],[41] I. Bizjak[hh],[28] K.R. Bland,[5] B. Blumenfeld,[23] A. Bocci,[14] A. Bodek,[47] D. Bortoletto,[46] J. Boudreau,[45] A. Boveia,[11] B. Brau[a],[15] L. Brigliadori[aa],[6] A. Brisuda,[12] C. Bromberg,[33] E. Brucken,[21] M. Bucciantonio[cc],[44] J. Budagov,[13] H.S. Budd,[47] S. Budd,[22] K. Burkett,[15] G. Busetto[bb],[41] P. Bussey,[19] A. Buzatu,[31] C. Calancha,[29] S. Camarda,[4] M. Campanelli,[33] M. Campbell,[32] F. Canelli[12],[15] A. Canepa,[43] B. Carls,[22] D. Carlsmith,[58] R. Carosi,[44] S. Carrillo[k],[16] S. Carron,[15] B. Casal,[9] M. Casarsa,[15] A. Castro[aa],[6] P. Catastini,[15] D. Cauz,[52] V. Cavaliere[cc],[44] M. Cavalli-Sforza,[4] A. Cerri[f],[26] L. Cerrito[q],[28] Y.C. Chen,[1] M. Chertok,[7] G. Chiarelli,[44] G. Chlachidze,[15] F. Chlebana,[15] K. Cho,[25] D. Chokheli,[13] J.P. Chou,[20] W.H. Chung,[58] Y.S. Chung,[47] C.I. Ciobanu,[42] M.A. Ciocci[dd],[44] A. Clark,[18] G. Compostella[bb],[41] M.E. Convery,[15] J. Conway,[7] M.Corbo,[42] M. Cordelli,[17] C.A. Cox,[7] D.J. Cox,[7] F. Crescioli[cc],[44] C. Cuenca Almenar,[59] J. Cuevas[w],[9] R. Culbertson,[15] D. Dagenhart,[15] N. d'Ascenzo[u],[42] M. Datta,[15] P. de Barbaro,[47] S. De Cecco,[49] G. De Lorenzo,[4] M. Dell'Orso[cc],[44] C. Deluca,[4] L. Demortier,[48] J. Deng[c],[14] M. Deninno,[6] F. Devoto,[21] M. d'Errico[bb],[41] A. Di Canto[cc],[44] B. Di Ruzza,[44] J.R. Dittmann,[5] M. D'Onofrio,[27] S. Donati[cc],[44] P. Dong,[15] M. Dorigo,[52] T. Dorigo,[41] K. Ebina,[56] A. Elagin,[51] A. Eppig,[32] R. Erbacher,[7] D. Errede,[22] S. Errede,[22] N. Ershaidat[z],[42] R. Eusebi,[51] H.C. Fang,[26] S. Farrington,[40] M. Feindt,[24] J.P. Fernandez,[29]

Operated by Fermi Research Alliance, LLC under Contract No. DE-AC02-07CH11359 with the United States Department of Energy

C. Ferrazza[ee],[44] R. Field,[16] G. Flanagan[s],[46] R. Forrest,[7] M.J. Frank,[5] M. Franklin,[20] J.C. Freeman,[15]
Y. Funakoshi,[56] I. Furic,[16] M. Gallinaro,[48] J. Galyardt,[10] J.E. Garcia,[18] A.F. Garfinkel,[46] P. Garosi[dd],[44]
H. Gerberich,[22] E. Gerchtein,[15] S. Giagu[ff],[49] V. Giakoumopoulou,[3] P. Giannetti,[44] K. Gibson,[45] C.M. Ginsburg,[15]
N. Giokaris,[3] P. Giromini,[17] M. Giunta,[44] G. Giurgiu,[23] V. Glagolev,[13] D. Glenzinski,[15] M. Gold,[35] D. Goldin,[51]
N. Goldschmidt,[16] A. Golossanov,[15] G. Gomez,[9] G. Gomez-Ceballos,[30] M. Goncharov,[30] O. González,[29] I. Gorelov,[35]
A.T. Goshaw,[14] K. Goulianos,[48] S. Grinstein,[4] C. Grosso-Pilcher,[11] R.C. Group,[55] J. Guimaraes da Costa,[20]
Z. Gunay-Unalan,[33] C. Haber,[26] S.R. Hahn,[15] E. Halkiadakis,[50] A. Hamaguchi,[39] J.Y. Han,[47] F. Happacher,[17]
K. Hara,[53] D. Hare,[50] M. Hare,[54] R.F. Harr,[57] K. Hatakeyama,[5] C. Hays,[40] M. Heck,[24] J. Heinrich,[43]
M. Herndon,[58] S. Hewamanage,[5] D. Hidas,[50] A. Hocker,[15] W. Hopkins[g],[15] D. Horn,[24] S. Hou,[1] R.E. Hughes,[37]
M. Hurwitz,[11] U. Husemann,[59] N. Hussain,[31] M. Hussein,[33] J. Huston,[33] G. Introzzi,[44] M. Iori[ff],[49] A. Ivanov[o],[7]
E. James,[15] D. Jang,[10] B. Jayatilaka,[14] E.J. Jeon,[25] M.K. Jha,[6] S. Jindariani,[15] W. Johnson,[7] M. Jones,[46]
K.K. Joo,[25] S.Y. Jun,[10] T.R. Junk,[15] T. Kamon,[51] P.E. Karchin,[57] Y. Kato[n],[39] W. Ketchum,[11] J. Keung,[43]
V. Khotilovich,[51] B. Kilminster,[15] D.H. Kim,[25] H.S. Kim,[25] H.W. Kim,[25] J.E. Kim,[25] M.J. Kim,[17] S.B. Kim,[25]
S.H. Kim,[53] Y.K. Kim,[11] N. Kimura,[56] M. Kirby,[15] S. Klimenko,[16] K. Kondo,[56] D.J. Kong,[25] J. Konigsberg,[16]
A.V. Kotwal,[14] M. Kreps,[24] J. Kroll,[43] D. Krop,[11] N. Krumnack[l],[5] M. Kruse,[14] V. Krutelyov[d],[51] T. Kuhr,[24]
M. Kurata,[53] S. Kwang,[11] A.T. Laasanen,[46] S. Lami,[44] S. Lammel,[15] M. Lancaster,[28] R.L. Lander,[7] K. Lannon[v],[37]
A. Lath,[50] G. Latino[cc],[44] T. LeCompte,[2] E. Lee,[51] H.S. Lee,[11] J.S. Lee,[25] S.W. Lee[x],[51] S. Leo[cc],[44] S. Leone,[44]
C.M. Lester,[43] J.D. Lewis,[15] A. Limosani[r],[14] C.-J. Lin,[26] J. Linacre,[40] M. Lindgren,[15] E. Lipeles,[43] A. Lister,[18]
D.O. Litvintsev,[15] C. Liu,[45] Q. Liu,[46] T. Liu,[15] S. Lockwitz,[59] N.S. Lockyer,[43] A. Loginov,[59] D. Lucchesi[bb],[41]
J. Lueck,[24] P. Lujan,[26] P. Lukens,[15] G. Lungu,[48] J. Lys,[26] R. Lysak,[12] R. Madrak,[15] K. Maeshima,[15] K. Makhoul,[30]
P. Maksimovic,[23] S. Malik,[48] G. Manca[b],[27] A. Manousakis-Katsikakis,[3] F. Margaroli,[46] C. Marino,[24] M. Martínez,[4]

R. Martínez-Ballarín,[29] P. Mastrandrea,[49] M. Mathis,[23] M.E. Mattson,[57] P. Mazzanti,[6] K.S. McFarland,[47] P. McIntyre,[51] R. McNulty[i],[27] A. Mehta,[27] P. Mehtala,[21] A. Menzione,[44] C. Mesropian,[48] T. Miao,[15] D. Mietlicki,[32] A. Mitra,[1] H. Miyake,[53] S. Moed,[20] N. Moggi,[6] M.N. Mondragon[k],[15] C.S. Moon,[25] R. Moore,[15] M.J. Morello,[15] J. Morlock,[24] P. Movilla Fernandez,[15] A. Mukherjee,[15] Th. Muller,[24] P. Murat,[15] M. Mussini[aa],[6] J. Nachtman[m],[15] Y. Nagai,[53] J. Naganoma,[56] I. Nakano,[38] A. Napier,[54] J. Nett,[51] C. Neu,[55] M.S. Neubauer,[22] J. Nielsen[e],[26] L. Nodulman,[2] O. Norniella,[22] E. Nurse,[28] L. Oakes,[40] S.H. Oh,[14] Y.D. Oh,[25] I. Oksuzian,[55] T. Okusawa,[39] R. Orava,[21] L. Ortolan,[4] S. Pagan Griso[bb],[41] C. Pagliarone,[52] E. Palencia[f],[9] V. Papadimitriou,[15] A.A. Paramonov,[2] J. Patrick,[15] G. Pauletta[gg],[52] M. Paulini,[10] C. Paus,[30] D.E. Pellett,[7] A. Penzo,[52] T.J. Phillips,[14] G. Piacentino,[44] E. Pianori,[43] J. Pilot,[37] K. Pitts,[22] C. Plager,[8] L. Pondrom,[58] K. Potamianos,[46] O. Poukhov*,[13] F. Prokoshin[y],[13] A. Pronko,[15] F. Ptohos[h],[17] E. Pueschel,[10] G. Punzi[cc],[44] J. Pursley,[58] A. Rahaman,[45] V. Ramakrishnan,[58] N. Ranjan,[46] I. Redondo,[29] P. Renton,[40] M. Rescigno,[49] F. Rimondi[aa],[6] L. Ristori[45],[15] A. Robson,[19] T. Rodrigo,[9] T. Rodriguez,[43] E. Rogers,[22] S. Rolli,[54] R. Roser,[15] M. Rossi,[52] F. Rubbo,[15] F. Ruffini[dd],[44] A. Ruiz,[9] J. Russ,[10] V. Rusu,[15] A. Safonov,[51] W.K. Sakumoto,[47] Y. Sakurai,[56] L. Santi[gg],[52] L. Sartori,[44] K. Sato,[53] V. Saveliev[u],[42] A. Savoy-Navarro,[42] P. Schlabach,[15] A. Schmidt,[24] E.E. Schmidt,[15] M.P. Schmidt*,[59] M. Schmitt,[36] T. Schwarz,[7] L. Scodellaro,[9] A. Scribano[dd],[44] F. Scuri,[44] A. Sedov,[46] S. Seidel,[35] Y. Seiya,[39] A. Semenov,[13] F. Sforza[cc],[44] A. Sfyrla,[22] S.Z. Shalhout,[7] T. Shears,[27] P.F. Shepard,[45] M. Shimojima[t],[53] S. Shiraishi,[11] M. Shochet,[11] I. Shreyber,[34] A. Simonenko,[13] P. Sinervo,[31] A. Sissakian*,[13] K. Sliwa,[54] J.R. Smith,[7] F.D. Snider,[15] A. Soha,[15] S. Somalwar,[50] V. Sorin,[4] P. Squillacioti,[15] M. Stancari,[15] M. Stanitzki,[59] R. St. Denis,[19] B. Stelzer,[31] O. Stelzer-Chilton,[31] D. Stentz,[36] J. Strologas,[35] G.L. Strycker,[32] Y. Sudo,[53] A. Sukhanov,[16] I. Suslov,[13] K. Takemasa,[53] Y. Takeuchi,[53] J. Tang,[11] M. Tecchio,[32] P.K. Teng,[1] J. Thom[g],[15] J. Thome,[10] G.A. Thompson,[22] E. Thomson,[43] P. Ttito-Guzmán,[29] S. Tkaczyk,[15] D. Toback,[51] S. Tokar,[12]

K. Tollefson,[33] T. Tomura,[53] D. Tonelli,[15] S. Torre,[17] D. Torretta,[15] P. Totaro,[41] M. Trovato[ee],[44] Y. Tu,[43] F. Ukegawa,[53] S. Uozumi,[25] A. Varganov,[32] F. Vázquez[k],[16] G. Velev,[15] C. Vellidis,[3] M. Vidal,[29] I. Vila,[9] R. Vilar,[9] J. Vizán,[9] M. Vogel,[35] G. Volpi[cc],[44] P. Wagner,[43] R.L. Wagner,[15] T. Wakisaka,[39] R. Wallny,[8] S.M. Wang,[1] A. Warburton,[31] D. Waters,[28] M. Weinberger,[51] W.C. Wester III,[15] B. Whitehouse,[54] D. Whiteson[c],[43] A.B. Wicklund,[2] E. Wicklund,[15] S. Wilbur,[11] F. Wick,[24] H.H. Williams,[43] J.S. Wilson,[37] P. Wilson,[15] B.L. Winer,[37] P. Wittich[g],[15] S. Wolbers,[15] H. Wolfe,[37] T. Wright,[32] X. Wu,[18] Z. Wu,[5] K. Yamamoto,[39] J. Yamaoka,[14] T. Yang,[15] U.K. Yang[p],[11] Y.C. Yang,[25] W.-M. Yao,[26] G.P. Yeh,[15] K. Yi[m],[15] J. Yoh,[15] K. Yorita,[56] T. Yoshida[j],[39] G.B. Yu,[14] I. Yu,[25] S.S. Yu,[15] J.C. Yun,[15] A. Zanetti,[52] Y. Zeng,[14] and S. Zucchelli[aa6]

(CDF Collaboration[†])

[1]*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*

[2]*Argonne National Laboratory, Argonne, Illinois 60439, USA*

[3]*University of Athens, 157 71 Athens, Greece*

[4]*Institut de Fisica d'Altes Energies, ICREA, Universitat Autonoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*

[5]*Baylor University, Waco, Texas 76798, USA*

[6]*Istituto Nazionale di Fisica Nucleare Bologna, [aa]University of Bologna, I-40127 Bologna, Italy*

[7]*University of California, Davis, Davis, California 95616, USA*

[8]*University of California, Los Angeles, Los Angeles, California 90024, USA*

[9]*Instituto de Fisica de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*

[10]*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*

[11]*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*

[12]*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*

[13]*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*

[14]*Duke University, Durham, North Carolina 27708, USA*

[15]*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*

[16]*University of Florida, Gainesville, Florida 32611, USA*

[17]*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*

[18]*University of Geneva, CH-1211 Geneva 4, Switzerland*

[19]*Glasgow University, Glasgow G12 8QQ, United Kingdom*

[20]*Harvard University, Cambridge, Massachusetts 02138, USA*

[21]*Division of High Energy Physics, Department of Physics,*

University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland

[22]*University of Illinois, Urbana, Illinois 61801, USA*

[23]*The Johns Hopkins University, Baltimore, Maryland 21218, USA*

[24]*Institut für Experimentelle Kernphysik, Karlsruhe Institute of Technology, D-76131 Karlsruhe, Germany*

[25]*Center for High Energy Physics: Kyungpook National University,*

Daegu 702-701, Korea; Seoul National University, Seoul 151-742,

Korea; Sungkyunkwan University, Suwon 440-746,

Korea; Korea Institute of Science and Technology Information,

Daejeon 305-806, Korea; Chonnam National University, Gwangju 500-757,

Korea; Chonbuk National University, Jeonju 561-756, Korea

[26]*Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*

[27]*University of Liverpool, Liverpool L69 7ZE, United Kingdom*

[28]*University College London, London WC1E 6BT, United Kingdom*

[29]*Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain*

[30]*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

[31]*Institute of Particle Physics: McGill University, Montréal, Québec,*

Canada H3A 2T8; Simon Fraser University, Burnaby, British Columbia,

Canada V5A 1S6; University of Toronto, Toronto, Ontario,

Canada M5S 1A7; and TRIUMF, Vancouver, British Columbia, Canada V6T 2A3

[32]*University of Michigan, Ann Arbor, Michigan 48109, USA*

[33]*Michigan State University, East Lansing, Michigan 48824, USA*

[34]*Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia*

[35]*University of New Mexico, Albuquerque, New Mexico 87131, USA*

[36]*Northwestern University, Evanston, Illinois 60208, USA*

[37]*The Ohio State University, Columbus, Ohio 43210, USA*

[38]*Okayama University, Okayama 700-8530, Japan*

[39]*Osaka City University, Osaka 588, Japan*

[40]*University of Oxford, Oxford OX1 3RH, United Kingdom*

[41]*Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento,* [bb]*University of Padova, I-35131 Padova, Italy*

[42]*LPNHE, Universite Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France*

[43]*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*

[44]*Istituto Nazionale di Fisica Nucleare Pisa,* [cc]*University of Pisa,*

[dd]*University of Siena and* [ee]*Scuola Normale Superiore, I-56127 Pisa, Italy*

[45]*University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA*

[46]*Purdue University, West Lafayette, Indiana 47907, USA*

[47]*University of Rochester, Rochester, New York 14627, USA*

[48]*The Rockefeller University, New York, New York 10065, USA*

[49]*Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1,*

[ff]*Sapienza Università di Roma, I-00185 Roma, Italy*

[50]*Rutgers University, Piscataway, New Jersey 08855, USA*

[51]*Texas A&M University, College Station, Texas 77843, USA*

[52]*Istituto Nazionale di Fisica Nucleare Trieste/Udine,*

I-34100 Trieste, [gg]*University of Trieste/Udine, I-33100 Udine, Italy*

[53] *University of Tsukuba, Tsukuba, Ibaraki 305, Japan*

[54] *Tufts University, Medford, Massachusetts 02155, USA*

[55] *University of Virginia, Charlottesville, VA 22906, USA*

[56] *Waseda University, Tokyo 169, Japan*

[57] *Wayne State University, Detroit, Michigan 48201, USA*

[58] *University of Wisconsin, Madison, Wisconsin 53706, USA*

[59] *Yale University, New Haven, Connecticut 06520, USA*

[60] *Istituto Nazionale di Fisica Nucleare Pisa,* [bb] *University of Pisa,*

[cc] *University of Siena and* [dd] *Scuola Normale Superiore, I-56127 Pisa, Italy*

(Dated: April 11, 2011, version 1.1)

We present a search for the rare radiative decay $W^{\pm} \to \pi^{\pm}\gamma$ using data corresponding to an integrated luminosity of 4.3 fb^{-1} of proton-antiproton collisions at a center of mass energy of 1.96 TeV collected by the CDF experiment at Fermilab. As no statistically significant signal is observed, we set a 95% confidence level upper limit on the relative branching fraction $\Gamma(W^{\pm} \to \pi^{\pm}\gamma)/\Gamma(W^{\pm} \to e^{\pm}\nu)$ at 6.4×10^{-5}, a factor of 10 improvement over the previous limit.

PACS numbers: 14.70.Fm, 12.15.Ji, 13.38.Be

Since their discovery by the UA1 [1] and UA2 [2] ex-

*Deceased

†With visitors from [a]University of Massachusetts Amherst, Amherst, Massachusetts 01003, [b]Istituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy, [c]University of California Irvine, Irvine, CA 92697, [d]University of California Santa Barbara, Santa Barbara, CA 93106 [e]University of California Santa Cruz, Santa Cruz, CA 95064, [f]CERN,CH-1211 Geneva, Switzerland, [g]Cornell University, Ithaca, NY 14853, [h]University of Cyprus, Nicosia CY-1678, Cyprus, [i]University College Dublin, Dublin 4, Ireland, [j]University of Fukui, Fukui City, Fukui Prefecture, Japan 910-0017, [k]Universidad Iberoamericana, Mexico D.F., Mexico, [l]Iowa State University, Ames, IA 50011, [m]University of Iowa, Iowa City, IA 52242, [n]Kinki University, Higashi-Osaka City, Japan 577-8502, [o]Kansas State University, Manhattan, KS 66506, [p]University of Manchester, Manchester M13

periments, the properties and decays of the massive mediators of the electroweak interaction, the W and Z vector bosons, have been studied extensively. The exclusive leptonic final states of the boson decays, $Z \to l^+l^-$ and $W^\pm \to l^\pm\nu$, have been observed and are well-understood. Hadronic decays of the bosons have also been observed, but no exclusive final states predicted by the standard model (SM) have been identified. A class of these unobserved vector boson decay channels, radiative decays ($V \to P + \gamma$), where P is a pseudoscalar meson, are sensitive probes of the strong dynamics involved in meson formation as well as of the vector boson couplings to the photon. These radiative decays are rare because their branching ratios are suppressed by a factor of $(f_P/M_{W,Z})^2$, where f_P is the meson form factor. Standard model predictions for the branching ratio of the decay of the W boson to charged pion and photon ($W^\pm \to \pi^\pm\gamma$) range from $\sim 10^{-6}$ to $\sim 10^{-8}$ [3]. However, the sensitivity of the Tevatron experiments to observing these rare decays, especially $W^\pm \to \pi^\pm\gamma$, is approaching the upper range of the SM predictions. Measurement of the cross section of these radiative W and Z boson decays would provide information about these poorly understood SM processes, especially involving meson form factors at high energies.

Previous searches for the $W^\pm \to \pi^\pm\gamma$ decay by the UA1, UA2, and CDF experiments have successively brought the upper limit on $\Gamma(W^\pm \to \pi^\pm\gamma)/\Gamma(W^\pm \to e^\pm\nu)$ down to the level of 7×10^{-4} at 95% confidence level (C.L.)[4, 5].

In this Letter we present a search for the $W^\pm \to \pi^\pm\gamma$ decay using data corresponding to $4.3\,\mathrm{fb}^{-1}$ of integrated luminosity recorded by the CDF experiment in $p\bar{p}$ collisions at 1.96 TeV. Compared to the previous CDF search [5], this represents an increase in the size of the data set

9PL, England, [q]Queen Mary, University of London, London, E1 4NS, England, [r]University of Melbourne, Victoria 3010, Australia, [s]Muons, Inc., Batavia, IL 60510, [t]Nagasaki Institute of Applied Science, Nagasaki, Japan, [u]National Research Nuclear University, Moscow, Russia, [v]University of Notre Dame, Notre Dame, IN 46556, [w]Universidad de Oviedo, E-33007 Oviedo, Spain, [x]Texas Tech University, Lubbock, TX 79609, [y]Universidad Tecnica Federico Santa Maria, 110v Valparaiso, Chile, [z]Yarmouk University, Irbid 211-63, Jordan, [hh]On leave from J. Stefan Institute, Ljubljana, Slovenia,

by a factor of ≈ 50.

The CDF II detector is a general purpose particle detector and is described in detail elsewhere [6]. The z-axis of the detector coordinate system points along the direction of the proton beam. The event geometry and kinematics are described using the azimuthal angle ϕ and the pseudorapidity $\eta = -\ln(\tan\theta/2)$, where θ is the polar angle with respect to the beam axis. The transverse energy and momentum of the reconstructed particles are defined as $E_T = E\sin\theta$, $p_T = p\sin\theta$, where E is the energy and p is the momentum.

The analysis presented below uses electron ($e^{\pm}$), photon (γ), and charged pion ($\pi^{\pm}$) candidates, reconstructed and identified using information from the central outer tracker (COT) [7], central electromagnetic (CEM) and hadronic calorimeters (CHA), and the central shower maximum detector (CES) [8, 9].

Candidate $\pi^{\pm}\gamma$ events are collected using an inclusive photon trigger that has no tracking requirements. As a result, the trigger has similar efficiencies for selecting high-p_T electrons and photons. The kinematic properties of the $W^{\pm} \to \pi^{\pm}\gamma$ decay are also in many respects close to the kinematics properties of the $W^{\pm} \to e^{\pm}\nu$ decay, which has been extensively studied at the Tevatron [10]. Consequently, many common systematic uncertainties cancel in the ratio $\Gamma(W^{\pm} \to \pi^{\pm}\gamma)/\Gamma(W^{\pm} \to e^{\pm}\nu)$, and we use sample of $W^{\pm} \to e^{\pm}\nu$ events collected with the same trigger to normalize the results of the search.

The three level calorimeter-based trigger selects events with at least one central calorimeter cluster with $E_T >$ 25 GeV and a large fraction of energy deposited in the CEM: $E_{CHA}/E_{CEM} < 0.0055 + 0.00045 \times E$. A cluster is required to be isolated such that the additional energy in a cone of $\Delta R = \sqrt{\Delta\phi^2 + \Delta\eta^2} = 0.4$ is less than 10% of the cluster energy. To reduce background from the multi-photon decays of neutral mesons, most importantly π^0 mesons, the lateral profile of the CES shower is required to be consistent with that produced by a single photon. Offline $\pi^{\pm}\gamma$ selection requires each event passing the trigger to have a photon candidate with $E_T > 25$ GeV and a charged pion candidate with $p_T > 25$ GeV/c and with significant azimuthal separation ($\Delta\phi > 2$ radians) between them.

Photon candidates are reconstructed from CEM en-

ergy clusters with $|\eta| \leq 1.1$ and within the CES fiducial. A cluster may be pointed to by at most one track; this track must have p_T less than $1\,\mathrm{GeV/c} + 0.005 \times E_T^{\gamma}/c$. To reject background from neutral mesons decaying into photons, we veto candidates which have additional CES clusters with energy above $2.4\,\mathrm{GeV} + 0.01 \times E_T^{\gamma}$ in the same CES chamber. As photons from $W^{\pm} \to \pi^{\pm}\gamma$ decays are expected to be isolated, we require the total additional energy deposited in the calorimeter within a cone of $\Delta R = 0.4$ around the photon candidate to be less than $1.6\,\mathrm{GeV} + 0.02 \times E_T^{\gamma}$ and the sum of transverse momenta of the COT tracks within the same cone to be less than $2.0\,\mathrm{GeV/c} + 0.005 \times E_T^{\gamma}/c$.

Charged pion candidates are reconstructed as COT tracks with impact parameter ≤ 0.2 cm with respect to the beam axis. The track must be consistent with originating from the primary event vertex, which is required to be within 60 cm of the center of the detector along the beam line. The track must point to a narrow calorimeter cluster and extrapolate to a fiducial region of the CES detector. As with photons, pions from $W^{\pm} \to \pi^{\pm}\gamma$ decays are expected to be isolated, and the pion candidates are required to satisfy a number of tracking and calorimeter isolation criteria. A pion candidate is rejected if there is another track originating from the same vertex having $p_T > 1$ GeV/c within a cone of 10° around the pion track. The total additional calorimeter energy within the cone of $\Delta R = 0.4$ around the cluster is required to be less than 10% of the track p_T and the sum of transverse momenta of all tracks within $\Delta R = 0.4$ of the pion track is required to be less than 5% of its p_T. To reject background from QCD jets, which often contain neutral particles, no CES clusters with energy greater than 500 MeV may be found within 30° of the track. Finally, pion candidates passing electron identification criteria are excluded from the analysis. This selection yields a total of 1398 $\pi^{\pm}\gamma$ candidate events.

The acceptance for the $W^{\pm} \to \pi^{\pm}\gamma$ decay is determined using a sample of Monte Carlo (MC) events generated with the PYTHIA event generator [11] and the CTEQ5L parton distribution functions [12]. The angular distribution of pions in the W boson rest frame is described by the formula $dN/d\cos\theta = 1 + \cos^2\theta$, where θ is the angle of the pion with respect to the W boson line

of flight axis. This distribution describes the decay of a spin-one particle (W) into a photon and a pseudoscalar pion averaged over W boson polarizations and summed over photon polarizations. The detector response is simulated with a GEANT3 based simulation program [13].

The photon identification efficiency is determined using MC simulated events and corrected using $Z^0 \to e^+e^-$ data events, where only probe electrons with suppressed bremsstrahlung, $|E/p - 1| < 0.1$, are selected. The method assumes that the detector response to photons is the same as that to non-radiating electrons. Comparing simulations of electron and photon showers in the CEM, we verified that the accuracy of this assumption is better than 1%.

The identification efficiency for pions from the $W^\pm \to \pi^\pm\gamma$ decay is also calculated using MC simulation. The correction factor to this efficiency, which takes into account the effects of calorimeter shower mismodeling and instantaneous luminosity, is determined by comparing properties of the reconstructed jets in photon+jet data to those in photon+jet MC.

The total corrected acceptance times efficiency for $\pi^\pm\gamma$ selection is $A \times \epsilon = 0.0503 \pm 0.0006(stat) \pm 0.0011(sys)$. The uncertainty is dominated by the systematic uncertainty on the pion identification efficiency.

The dominant backgrounds to this search come from photon+jet events, where the jet fragments into a single charged particle, and from multi-jet events, where one of the jets fragments into a single charged particle and another is misidentified as a photon. Drell-Yan pair production and W/Z decays, especially to τ leptons, also contribute to the background at a level of $\approx 10\%$. The signal from the $W^\pm \to \pi^\pm\gamma$ decay would appear as a peak in the $\pi^\pm\gamma$ invariant mass spectrum centered at the W boson mass with a resolution of 2.5 GeV/c^2, which includes the experimental resolution and the full width of the W boson, 2.1 GeV/c^2 [14]. We therefore define the signal region as $75 < M_{\pi^\pm\gamma} < 85$ GeV/c^2, which includes 90% of $W^\pm \to \pi^\pm\gamma$ decays. The background within the signal region is estimated by fitting the sidebands, $67.5 < M_{\pi^\pm\gamma} < 75$ GeV/c^2 and $85 < M_{\pi^\pm\gamma} < 120$ GeV/c^2, with an exponential function using a χ^2 minimization.

Figure 1 shows the $M_{\pi^\pm\gamma}$ distribution as well as the

sideband fit for all events passing selection. The fit residuals, plotted in Figure 2, show that the exponential fit is an adequate model of the background shape with the present level of statistics. The statistical uncertainty on the fit results in a $\approx 5\%$ uncertainty on the background expectation.



FIG. 1: $M_{\pi^{\pm}\gamma}$ and background expectation for 1398 events passing full selection. The signal expectation at the 95% C.L. upper limit is included as the dashed curve. The uncertainties shown are purely statistical.

From the sideband fit, a total of 219 ± 10 events are expected in the signal region from the fit and 206 are observed. Since the data in the signal region are consistent with the expected background, we set a 95% C.L. upper limit on the $\sigma(p\bar{p} \to W) \times BR(W^{\pm} \to \pi^{\pm}\gamma)$. We divide the signal region into four 2.5 GeV bins. The limits calculated for each bin are then combined assuming that uncertainties across the bins are 100% correlated. This provides a gain in sensitivity by using information about the shape of the expected $W^{\pm} \to \pi^{\pm}\gamma$ mass peak.



FIG. 2: The fit residual divided by bin uncertainty for Figure1. The signal and sideband (SB) regions are noted with the vertical lines.

For the normalization measurement in the $W^{\pm} \to e^{\pm}\nu$ channel we select events which have a central ($|\eta| < 1.2$) electron with $E_T > 25\,\mathrm{GeV}$ and missing transverse energy $\not{E}_T > 25\,\mathrm{GeV}$ [15]. The electron reconstruction and identification algorithms used in this analysis are discussed in [10]. In addition, a standalone measurement of electron energy in the CES helps to identify electron candidates that radiate a significant fraction of their en-

ergy (due to bremsstrahlung) and improves overall electron identification efficiency. This selection results in a sample with $\leq 5\%$ background, the dominant sources of which are $W^{\pm} \to \tau^{\pm}\nu$, $Z^0 \to e^+e^-$ decays and multijet events with one of the jets misidentified as an electron. The backgrounds are subtracted using MC simulation and the resulting $\sigma \times BR(W^{\pm} \to e^{\pm}\nu)$ is found to be consistent with previous measurements [10]. The dominant source of uncertainty on the cross section is the uncertainty on the luminosity [16], which cancels in the ratio $\Gamma(W^{\pm} \to \pi^{\pm}\gamma)/\Gamma(W^{\pm} \to e^{\pm}\nu)$.

Other sources of the systematic uncertainties that cancel almost entirely in the measurement of $\Gamma(W^{\pm} \to \pi^{\pm}\gamma)/\Gamma(W^{\pm} \to e^{\pm}\nu)$ include the uncertainties on the trigger efficiency as well as the uncertainties on parton distribution functions [17] and on the W transverse momentum, which affect the experimental acceptance. The uncertainties on the electron and photon identification efficiences cancel partially but result in a negligible uncertainty on the measurement of $\Gamma(W^{\pm} \to \pi^{\pm}\gamma)/\Gamma(W^{\pm} \to e^{\pm}\nu)$. The dominant remaining sources of uncertainties are the statistical and systematic uncertainties on $A \times \epsilon$ ($\sim$ 2%) and the statistical uncertainty on the $\pi^{\pm}\gamma$ background fit ($\sim 5\%$).

The 95% C.L. upper limit on the ratio of partial widths $\Gamma(W^{\pm} \to \pi^{\pm}\gamma)/\Gamma(W^{\pm} \to e^{\pm}\nu)$, calculated using a Bayesian approach that takes into account the effect of systematic uncertainties [18], is 6.4×10^{-5}. This improves the previous world's best limit [5] by more than a factor of 10. The corresponding 95% C.L. upper limit on the branching ratio $BR(W^{\pm} \to \pi^{\pm}\gamma)$, calculated using the world average $BR(W^{\pm} \to e^{\pm}\nu) = 0.1075 \pm 0.0013$ [14], is 7.0×10^{-6}. While no $W^{\pm} \to \pi^{\pm}\gamma$ signal was observed, the sensitivity of the CDF experiment to this decay channel has reached a level comparable to theoretical predictions. With increases in the dataset size, we expect the sensitivity to improve as $1/\sqrt{L}$, where L is the integrated luminosity.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Min-

istry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korean World Class University Program, the National Research Foundation of Korea; the Science and Technology Facilities Council and the Royal Society, UK; the Institut National de Physique Nucleaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010, Spain; the Slovak R&D Agency; the Academy of Finland; and the Australian Research Council (ARC).

[1] G. Arnison *et al.*, Phys. Lett. B **122**, 103 (1983); G. Arnison *et al.*, Phys. Lett. B **126**, 398 (1983).

[2] M. Banner *et al.*, Phys. Lett. B **122**, 476(1983); P. Bagnaia *et al.*, Phys. Lett. B **129**, 130(1983).

[3] L. Arnellos, W. Marciano, and Z. Parsa, Nucl. Phys. B **196**, 378 (1982); Y. Y. Keum and X. Y. Pham, Mod. Phys. Lett. A **9**, 1545 (1994).

[4] C. Albajar *et al.* (UA1 Collaboration), Phys. Lett. B **241**, 283 (1990); J. Alitti *et al.* (UA2 Collaboration), Phys. Lett. B **277**, 203 (1992); F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **69**, 2160 (1992); F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **76**, 2852 (1996).

[5] F. Abe *et al.* (CDF Collaboration), Phys. Rev. D **58**, 031101 (1998).

[6] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 032001 (2005).

[7] T. Affolder *et al.*, Nucl. Instrum. Methods A **526**, 249 (2004).

[8] L. Balka *et al.*, Nucl. Instrum. Methods A **267**, 272 (1988); G. Apollinari, K. Goulianos, P. Melese, and M. Lindgren, Nucl. Instrum. Methods A **412** 515 (1998).

[9] S. Bertolucci *et al.*, Nucl. Instrum. Methods A **267**, 301 (1988).

[10] A. Abulencia *et al.* (CDF Collaboration), J. Phys. G: Nucl. Part. Phys. **34**, 2457 (2007).

[11] T. Sjöstrand *et al.*, Comput. Phys. Commun. **135**, 238 (2001).

[12] H. L. Lai *et al.*, Eur. Phys. J. C **12**, 375 (2000).

[13] R. Brun and F. Carminati, CERN Programming Library Long Writeup W5013 (1993).

[14] K. Nakamura *et al.* (Particle Data Group), J. Phys. G

37, 075021 (2010).

[15] Missing E_T ($\not\!\!E_T$) is defined by $\vec{\not\!\!E}_T = -\Sigma_i E_T^i \hat{n}_i$, where i is the calorimeter tower number for $|\eta| < 3.6$, and $\hat{n}_i$ is a unit vector perpendicular to the beam axis and pointing at the ith tower ($\not\!\!E_T = |\vec{\not\!\!E}_T|$).

[16] S. Klimenko, J. Konigsberg, and T.M. Liss, FERMILAB-FN-0741 (2003).

[17] A. Stump *et al.*, J. High Energy Phys. 046 (2003) 0310.

[18] J. Heinrich and L. Lyons, Annual Review of Nuclear and Particle Science, **57**, 145 (2007).