

Study of secondary neutron interactions with ^{232}Th , ^{129}I , and ^{127}I nuclei with the uranium assembly "QUINTA" at 2, 4, and 8 GeV deuteron beams of the JINR Nuclotron accelerator¹²

J. Adam^{1,2}, V.V. Chilap³, V.I. Furman¹, M.G. Kadykov¹, J. Khushvaktov¹, V.S. Pronskikh^{1,4}, A.A. Solnyshkin¹, V.I. Stegailov¹, M. Suchopar², V.M. Tsoupko-Sitnikov¹, S.I. Tyutyunnikov¹, J. Vrzalova¹, V. Wagner², L. Zavorka¹

¹*Joint Institute for Nuclear Research, Dubna, Russia.*

²*Nuclear Physics Institute ASCR PRI, Czech Republic.*

³*Center of Physical and Technical Projects "Atomenergomash", Moscow, Russia.*

⁴*Fermi National Accelerator Laboratory, Batavia IL, USA*

Abstract

The natural uranium assembly, "QUINTA", was irradiated with 2, 4, and 8 GeV deuterons. The ^{232}Th , ^{127}I , and ^{129}I samples have been exposed to secondary neutrons produced in the assembly at a 20-cm radial distance from the deuteron beam axis. The spectra of gamma rays emitted by the activated ^{232}Th , ^{127}I , and ^{129}I samples have been analyzed and several tens of product nuclei have been identified. For each of those products, neutron-induced reaction rates have been determined. The transmutation power for the ^{129}I samples is estimated. Experimental results were compared to those calculated with well-known stochastic and deterministic codes.

¹ Work supported by Fermi Research Alliance, LLC under contract No. DE-AC02-07CH11359 with the U.S. Department of Energy.

² Accepted to Applied Radiation and Isotopes, 2015.

Study of secondary neutron interactions with ^{232}Th , ^{129}I , and ^{127}I nuclei with the uranium assembly "QUINTA" at 2, 4, and 8 GeV deuteron beams of the JINR Nuclotron accelerator

J. Adam^{1,2}, V.V. Chilap³, V.I. Furman¹, M.G. Kadykov¹, J. Khushvaktov¹, V.S. Pronskikh^{1,4},
A.A. Solnyshkin¹, V.I. Stegailov¹, M. Suchopar², V.M. Tsoupko-Sitnikov¹,
S.I. Tyutyunnikov¹, J. Vrzalova¹, V. Wagner², L. Zavorka¹,

¹*Joint Institute for Nuclear Research, Dubna, Russia.*

²*Nuclear Physics Institute ASCR PRI, Czech Republic.*

³*Center of Physical and Technical Projects "Atomenergomash", Moscow, Russia.*

⁴*Fermi National Accelerator Laboratory, Batavia IL, USA*

INTRODUCTION

Interest in the international scientific community for research of this kind, is primarily concerned with the problem of transmutation of long-lived radioactive waste [1,2] and the creation of subcritical nuclear power plants with uranium-thorium cycle, controlled high-energy particle accelerators (Accelerator Driven Subcritical Systems) [3,4]. Such research is actively conducted throughout the world has been for the last two decades: PNF (Poahng) [5], n-ToF (CERN) [6], MYRRHA (Belgium) [7] and «Energy + Transmutation» setup at JINR (Dubna) [8, 9, 10, 11]. Currently working in this direction and develops a number of programs: SINQ (PSI) [12], KEK (Japan) [13], MYRRHA (Belgium), n-ToF (CERN) and a cluster of other research programs at LANL (USA) [14] – for obtaining data and developing new materials to create prototypes industrial ADS-systems.

During the past several years, such studies have been conducted and are ongoing with beams of particles Nuclotron VBLHEP JINR (Dubna) under the program Energy plus Transmutation of Radioactive Waste. This program was carried out a large number of experiments on subcritical uranium-lead target "QUINTA" [15, 16], as well as lead-graphite target "GAMMA-3" [17]. Several experiments were carried out using a solid lead target "GENERATOR" [18, 19, 20] with proton beam Phasotron DLNP JINR.

In this paper we present experimental data in comparison with the calculations obtained in the last two years in studying the interaction of secondary neutrons with

nucleus ^{232}Th , ^{129}I , ^{127}I on the "QUINTA" VBLHEP JINR on deuteron beams with energies 2, 4, 8 GeV.

STRUCTURE SETUP "QUINTA"

Uranium assembly "Quinta" is presented in Fig.1. It consists of five sections, formed in the shape of a hexahedron (aluminum containers with an inscribed diameter of 284 mm). Containers filled cylindrical rods of natural uranium metal, having a sealed aluminum shell (external dimensions: diameter 3.6 cm, length 10.4 cm, weight 1.72 kg of uranium). The ends are made of aluminum sections, 6 mm thick. The first section, the first located along the beam contains 54 uranium rod and has a through central opening 80 mm in diameter for the input beam into the target, made in order to reduce its albedo and reduce the leakage of neutrons from the target. Four subsequent sections are structurally identical and contain 61 uranium rods. Mass of uranium in one section is $61 \times 1.72 = 104.92$ kg, and the total mass of uranium entire target $298 \times 1.72 = 512.56$ kg. The fill factor of uranium 2, 3, 4 or 5 sections about 0.8, and the whole assembly of uranium ~ 0.6 .




Fig. 1. The general scheme of the setup "QUINTA"

Uranium target surrounded with lead shield thickness of 10 cm and a weight of 2545 kg with a window to enter the beam dimensions of 15 x 15 cm² (see Fig. 2). In the side wall of lead shield on opposite to the third section there is window hole in the size 15 x 5 cm² to accommodate transmutation samples. The upper part of the lead assembly is provided with a special hole for mounting and dismantling of detector probes and installation of the samples inside the uranium assemblies between sections.




Fig. 2. The general view of setup "QUINTA", where 1 – window for placement of transmutation samples, 2 - mine for mounting and dismantling of detector probes and installation of the samples inside the uranium assemblies between sections, 3 - cover lead assembly, 4 - lead assembly, 5 - input box beam 15x15 cm².

EXPERIMENT

In our experiments, transmutation samples (¹²⁷I, ¹²⁹I, ^{nat}Th, ²³³U, ²³⁵U, ^{nat}U, ²³⁷Np, ²³⁸Pu, ²³⁹Pu, and ²⁴¹Am) were placed inside the window 1 (see Fig. 2). Irradiation was carried out on three deuteron energies (2, 4 and 8 GeV). Before entering the deuteron beam at the target were installed aluminum and copper foil. It is possible to determine the integral flux of deuterons and beam shape. Used the widely known method of activation of aluminum foil in the reaction ²⁷Al(d,x)²⁴Na and copper foil in the reaction ^{nat}Cu(d,x)²⁴Na. The following Table 1 shows the irradiation conditions and the characteristics of the samples that were used in our studies (^{nat}Th, ¹²⁹I and ¹²⁷I).

Table.1. Data on the irradiation conditions and characteristics of the samples.

Deuterons energy, GeV	2		4		8	
Irradiation time, min.	376		561		970	
Integral number of deuterons	3.02(10)E+13		2.73(10)E+13		9.10(40)E+12	
Coordinates of the center of beam, cm *	Xc	Yc	Xc	Yc	Xc	Yc
	1.5(2)	0.1(1)	1.8(1)	-0.3(1)	0.9(1)	0.1(1)
FWHM (full width at half maximum), cm *	FWHM _X	FWHM _Y	FWHM _X	FWHM _Y	FWHM _X	FWHM _Y
	2.0(1)	1.7(2)	1.5(2)	1.1(1)	1.0(1)	1.3(1)
Samples	Th-nat		Th-nat		Th-nat	
Mass, g.	0.975		1.000		0.249	
Diameter of samples, cm	1.3		1.3		1.3	
Samples	I-129	I-127	I-129	I-127	I-129	I-127
Mass (I-129), g.	0.591	-	0.339	-	0.218	-
Mass (I-127), g.	0.129	1.550	0.074	1.270	0.048	1.980
Mass (Na-23), g.	0.118	0.290	0.067	0.230	0.043	0.360
Mass (Al-27), g. **	17.6	-	17.6	-	17.6	-
Diameter of samples, cm	2.1	2.0	2.1	2.0	2.1	2.0

* Deuteron beam parameters (plata0) from group I.V. Zhuk (Minsk). Determined using solid-state nuclear track detectors with radiators from a natural lead, called sensors [private message].

** Container for radioactive ¹²⁹I.

After each session of irradiation the studying samples were transported to the DLNP by complex YASNAPP-2 where measured γ -spectra with the three spectrometers based HPGe-detector ORTEC (single detector efficiency - 33% energy resolution - 1.8 keV at line 1.33 MeV ⁶⁰Co) and CANBERRA (two detector efficiency - 18% and 30%, the energy resolutions - 1.9 keV and 1.8 keV 1.33 MeV at line ⁶⁰Co). For each sample at various time intervals was measured from 13 to 16 γ -spectra and decay time to the first spectrum from 79 to 157 min. Calibration of the detectors by energy and efficiency was performed using a standard set of sources (⁵⁴Mn, ⁵⁷Co, ⁶⁰Co, ⁸⁸Y, ¹¹³Sn, ¹³³Ba, ¹³⁷Cs, ¹³⁹Ce, ¹⁵²Eu, ²²⁸Th, ²⁴¹Am).

PROCESSING OF γ - SPECTRA

Processing of the measured γ -spectra were using DEIMOS32 [21]. The program allows to define the area under the γ -peaks and their position (channel number). Next, using a special software package [22] was calibrated for energy,

corrects for detector efficiency and identified separate γ -lines corresponding nuclei products that were emitted in the sample as a result of interaction with secondary neutrons. In determining the intensities of γ -transitions also introduced corrections to the nuclear decay products during irradiation, corrections for self-absorption for registered γ -rays in the sample, to the geometric dimensions of the sample, corrections to breaks during irradiation and the change in intensity of the deuteron beam (on-line measurements of fast ionization chambers). All these procedures are described in detail in [9, 22, 23].

METHOD OF ANALYSIS AND CALCULATIONS

In determining the reaction rates (experimental data) used the following relation (1) [9]:

$$R(A_r, Z_r) = \frac{Q(A_r, Z_r)}{N_t N_d} , \quad (1)$$

where, $Q(A_r, Z_r)$ – rate of production of radioactive nucleus (r), N_t – number of atoms in the sample, N_d - number of incident deuterons on the target.

Values of reaction rates (in calculation) calculated by the formula:

$$R(A_r, Z_r) = \int_{E_{thr}(A_r, Z_r)}^{\infty} \sigma(A_r, Z_r, E_n) \varphi(E_n) dE_n \quad (2)$$

where, $\sigma(A_r, Z_r, E_n)$ – reaction cross section, $\varphi(E_n)$ – neutron fluence.

Calculations of reaction rates (Calc.1) were performed with program MARS15 [24] and the reaction products with neutrons were modeled using LAQGSM03.03 [25].

Calculation of neutron fluence (Calc.2) was carried with program MCNPX2.7 [26] using models INCL4 (intranuclear physics model) [27] and ABLA (fission-evaporation model) [28]. For the reaction (n, γ) in the ^{232}Th , the reaction cross sections were calculated by the program TALYS1.4 [29], and (n,fission) were taken from the nuclear data library TENDL-2009 [30], as in the TENDL-2009, thorium fission reaction cross sections with neutrons calculated up to neutron energy 200 MeV and are in good agreement with the data from the library JEFF3.1.2 [31], as well as with the experimental data [32] (see Figure.3). Reaction cross sections for (n, γ), (n,4n) and (n,6n) in ^{129}I to 40 MeV from TENDL-2011, from 40 to 200 MeV, the calculation was performed with program TALYS1.4. In ^{127}I reaction cross sections for (n, γ), (n,2n) and (n,4n) chosen the calculations with program TALYS1.4 lack of data in libraries up to 200 MeV.




Fig.3. Comparison of the calculated fission cross sections of ^{232}Th with experiment [32], depending on the neutron energy.

RESULTS OF ^{232}Th

In results processing of obtained data in the experiment for thorium has been identified a large number of product nuclei (at 2 GeV – 19, at 4 GeV - 30 and at 8 GeV - 27) for three values of the deuteron energy. For each of these are obtained reaction rates. Table 2 summarizes the results of ^{232}Th for all energies for all registered nuclei. I_g – the yield of gamma rays (%), $T_{1/2}(\text{Exper})$ – the observed half-lives of the radionuclides, R – reaction rate, $\langle R \rangle$ - the average value of the reaction rate ($\text{atoms}^{-1} * \text{deuteron}^{-1}$). The results show that with increasing deuteron energy increases and the value of the reaction rates for almost all product nuclei. Obviously, this growth is due to the increased flow and energy of secondary neutrons with increasing energy deuterons. Fig.4 shows the ratio of reaction rate $R(4 \text{ GeV}) / R(2 \text{ GeV})$ and $R(8 \text{ GeV}) / R(2 \text{ GeV})$ for the identified product nuclei generated in reactions with secondary neutrons at all three deuteron energies (2, 4, 8 GeV).




Fig.4. Experimental values of the ratio of reaction rate $R(4 \text{ GeV}) / R(2 \text{ GeV})$ and $R(8 \text{ GeV}) / R(2 \text{ GeV})$ for ^{232}Th with secondary neutrons for product nuclei at energies of deuterons 2, 4, 8 GeV.

Presented in Fig.4 nuclei products: $^{85\text{m}}\text{Kr}$, ^{87}Kr , ^{88}Kr , $^{91\text{m}}\text{Y}$, ^{91}Sr , ^{92}Y , ^{92}Sr , ^{93}Y , ^{97}Nb , ^{97}Zr , ^{133}I , ^{134}I , ^{135}I , ^{135}Xe , ^{138}Cs , ^{142}La and ^{143}Ce – produced by fission the ^{232}Th ; radionuclides: ^{66}Ga , ^{88}Y , $^{92\text{m}}\text{Nb}$, ^{105}Ru , ^{115}Cd , $^{115\text{m}}\text{In}$, ^{117}In , ^{126}Sb , ^{128}Sb , ^{129}Sb , ^{132}Te и ^{132}I – products of the (n,spallation) reactions (the ratio of the reaction rate to yield R / Y for these radionuclides 10-20 times differed from fission products). ^{224}Ac is product of reaction $^{228}\text{Th}(n,2nt)^{224}\text{Ac}$ ($E_{\text{thr}} = 16.56 \text{ MeV}$), ^{233}Pa produced in the reaction $^{232}\text{Th}(n,\gamma)^{233}\text{Th}$ (β^- decay, $T_{1/2}=22.3 \text{ min.}$) \rightarrow ^{233}Pa (β^- decay, $T_{1/2}=26.967 \text{ day}$) \rightarrow ^{233}U .

Fig.5 shows the experimental values of the reaction rate (n, γ) and fission (n,f) depending on the deuteron energy in comparison with the calculations Calc.2 (MCNPX). In both cases, there is a proportional increase in the values of reaction rate with increasing energy deuterons.




Fig.5. a) Comparison of experimental and calculated values of the reaction rate (n,γ) ^{232}Th in the interaction with the secondary neutrons, depending from deuteron energy, b) Comparison of experimental and calculated values of the fission reaction rate (n,f) in the interaction of ^{232}Th with secondary neutrons depending from deuteron energy.

Calculations of fission reaction rates $R(n,f)$ from the experimental data were carried out as follows. Cumulative yield Y for fission products of ^{232}Th were taken from the library JEFF3.1 for neutrons with an energy 0.4 MeV. The average value of the ratio R/Y for fission products such as nuclei ^{85m}Kr , ^{87}Kr , ^{88}Kr , ^{91m}Y , ^{91}Sr , ^{92}Y , ^{92}Sr , ^{93}Y , ^{97}Nb , ^{97}Zr , ^{133}I , ^{134}I , ^{135}I , ^{135}Xe , ^{138}Cs , ^{142}La and ^{143}Ce in experiment for deuteron energies at 2 GeV is $0.58(6)\text{E-}27$, at 4 GeV is $1.36(9)\text{E-}27$ and at 8 GeV is $2.92(14)\text{E-}27$. These numbers is the fission reaction rates $R(n,f)$ for ^{232}Th .

On the experimental results values of ratio $(n,\gamma)/(n,f)$ for different energies of deuterons: 2 GeV – $16.4(24)$, at 4 GeV – $13.0(20)$, at 8 GeV – $14.8(22)$. Calculated values (Cal.2.) for these ratios $(n,\gamma)/(n,f)$ are at 2 GeV - 12.3, at 4 GeV - 12.2 and at 8 GeV - 11.2.

Table.2. Values of the reaction rates ^{232}Th with secondary neutrons for product nuclei at energies of deuterons 2, 4, 8 GeV.
 (*) denotes mixing due to other nuclide.

Isotope	I _g [%]	2 GeV		4 GeV		8 GeV	
		T _{1/2} (Library) T _{1/2} (Exper.)	<R> R	T _{1/2} (Library) T _{1/2} (Exper.)	<R> R	T _{1/2} (Library) T _{1/2} (Exper.)	<R> R
Ga-66				9.49(7) h		9.49(7) h	
1039.231	37			12.6(20) h	2.68(62)E-29	6(5) h	1.05(23)E-28
Kr-85m		4.48(1) h		4.48(1) h			
151.159	75	7(3) d	1.47(48)E-29				
304.870	14			3.2(11) h	5.52(85)E-29		
Kr-87		76.3(6) m		76.3(6) m		76.3(6) m	
402.586	49.6	1.08(12) h	4.32(47)E-29	1.29(19) h	1.02(7)E-28	1.7(7) h	2.64(69)E-28
Kr-88		2.84(3) h	2.90(23)E-29	2.84(3) h	8.61(94)E-29	2.84(3) h	
196.301	26	2.3(4) h	2.85(25)E-29				
1529.770	10.9			3.9(1) h	1.08(11)E-28	12(5) h	3.2(11)E-28
2195.842	13.2		3.36(71)E-29		7.4(10)E-29		
2392.110	34.6	2.5(22) h	3.22(95)E-29	2.11(22) h	8.17(93)E-29		
Y-88				106.65(4) d		106.65(4) d	
898.042	93.7						
1836.063	99.2				2.99(36)E-28		1.11(18)E-27
Y-91mD		49.71(4) m		49.71(4) m		49.71(4) m	
555.570	95	10.3(12) h	2.78(68)E-29		1.61(9)E-28		1.74(27)E-28
Sr-91		9.63(5) h	4.00(90)E-29	9.63(5) h	1.01(3)E-28	9.63(5) h	1.89(29)E-28
652.900	8			14(10) h	1.15(11)E-28		
749.800	23.6	10.8(10) h	4.85(25)E-29	9.1(7) h	1.01(4)E-28	1.15(25) d	2.09(31)E-28
1024.300	33	8.8(15) h	2.82(29)E-29	9.9(4) h 7	1.01(3)E-28	9.1(16) h	1.83(26)E-28
Sr-92		2.71(1) h		2.71(1) h		2.71(1) h	
1383.930	90	2.62(10) h	3.81(16)E-29	2.68(11) h	8.81(32)E-29	2.74(27) h	1.82(13)E-28
Y-92				3.54(1) h	1.45(13)E-28	3.54(1) h	4.7(10)E-28
934.460	13.9				1.63(36)E-28		4.4(14)E-28
1405.280	4.8			5.1(12) h	1.42(14)E-28		4.9(16)E-28

Nb-92m				10.15(2) d		10.15(2) d	
934.460	99				2.43(19)E-28		3.16(69)E-28
Y-93		10.18(8) h					
266.900	7.3	9(3) h	4.28(56)E-29				
Zr-97		16.91(5) h		16.91(5) h		16.91(5) h	
743.360	93	14.7(8) h	2.88(14)E-29	15.7(6) h	6.76(18)E-29	17.2(24) h	1.26(9)E-28
Nb-97				72.1(7) m			
658.080	98				6.89(46)E-29		
Ru-105		4.44(2) h		4.44(2) h	2.66(28)E-29	4.44(2) h	
469.370	17.5				2.03(78)E-29		
724.210	47	9(6) h	1.15(11)E-29	5.7(26) h	2.74(30)E-29		8.9(22)E-29
Cd-115				53.46(1) h		53.46(1) h	
336.240	45.9						
527.900	27.5			1.7(5) d	3.88(62)E-29		1.98(65)E-28
In-115m		4.87(1) h					
336.240	45.8	4.4(20) h	8.4(15)E-30				
In-117D		43.2(3) m	1.90(34)E-29	43.2(3) m		43.2(3) m	
158.562	87		1.69(37)E-29				
553.000	100		2.12(32)E-29		7.26(95)E-29	26.5(1) m	1.04(27)E-28
Sb-126				12.46(3) d	4.81(51)E-29	12.46(3) d	3.74(52)E-28
414.810	83.3						
666.331	100				4.0(11)E-29		4.38(77)E-28
695.030	100				5.06(58)E-29		3.03(88)E-28
Sn-127						91.1(5) m	
1114.300	39					2.08(1) h	6.9(20)E-29
Sb-128		9.01(3) h		9.01(3) h		9.01(3) h	7.1(16)E-29
314.120	61						4.6(23)E-29
526.570	45			4.5(22) h	2.76(21)E-29		7.9(22)E-29
743.220	100	14.7(8) h	2.08(10)E-29		5.56(59)E-29*		8.0(14)E-29
Sb-129				4.40(1) h			
812.800	43			6(4) h	7.9(16)E-30		
Te-132				3.20(1) d		3.20(1) d	

228.160	88			15(10) d	4.64(94)E-29		1.46(38)E-28
I-132				2.29(1) h	1.49(65)E-29		
667.718	99				1.35(77)E-29		
954.550	17.6				1.81(70)E-29		
I-133		20.8(1) h		20.8(1) h		20.8(1) h	
529.872	87	1.06(15) d	2.02(14)E-29	21.2(2) h	6.49(20)E-29	1.24(15) d	1.27(9)E-28
I-134		52.5(2) m		52.5(2) m	1.69(11)E-28	52.5(2) m	2.62(27)E-28
847.025	95.4	1.15(7) h	5.75(76)E-29	1.06(2) h	1.53(17)E-28	1.33(19) h	2.51(57)E-28
884.090	64.9			1.11(7) h	1.59(18)E-28		2.57(32)E-28
1072.550	14.9			1.12(2) h	1.94(21)E-28		3.4(13)E-28
1136.160	9.1				1.98(33)E-28		
I-135		6.57(2) h	3.09(14)E-29	6.57(2) h	7.04(26)E-29	6.57(2) h	1.72(28)E-28
1131.511	22.7	6.8(8) h	3.11(21)E-29	5.7(6) h	6.99(52)E-29	12.9(1) h	1.77(26)E-28
1260.409	28.9	7(5) h	3.08(18)E-29	6.7(4) h	6.89(28)E-29	7(2) h	1.67(73)E-28
1791.196	7.8			8.1(16) h	7.29(70)E-29		
Xe-135		9.14(2) h		9.14(2) h		9.14(2) h	
249.770	90	16.8(21) h	2.85(55)E-29	16.8(19) h	7.8(12)E-29	20(3) h	1.95(34)E-28
Cs-138				33.4(2) m		33.4(2) m	
1435.795	76.3				1.31(13)E-28		2.03(43)E-28
La-142		91.1(5) m		91.1(5) m	7.73(82)E-29	91.1(5) m	
641.285	47	1.86(19) h	3.91(31)E-29	1.9(5) h	8.4(13)E-29	2.08(1) h	1.66(24)E-28
894.900	8.3			3.46(2) h	7.3(14)E-29		
2397.800	13.3				7.3(15)E-29		
Ce-143		1.38(2) d		1.38(2) d		1.38(2) d	
293.266	42.8	1.68(3) d	3.32(22)E-29	1.50(18) d	7.80(35)E-29	3.1(7) d	2.51(35)E-28
At-208				1.63(3) h		1.63(3) h	
177.595	48.6						6.1(23)E-28
845.044	19.7				3.83(64)E-29		
Ac-224				2.78(17) h		2.78(17) h	
131.613	26.9						1.13(51)E-26
215.983	52.3				2.84(80)E-29		
Pa-233		26.97(1) d		26.97(1) d		26.97(1) d	

312.170	38.6	17.8(16) d	9.55(55)E-27		1.78(3)E-26		4.30(20)E-26
---------	------	------------	---------------------	--	--------------------	--	---------------------

RESULTS OF ^{129}I

During irradiation the samples ^{129}I (coated aluminum weighing 17.6 g) and ^{127}I (in a shell made of plexiglass weighing 2.53 g) were installed on the side of the section №3 uranium assembly Fig.2. In samples ^{129}I impurity present stable isotope ^{127}I . To correct account the contribution of ^{127}I , samples ^{129}I were irradiated simultaneously with the samples, containing only isotope ^{127}I .




Fig.6. Experimental values of the ratio of reaction rate $R(4 \text{ GeV}) / R(2 \text{ GeV})$ and $R(8 \text{ GeV}) / R(2 \text{ GeV})$ for $^{23}\text{Na} + ^{27}\text{Al} + ^{129}\text{I}$ with secondary neutrons for product nuclei at energies of deuterons 2, 4, 8 GeV. Na-22* - product $^{23}\text{Na}(n,2n)^{22}\text{Na}$. Na-22** - product $^{27}\text{Al}(n,\alpha 2n)^{22}\text{Na}$. Na-24* - product $^{23}\text{Na}(n,\gamma)^{24}\text{Na}$. Na-24** - product $^{27}\text{Al}(n,\alpha)^{24}\text{Na}$. Values of reaction rates for ^{82}Br obtained from calculation Calc.2 due to lack of weight values ^{81}Br .

^{22}Na produced simultaneously in two reactions $^{27}\text{Al}(n,\alpha 2n)^{22}\text{Na}$ ($E_{\text{thr}}=23.35 \text{ M}\text{eV}$) and $^{23}\text{Na}(n,2n)^{22}\text{Na}$ ($E_{\text{thr}}=12.96 \text{ M}\text{eV}$). ^{24}Na is generated from $^{27}\text{Al}(n,\alpha)^{24}\text{Na}$ ($E_{\text{thr}}=3.25 \text{ M}\text{eV}$) and $^{23}\text{Na}(n,\gamma)^{24}\text{Na}$ reactions. Share ^{24}Na generated from the (n, γ) reaction at deuteron energies 2 GeV - 2.1 %, 4 GeV - 0.9 % and 8 GeV - 0.6 %.

^{22}Na and ^{24}Na produced mainly from ^{27}Al due to large mass of ^{27}Al . Contribution ^{24}Na produced from ^{23}Na calculated using the values of reaction rates for ^{24}Na in the sample ^{127}I .

We assume that in the composition of samples ^{129}I , present admixture of ^{81}Br and ^{82}Br is a product of $^{81}\text{Br}(n,\gamma)^{82}\text{Br}$ reaction. The admixture of ^{81}Br may be, by our estimates (Calc.2), in ^{129}I no more than 1.5(5)%. ^{82}Br in the samples ^{127}I was not observed. ^{123}I , ^{124}I , and ^{126}I are products of (n,7n), (n,6n) and (n,4n) reactions. ^{130}I is product (n, γ) reaction.




Fig.7. Experimental values of the ratio of reaction rate $R(4 \text{ GeV}) / R(2 \text{ GeV})$ and $R(8 \text{ GeV}) / R(2 \text{ GeV})$ for $^{23}\text{Na} + ^{127}\text{I}$ with secondary neutrons for product nuclei at energies of deuterons 2, 4, 8 GeV.

^7Be is produced in the shell of the sample (plexiglass) from ^{12}C and ^{16}O . ^{22}Na and ^{24}Na produced from ^{23}Na in the reactions $(n,2n)$ and (n,γ) . $^{118\text{m}}\text{Sb}$, $^{120\text{m}}\text{Sb}$ and ^{122}Sb are products of $(n,\alpha 6n)$ ($E_{\text{thr}}=44.11 \text{ MeV}$), $(n,\alpha 4n)$ ($E_{\text{thr}}=27.42 \text{ MeV}$) and $(n,\alpha 2n)$ ($E_{\text{thr}}=11.23 \text{ MeV}$) reactions. Radionuclides ^{119}Te , ^{121}Te and $^{123\text{m}}\text{Te}$ are products of $(n,t6n)$ ($E_{\text{thr}}=57.56 \text{ MeV}$), $(n,t4n)$ ($E_{\text{thr}}=39.92 \text{ MeV}$) and $(n,t2n)$ ($E_{\text{thr}}=23.01 \text{ MeV}$). ^{120}I , ^{121}I , ^{123}I , ^{124}I , and ^{126}I are products of $(n,8n)$ ($E_{\text{thr}}=62.18 \text{ MeV}$), $(n,7n)$ ($E_{\text{thr}}=51.53 \text{ MeV}$), $(n,5n)$ ($E_{\text{thr}}=33.59 \text{ MeV}$), $(n,4n)$ ($E_{\text{thr}}=26.01 \text{ MeV}$) and $(n,2n)$ ($E_{\text{thr}}=44.11 \text{ MeV}$) reactions. ^{128}I is product (n,γ) reaction. Table 2 summarizes the results of comparing the experimental and calculated data (Calc.1 and Calc.2) on ^{127}I and ^{129}I .

Table.2. Comparison of results for ^{127}I and ^{129}I with calculations.

Nuclear reactions on ^{127}I samples	2 GeV		4 GeV		8 GeV	
	Exp/Calc. 1	Exp/Calc. 2	Exp/Calc. 1	Exp/Calc. 2	Exp/Calc. 1	Exp/Calc. 2
$^{127}\text{I}(n,\gamma)^{128}\text{I}$		0.64(4)		0.52(3)		0.53(3)
$^{127}\text{I}(n,2n)^{126}\text{I}$	1.10(4)	0.66(2)	0.85(3)	0.57(2)	0.69(3)	0.53(1)
$^{127}\text{I}(n,4n)^{124}\text{I}$	1.18(9)	0.68(2)	0.89(8)	0.64(7)	0.79(7)	0.62(4)
Nuclear reactions on ^{129}I samples						
$^{129}\text{I}(n,\gamma)^{130}\text{I}$		0.61(1)		0.71(1)		0.71(1)
$^{129}\text{I}(n,4n)^{126}\text{I}$	1.45(12)	0.72(3)	2.46(17)	1.56(6)	1.74(27)	1.27(6)

$^{129}\text{I}(n,6n)^{124}\text{I}$	1.95(26)	0.88(5)	2.49(32)	1.49(19)	1.43(23)	0.94(18)
--------------------------------------	----------	---------	----------	----------	----------	----------




Fig.8. Comparison of the experimental results on ^{127}I - a) and ^{129}I - b) with [18].

Previously, we performed an experiment [18], in which for production of secondary neutrons used massive lead assembly installed in the proton beam (660 MeV) in Phasotron DLNP. Data for ^{127}I and ^{129}I , obtained in this experiment, we compare with data from experiments at the Nuclotron VBLHEP (see Fig.8). It can be seen that the results for ^{127}I and ^{129}I in good agreement for almost all the observed (n,xn) reactions (in this work have not been identified $(n, 9n)$, $(n, 10n)$ reactions in

^{127}I , $(n, 9n)$, $(n, 10n)$ and $(n, 11n)$ reactions in ^{129}I , as the time from end of irradiation up to start of measurement 80 -160 min.), except the (n,γ) reaction, which is caused by the presence on setup "QUINTA" lead shield that reflects neutrons.

Table.3. Values of the reaction rates ^{127}I with secondary neutrons for product nuclei at energies of deuterons 2, 4, 8 GeV.
 (*) denotes mixing due to other nuclide.

Isotope Energy [keV]	I _g [%]	2 GeV		4 GeV		8 GeV	
		T _{1/2} (Library) T _{1/2} (Exper.)	<R> R	T _{1/2} (Library) T _{1/2} (Exper.)	<R> R	T _{1/2} (Library) T _{1/2} (Exper.)	<R> R
Be-7		53.12(7) d		53.12(7) d		53.12(7) d	
477.595	10.5	38(15) d	3.68(30)E-29	25(12) d	6.81(47)E-29		1.04(9)E-28
Na-22		2.60(1) y		2.60(1) y		2.60(1) y	
1274.530	99.94	100(110) d	5.58(51)E-29		1.01(24)E-28	7.0(15) d	2.79(75)E-28
Na-24		14.96(1) h	7.22(66)E-29	14.96(1) h	9.6(11)E-29	14.96(1) h	1.85(26)E-28
1368.633	100	15.18(18) h	7.04(15)E-29	15.21(10) h	9.40(24)E-29	15.17(14) h	1.79(5)E-28
2754.028	99.94	15.31(21) h	9.65(58)E-29	14.86(10) h	1.39(9)E-28	14.76(19) h	2.93(19)E-28
Ag-108m						418(21) y	2.70(63)E-30
433.937	90						2.57(73)E-30
614.276	89.8						2.85(76)E-30
722.907	90.8						6.9(13)E-30*
In-111		2.80(1) d	5.87(72)E-31				
171.280	90	2.8(4) d	5.26(40)E-31				
245.395	94	2.5(4) d	6.73(47)E-31				
Sb-118m		5.00(2) h	1.55(6)E-30	5.00(2) h	1.54(11)E-30	5.00(2) h	4.01(30)E-30
253.678	99	5.09(25) h	1.47(7)E-30	5.1(4) h	1.44(9)E-30	5.0(4) h	3.44(21)E-30
1050.650	97	5.8(7) h	1.64(11)E-30	4.6(7) h	1.76(16)E-30	4.7(4) h	4.51(35)E-30
1229.680	100	5.1(4) h	1.62(10)E-30	8.3(14) h	1.73(23)E-30	6.2(8) h	4.26(41)E-30
Te-119		16.03(5) h		16.03(5) h		16.03(5) h	
644.010	84	19.0(6) h	3.24(14)E-30	16.9(7) h	3.92(13)E-30	15.5(6) h	1.10(4)E-29
Te-119m		4.70(4) d	2.49(16)E-30	4.70(4) d	4.09(23)E-30	4.70(4) d	1.05(4)E-29
153.590	66	4.9(4) d	2.39(11)E-30	4.1(6) d	4.20(36)E-30	3.92(23) d	1.02(5)E-29
1212.730	66	5.4(7) d	2.75(17)E-30	11.3(28) d	4.02(30)E-30	4.6(8) d	1.11(8)E-29
I-120				81.0(6) m		81.0(6) m	
560.440	73			1.32(2) h	3.87(36)E-30	1.40(14) h	9.3(8)E-30
Sb-120m		5.76(2) d	1.58(8)E-30	5.76(2) d	1.61(19)E-30	5.76(2) d	4.43(24)E-30

197.300	87	12.3(21) d	1.50(13)E-30	5.5(7) d	1.51(21)E-30	5.6(5) d	3.83(33)E-30
1023.100	99.4	9(4) d	1.63(16)E-30	1.6(10) d	1.86(57)E-30	5.4(5) d	4.44(33)E-30
1171.300	100	5.4(13) d	1.63(12)E-30			5.0(15) d	4.52(92)E-30
I-121		2.12(1) h		2.12(1) h		2.12(1) h	
212.189	84	2.14(9) h	9.48(73)E-30	2.05(10) h	1.07(7)E-29	1.99(9) h	2.71(21)E-29
Te-121m		154(7) d		154(7) d		154(7) d	
212.189	81		8.7(16)E-30		1.01(20)E-29		2.33(42)E-29
Te-121		16.78(35) d	1.26(26)E-29	16.78(35) d		16.78(35) d	3.47(20)E-29
507.591	17.7	23(8) d	2.37(22)E-29			13(5) d	6.23(73)E-29*
573.139	80.3	22.7(10) d	1.20(5)E-29	19.3(13) d	1.60(11)E-29	16.6(19) d	3.47(20)E-29
Sb-122		2.72(1) d		2.72(1) d		2.72(1) d	
564.119	71	2.64(23) d	2.29(11)E-30	2.4(3) d	3.08(16)E-30	2.89(22) d	7.19(36)E-30
I-123		13.27(8) h		13.27(8) h		13.27(8) h	
158.970	83	13.0(6) h	3.59(41)E-29	12.3(6) h	4.18(45)E-29	12.5(7) h	9.5(11)E-29
Te-123m		119.7(1) d		119.7(1) d		119.7(1) d	
158.970	84		7.7(13)E-30		1.55(21)E-29		3.47(47)E-29
I-124		4.18(2) d	5.54(18)E-29	4.18(2) d	7.93(85)E-29	4.18(2) d	1.71(12)E-28
602.729	63	4.18(15) d	5.56(15)E-29	4.13(5) d	7.29(30)E-29	3.96(9) d	1.56(5)E-28
722.786	10.3	4.14(22) d	5.26(17)E-29	17(7) d	1.21(10)E-28	5.1(11) d	2.44(21)E-28
1325.512	1.6	3.4(10) d	7.40(81)E-29	4.8(28) d	1.03(10)E-28	3.4(3) d	2.46(20)E-28
1509.470	3.1			5.5(7) d	6.9(12)E-29	4.5(5) d	1.97(22)E-28
1690.983	10.9	4.6(4) d	5.84(23)E-29	4.9(3) d	7.02(39)E-29	4.07(17) d	1.63(6)E-28
Xe-125		16.9(2) h	7.93(70)E-31	16.9(2) h	1.31(10)E-30	16.9(2) h	2.45(22)E-30
188.418	54	17.1(26) h	7.57(61)E-31	15.3(23) h	1.28(10)E-30	20(6) h	2.10(20)E-30
243.378	30	1.8(5) d	9.3(12)E-31		1.61(30)E-30		2.89(63)E-30
I-126		13.11(5) d	2.32(7)E-28	13.11(5) d	3.35(12)E-28	13.11(5) d	6.90(18)E-28
388.633	34.1	13.3(6) d	2.49(5)E-28	12.37(16) d	3.44(16)E-28	10.7(5) d	6.85(22)E-28
491.243	2.8	12.8(6) d	2.33(8)E-28	11.8(6) d	3.27(18)E-28	11.1(11) d	6.44(28)E-28
666.331	33.1	13.8(7) d	2.27(5)E-28	11.98(29) d	3.36(11)E-28	10.8(6) d	6.73(16)E-28
753.819	4.2	14.0(7) d	2.16(6)E-28	11.5(6) d	3.14(17)E-28	11.0(11) d	6.53(25)E-28
879.876	0.7	15(4) d	2.57(29)E-28	12.8(18) d	4.74(48)E-28	9.1(12) d	9.68(97)E-28*
I-128		24.99(2) m		24.99(2) m	1.95(12)E-26	24.99(2) m	3.38(17)E-26

442.901	17	25.2(1) m	1.30(8)E-26	25.06(2) m	1.97(12)E-26	24.90(2) m	3.45(6)E-26
526.557	1.6			31.1(1) m	1.87(26)E-26	24.68(2) m	3.29(8)E-26

Table.4. Values of the reaction rates ^{129}I with secondary neutrons for product nuclei at energies of deuterons 2, 4, 8 GeV.

(*) denotes mixing due to other nuclide.

Isotope Energy [keV]	I_g [%]	2 GeV		4 GeV		8 GeV	
		$T_{1/2}(\text{Library})$ $T_{1/2}(\text{Exper.})$	$\langle R \rangle$ R	$T_{1/2}(\text{Library})$ $T_{1/2}(\text{Exper.})$	$\langle R \rangle$ R	$T_{1/2}(\text{Library})$ $T_{1/2}(\text{Exper.})$	$\langle R \rangle$ R
Na-22*		2.60(1) y		2.60(1) y		2.60(1) y	
1274.530	99.94	40(24) d	5.6(34)E-29	40(24) d	1.01(23)E-28		2.79(20)E-28
Na-22**		2.60(1) y		2.60(1) y		2.60(1) y	
1274.530	99.94	40(24) d	8.1(49)E-30	40(24) d	5.7(13)E-30		6.42(46)E-28
Na-24*		14.96(1) h	7.22(43)E-29	14.96(1) h	9.60(79)E-29	14.96(1) h	1.85(14)E-28
1368.633	100	15.02(3) h		15.6(4) h		15.0(4) h	
2754.028	99.94	15.05(3) h		15.6(5) h		14.9(5) h	
Na-24**		14.96(1) h	2.72(16)E-29	14.96(1) h	4.83(40)E-29	14.96(1) h	9.47(74)E-29
1368.633	100	15.02(3) h		15.6(4) h		15.0(4) h	
2754.028	99.94	15.05(3) h		15.6(5) h		14.9(5) h	
Br-82		35.30(2) h	5.09(13)E-29	35.30(2) h	6.37(12)E-28	35.30(2) h	1.11(4)E-27
554.348	70.8	1.08(5) d*	6.90(77)E-29	1.45(5) d	6.82(20)E-28	1.25(5) d	1.17(10)E-27
619.106	43.4	1.42(3) d	4.70(12)E-29	1.58(6) d	5.94(22)E-28	1.33(5) d	1.02(6)E-27
698.374	28.5	1.44(4) d	4.64(14)E-29	1.68(7) d	6.06(30)E-28	1.74(20) d	1.35(8)E-27
776.517	83.5	1.45(2) d	5.02(9)E-29	1.56(6) d	6.02(21)E-28	1.39(4) d	1.09(5)E-27
827.828	24	1.45(5) d	5.43(17)E-29	1.53(7) d	6.30(24)E-28	1.41(2) d	0.99(10)E-27
1044.002	27.2	1.48(5) d	5.31(18)E-29	1.52(6) d	6.55(25)E-28	1.7(3) d	1.13(11)E-27
1317.473	26.5	1.54(5) d	5.61(17)E-29	1.61(7) d	6.59(28)E-28	1.32(11) d	1.15(9)E-27
1474.880	16.3	1.45(6) d	5.39(20)E-29	1.58(6) d	6.56(26)E-28	1.36(12) d	1.11(8)E-27
I-123		13.27(8) h		13.27(8) h			
158.970	83	12.3(6) h	1.26(16)E-29	10(3) h	3.30(51)E-29		
I-124		4.18(2) d	2.42(15)E-29	4.18(2) d		4.18(2) d	

602.729	63	3.9(7) d	2.32(16)E-29	5.9(13) d	6.37(81)E-29	5.0(17) d	9.1(17)E-29
1690.983	10.9	4.1(19) d	2.81(33)E-29				
I-126		13.11(5) d	6.42(28)E-29	13.11(5) d	2.19(9)E-28	13.11(5) d	3.92(20)E-28
388.633	34.1	3.8(12) y	6.66(27)E-29	27(12) d	2.23(10)E-28		4.13(25)E-28
666.331	33.1	4.4(19) y	6.05(30)E-29	26(12) d	2.14(10)E-28		3.68(28)E-28
753.819	4.2	17(7) d	4.54(56)E-29				
I-130		12.36(3) h	4.59(10)E-27	12.36(3) h	1.07(2)E-26	12.36(3) h	1.84(4)E-26
418.010	34.2	12.47(2) h	4.15(7)E-27	12.8(4) h	1.01(4)E-26	12.27(29) h	1.76(6)E-26
457.720	0.2	10.1(18) h	3.98(39)E-27				
536.090	99	12.44(2) h	4.48(7)E-27	12.8(4) h	1.05(4)E-26	12.2(4) h	1.81(6)E-26
539.100	1.4	12.49(17) h	4.48(11)E-27	12.43(27) h	1.12(3)E-26	13.2(9) h	1.93(13)E-26
553.900	0.7	1.08(5) d*	8.31(8)E-27*				
586.050	1.7	12.51(12) h	4.34(10)E-27	12.9(5) h	1.03(4)E-26	12.5(7) h	1.70(8)E-26
603.500	0.6	13.8(5) h	5.35(20)E-27	13.4(18) h	1.13(9)E-26	2.37(15) d*	2.12(22)E-26
668.540	96	12.45(2) h	4.48(7)E-27	12.9(4) h	1.05(4)E-26	12.3(5) h	1.79(6)E-26
685.990	1.1	12.53(16) h	4.33(10)E-27	13.1(5) h	0.98(4)E-26	12.9(18) h	1.91(13)E-26
739.480	82	12.42(2) h	4.43(7)E-27	12.9(4) h	1.06(4)E-26	12.1(4) h	1.80(6)E-26
800.230	0.1	11.6(19) h	4.17(43)E-27				
808.290	0.2	13.0(6) h	4.22(20)E-27	14.9(20) h	0.92(8)E-26		
877.350	0.2	11.5(10) h	4.93(36)E-27				
967.020	0.9	12.47(24) h	4.51(11)E-27	13.3(8) h	1.08(5)E-26		2.21(24)E-26
1096.480	0.5	1.5(5) d	4.80(9)E-27	13.7(11) h	1.21(7)E-26		2.34(35)E-26
1122.150	0.2	12.7(10) h	5.60(30)E-27	16.4(25) h	1.44(19)E-26		
1157.470	11.3	12.37(4) h	5.31(10)E-27	13.1(5) h	1.11(4)E-26	11.8(4) h	1.97(6)E-26
1222.560	0.2	12.5(8) h	5.54(27)E-27				
1272.120	0.7	12.65(24) h	5.67(13)E-27	13.6(7) h	1.24(7)E-26	10(4) h	2.45(30)E-26
1403.900	0.3	12.3(3) h	5.08(16)E-27	14.1(20) h	1.08(12)E-26		

CONCLUSIONS

Is interesting to compare the results for ^{232}Th of this work with the results of other experiments. In [33] on setup «GAMMA-2»: Pb target ($d = 8$ cm, $l = 20$ cm), paraffin moderator (6 cm thick). ^{232}Th sample located on the end surface of the target. Installation was irradiated by protons with an energy of 1 GeV. The reaction rate (n,γ) $R = 9.80(6)\text{E-}27$, almost equal to the results of this work at a deuteron energy 2 GeV. In [9] on the setup "Energy plus Transmutation": overall length 48 cm (4 sections), the length of Pb target in four sections of 45.6 cm and a diameter of 8.4 cm target surrounded by uranium blanket thickness of two elements (see present work) and irradiated with deuterons 1.6 GeV. Samples located on the surface of the blanket. The reaction rate (n,γ) $R = 3.03(10)\text{E-}26$ and fission (n,f) $R = 5.89(60)\text{E-}27$ - 3 and 10 times higher than results of this work at 2 GeV. Also interesting comparison with the work [17], made at a deuteron energy of 2.33 GeV at the setup "GAMMA-3": a lead target ($d = 8$ cm, $l = 60$ cm) in the carbon moderator with large size, channels for samples at different distances from the center of the target. Results for $R - (n,\gamma)$ in a yield of ^{233}Pa (will eventually ^{233}U) is 30-40 times higher and fission (n, f) - 20 - 30 times higher than present work for the nearest window on carbon (24 cm from the axis target). This is due to the high ability of carbon to slowing of spallation neutrons coming from the target.

Ratio of the weight of produced ^{233}U to ^{232}Th at 2 GeV - $2.90(17)\text{E-}13$ at 4 GeV $4.88(8)\text{E-}13$ and at 8 GeV $3.93(18)\text{E-}13$. Ratio of the weight of produced ^{130}I to ^{129}I in the conditions of the present experiment at 2 GeV $1.40(3)\text{E-}13$ at 4 GeV $2.94(6)\text{E-}13$ and at 8 GeV $1.69(4)\text{E-}13$. If we calculate of transmutation for conditions: 10 mA and irradiate for 30 days. It is: 2 GeV - $0.082(2)\%$; 4 GeV - $0.190(4)\%$ and 8 GeV - $0.330(7)\%$. These calculations allow us to estimate the transmutation at high currents and irradiation time - in the tens of percent. By Calc.1 and experimental data $90(5)\%$ of the transmutation ^{129}I is (n,γ) reaction at all three deuteron energies (2, 4, 8 GeV).

These results are interesting to compare with those obtained previously. The first work was done with ^{129}I in the direct beam of protons Phasotron JINR $E_p = 660$ MeV and a current of 1.2 mA [34]. Samples with ^{129}I and ^{127}I were in direct proton beam. The paper presents number: 30 days and 10 mA. However, during its obtaining were taken coefficient "2" - taking into account the formation of stable and long-lived isotopes, as well as registration of gamma rays with energies lower than 300 keV (samples ^{129}I are measured with a filter: Pb - 10 mm, Cd - 2 mm and Cu - 1 mm). Therefore, comparisons should take a number (2-3)%. It should be borne in

mind that the experimental conditions: the direct beam of protons and not heavy target surrounded uranium, carbon or paraffin moderator - significantly different from the next. In experiments with beams of relativistic protons nuclotron JINR VBLHEP with energies: 1.0, 1.5 and 2.0 GeV [10] - transmutation is respectively, - 0.075, 0.132 and 0.153% at a current of 10 mA and 30 days of irradiation, which is close to the results of this work. In a similar study with deuterons 2.52 GeV [8], a transmutation of 0.132%. In [11] the results for protons with energy 2.0 GeV - transmutation is 0.13%. In [35, 36] at the proton beam with the energy of 3.67 GeV (Nuclotron JINR), lead target (d = 8 cm, l = 20cm, paraffin moderator) transmutation of ^{129}I was 0.9% (10 mA, 30 days). Grows of transmutation is associated from energy, obviously with increase the yield of spallation neutrons from the target ($70 \pm 20\%$) and including - thermal, making the main contribution to the reaction (n, γ) - more than 92%.

REFERENCES

1. C.D. Bowman et al. // Nucl. Instr. Meth. A. 1992. V.320. P.336.
2. A.J. Janssen. Transmutation of fission products in reactors and Accelerator-Driven Systems, ECN-R-94-001. 2004.
3. C. Rubbia, J.A. Rubio et al. Conceptual design of a fast neutron operated high power energy amplifier CERN/AT/95-44 (ET).1995. See also C. Rubbia. High gain energy amplifier operated with fast neutrons. AIP Conference 346, Proceedings of International Conference on Accelerator-Driven Transmutation Technologies and Applications, Las Vegas (NV), US.1994.
4. Thorium Fuel Cycle – Potential Benefits and Challenges, IAEA-TECDOC-1450 (2005).
5. G.N. Kim et al. // Nucl. Instr. Meth. A. 2002. V.485. P.458.
6. <http://pceet075.cern.ch/>
7. <http://www.sckcen.be/myrrha/>
8. M.I. Krivopustov, A.V. Pavlyuk, A.I. Malakhov et al. About the first experiment on investigation of ^{129}I , ^{237}Np , ^{238}Pu and ^{239}Pu transmutation at the Nuclotron 2.52 GeV deuteron beam in neutron field generated in U/Pb assembly “Energy plus Transmutation”. Journal of Radioanalytical and Nuclear Chemistry, Vol. 279, No.2 (2009) 567–584 // Preprint JINR E1-2007-7. 2007. Dubna.
9. J. Adam, K. Katovsky, M. Majerle et al. Transmutation of Th and U with neutrons produced in Pb target and U-blanket system by relativistic deuterons. Eur. Phys. J. A 43. No2. pp. 159-173, 2010. // Preprint JINR, E15-2008-118. Dubna, 2008, 29 p.
10. J. Adam, K. Katovsky, A. Balabekyan et al. Transmutation of ^{129}I , ^{237}Np , ^{238}Pu , ^{239}Pu and ^{241}Am Using Neutrons Produced in Target-Blanket System “Energy plus Transmutation” by relativistic Protons. Pramana Journal of Physics. Vol.68, N 2 (2007). PP. 201-212.
11. J. Adam, K. Katovsky A. R. Balabekyan et al. Transmutation of ^{129}I , ^{237}Np , ^{239}Pu and ^{241}Am Using Neutrons Produced in Pb-Target with ^{nat}U Blanket System “Energy plus Transmutation” Bombarded by 2.0 GeV Relativistic Protons.

- Proc. of the International Conference on Nuclear Data and technology. September 26-October 1, 2004, Santa Fe, New Mexico, USA. N.Y., 2005. V.2. P 1560-1562.
12. <http://sinq.web.psi.ch/>
 13. <http://www.kek.jp/intra-e/index.html>
 14. <http://lansce.lanl.gov/indexC.html>
 15. Коллаборация «Энергия и трансмутация радиоактивных отходов». Исследование пространственных распределений реакций деления и радиационного захвата нейтронов в массивной урановой мишени, облучаемой дейтронами с энергией 1-8 ГэВ (установка «Квинта»). Препринт ОИЯИ, P1-2012-147, Дубна, 2012, 20 с.
 16. N.L. Asquith, S.R. Hashemi-Nezhad, S. Tyutyunnikov, M. Kadykov, V. Chilap, J. Adam, W. Furman. *Annals of Nuclear Energy*. Vol. 63, 2014, P. 742–750
 17. J. Adam et al. A study of rates of (n,f), (n, γ), and (n,2n) reactions in ^{nat}U and ^{232}Th produced by the neutron fluence in the graphite setup (GAMMA-3) irradiated by 2.33 GeV. *Eur. Phys. J. A* 47, 85-104 (2011). (Preprint JINR. E1-2011-52, Dubna 2011)
 18. M. Majerle et al. Spallation experiment on thick lead target: analysis of experimental data with Monte Carlo codes. Preprint JINR, E15-2008-94. Dubna, 2008, 20 p.
 19. J. Adam et al. Investigation of the formation of residual nuclei in reactions induced by 660 MeV protons interacting with the radioactive ^{237}Np , ^{241}Am and ^{129}I targets. *Journal of Nuclear Science and Technology, Supplement 2*, p. 272-275, 2002.
 20. V. Henzl et al. Transmutation of ^{129}I with high energy neutrons produced in spallation reactions induced by protons in massive target. *Journal of Nuclear Science and Technology, Supplement 2*, p. 1248-1251, 2002.
 21. J. Frana. Program DEIMOS32 for gamma-ray spectra evaluation. *J. Radioanal. Nucl. Chem.* 2003, V. 257, 583.
 22. И. Адам и др. Система программ и дополнения к методу активационного анализа для определения сечений ядерных реакций. *Измерительная техника*. №1, 2001, 57-61. //Препринт ОИЯИ, P10-2000-28, Дубна, 2000, 22 с.
 23. И. Адам и др. Исследование образования продуктов протон-ядерных реакций в мишени ^{129}I при энергии протонов 660 МэВ. *Письма в ЭЧАЯ*. 2004. Т. 1, № 4(121). С. 53-64.
 24. N.V. Mokhov, C.C. James. *The MARS Code System User's Guide: Version 15*. Fermi National Accelerator Laboratory. P.O. Box 500, Batavia, Illinois 60510, January, 2014.
 25. S.G. Mashnik, et al., LANL Report LA-UR-08-2931, 2008.
 26. D.P. Pelowitz, *MCNPX User's Manual: Version 2.5.0*, Technical Report, LA-CP- 05-0369, Los Alamos National Laboratory and University of California, April 2005.
 27. A. Boudard, et al., *Physical Review C* 66 (2002) 044615.
 28. A.R. Junghans, et al., *Nuclear Physics A* 629 (1998) 635.
 29. A. J. Koning, *TALYS User Manual: Version 1.4*, Technical Report, NL-1755 ZG, Petten, The Netherlands, December 2011.
 30. TENDL-2009, Nuclear data library.
 31. JEFF3.1.2, Nuclear data library.
 32. O.A. Shcherbakov, A.Y. Donets, A.V. Evdokimov, A.V. Fomichev, T. Fukahori, A. Hasegawa, A.B. Laptev, V.M. Maslov, G.A. Petrov, Y.V. Tuboltsev, A.S. Vorobiev, *Journal of Nuclear Science and Technology Supplement 2 (1)* (2002) 230.
 33. J. Adam, A.R. Balabekyan, V.S. Barashenkov et al. Spallation neutron spectrum on a massive lead/paraffin target irradiated with 1 GeV protons. *Eur. Phys. J. A* 23, 61-68 (2005). // Preprint JINR E1-2004-16. 13 pp. Dubna. 2004.

34. И. Адам, А. Балабекян, В.С. Барашенков и др. Исследование образования продуктов протон-ядерных реакций в мишени ^{129}I при энергии протонов 660 МэВ. Труды 51 междунар. сов. по ядерной спектроскопии и структ. атомного ядра. Саратов, Россия, 2001, с.201. // Письма в ЭЧАЯ. 2004. т.1, № 4 (121).С.53-64.
35. M.I. Krivopustov, J. Adam, V. Bradnova et al. First Experiment on Transmutation Studies of Iodine - 129 and Neptunium -237 Using Relativistic Protons of 3.7 GeV. Preprint JINR, E1-97-59 (1997). Dubna.// J. of Radioanalytical and Nuclear Chemistry. Vol.222 (1997).P.267
36. J. S. Wan, M. Ochs, R. Brandt et al. Transmutation of radioactive waste by means of relativistic heavy ions. Kerntechnik 63 (1998) 4. // Communication JINR, E1-99-1, Dubna. 1999.