



CBPF - CENTRO BRASILEIRO DE PESQUISAS FÍSICAS

Rio de Janeiro

CBPF - NF - 045 - 02

Notas de Física

CBPF-NF-045/02

December 2002

On the Octonionic Superconformal M-algebra

Francesco Toppan



MCT - Ministério da Ciência e Tecnologia

On the Octonionic Superconformal M-algebra*

Francesco Toppan

Centro Brasileiro de Pesquisas Físicas – CCP/CBPF/MCT
Rua Dr. Xavier Sigaud 150,
22290-180 Rio de Janeiro, RJ Brazil

Abstract

It is shown that besides the standard real algebraic framework for M -theory a consistent octonionic realization can be introduced. The octonionic M -superalgebra and superconformal M -algebra are derived. The first one involves 52 real bosonic generators and presents a novel and surprising feature, its octonionic $M5$ (super-5-brane) sector coincides with the $M1$ and $M2$ sectors. The octonionic superconformal M -algebra is given by $OSp(1, 8|\mathbf{O})$ and admits 239 bosonic and 64 fermionic generators.

Keywords Supersymmetry; M-theory; Division algebras.

*Proceedings of the Second Londrina Winter School “Mathematical Methods in Physics” (August 25 - August 30, 2002, Londrina-Paraná, Brazil)

1 Introduction

The generalized supersymmetries going beyond the standard HLS scheme[1] admit the presence of bosonic abelian tensorial central charges associated with the dynamics of extended objects (branes). It is widely known since the work of Ref. 2 that supersymmetries are related to division algebras. Indeed, even for generalized supersymmetries, classification schemes based on the associative division algebras (\mathbf{R} , \mathbf{C} , \mathbf{H}) are now available, see Ref. 3. For what concerns the remaining division algebra, the octonions, much less is known due to the complications arising from non-associativity. Octonionic structures were, nevertheless, investigated in Ref. 4,5, in application to the superstring theory.

It must be said that the octonions are at the very heart of many exceptional structures in mathematics. It is very well known, e.g., that they can be held responsible for the existence of the 5 exceptional Lie algebras. Indeed, G_2 is the automorphism group of the octonions, while F_4 is the automorphism group of the 3×3 octonionic-valued hermitian matrices realizing the exceptional $J_3(\mathbf{O})$ Jordan algebra. F_4 and the remaining exceptional Lie algebras (E_6 , E_7 , E_8) are recovered from the so-called “magic square Tits’ construction” which associates a Lie algebra to any given pair of division algebras, if one of these algebras coincides with the octonionic algebra.[6]

There is a line of thought[7] suggesting that Nature prefers exceptional structures. Following this line, in Ref. 8, the already recalled exceptional Jordan algebra $J_3(\mathbf{O})$ was e.g. used to define a unique Chern-Simons type of theory in the loop quantum gravity approach. It is quite evident from these and a whole series of other reasons, not last the fact that the octonions are the maximal division algebra, that they deserve a careful investigation. In this talk I will discuss the investigation in Ref. 9,10 concerning the possibility of realizing general supersymmetries in terms of the non-associative division algebra of the octonions. In particular in Ref. 9 it was shown that, besides the standard realization of the M -algebra (which supposedly underlines the M -theory) and involves real spinors, an alternative formulation, requiring the introduction of the octonionic structure, is viable. This is indeed possible due to the existence of an octonionic description for the Clifford algebra defining the 11-dimensional Minkowskian spacetime and its related spinors. The features of this octonionic M -superalgebra are puzzling. It is not at all surprising that it contains fewer bosonic generators, 52, w.r.t. the 528 of the standard M -algebra (this is expected, after all the imposition of an extra structure puts a constraint on a theory). What is really unexpected is the fact that new conditions, not present in the standard M -theory, are now found. These conditions imply that the different brane-sectors are no longer independent. The octonionic 5-brane contains the same degrees of freedom and is equivalent to the $M1$ and the $M2$ sectors. We can write this equivalence, symbolically, as $M5 \equiv M1 + M2$. This result is indeed very intriguing. It implies that quite non-trivial structures are found when investigating the octonionic construction of the M -theory. The next passage consists in defining the closed algebraic structure realizing the superconformal M -algebra. We will see that the $OSp(1, 32)$ superconformal algebra of the real M -theory is in this case replaced by the $OSp(1, 8|\mathbf{O})$ superalgebra of supermatrices with octonionic-valued entries.

2 Octonionic Clifford algebras and spinors.

In the $D = 11$ Minkowskian spacetime, where the M -theory should be found, the spinors are real and have 32 components. Since the most general symmetric 32×32 matrix admits 528 components, one can easily prove that the most general supersymmetry algebra in $D = 11$ can be presented as

$$\{Q_a, Q_b\} = (C\Gamma_\mu)_{ab}P^\mu + (C\Gamma_{[\mu\nu]})_{ab}Z^{[\mu\nu]} + (C\Gamma_{[\mu_1\dots\mu_5]})_{ab}Z^{[\mu_1\dots\mu_5]} \quad (1)$$

(where C is the charge conjugation matrix), while $Z^{[\mu\nu]}$ and $Z^{[\mu_1\dots\mu_5]}$ are totally anti-symmetric tensorial central charges, of rank 2 and 5 respectively, which correspond to extended objects[11, 12], the p -branes. Please notice that the total number of 528 is obtained in the r.h.s as the sum of the three distinct sectors, i.e.

$$528 = 11 + 66 + 462. \quad (2)$$

The algebra (1) is called the M -algebra. It provides the generalization of the ordinary supersymmetry algebra, recovered by setting $Z^{[\mu\nu]} \equiv Z^{[\mu_1\dots\mu_5]} \equiv 0$.

In the next section we will prove the existence of an octonionic version of Eq. (1). For this purpose we need at first to introduce the octonionic realizations of Clifford algebras and spinors. They exist only in a restricted class of spacetime signatures which includes the Minkowskian $(10, 1)$ spacetime.

The most convenient way to construct realizations of Clifford algebras is to iteratively derive them with the help of the following algorithm, allowing the recursive construction of $D+2$ spacetime dimensional Clifford algebras by assuming known a D dimensional representation. Indeed, it is a simple exercise to verify that if γ_i 's denotes the d -dimensional Gamma matrices of a $D = p + q$ spacetime with (p, q) signature (namely, providing a representation for the $C(p, q)$ Clifford algebra) then $2d$ -dimensional $D + 2$ Gamma matrices (denoted as Γ_j) of a $D + 2$ spacetime are produced according to either

$$\Gamma_j \equiv \begin{pmatrix} 0 & \gamma_i \\ \gamma_i & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & 1_d \\ -1_d & 0 \end{pmatrix}, \quad \begin{pmatrix} 1_d & 0 \\ 0 & -1_d \end{pmatrix} \\ (p, q) \mapsto (p + 1, q + 1). \quad (3)$$

or

$$\Gamma_j \equiv \begin{pmatrix} 0 & \gamma_i \\ -\gamma_i & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & 1_d \\ 1_d & 0 \end{pmatrix}, \quad \begin{pmatrix} 1_d & 0 \\ 0 & -1_d \end{pmatrix} \\ (p, q) \mapsto (q + 2, p). \quad (4)$$

Some remarks are in order. The two-dimensional real-valued Pauli matrices τ_A, τ_1, τ_2 which realize the Clifford algebra $C(2, 1)$ are obtained by applying either (3) or (4) to the number 1, i.e. the one-dimensional realization of $C(1, 0)$. We have indeed

$$\tau_A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \tau_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \tau_2 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (5)$$

The above algorithms can be applied to “lift” the Clifford algebra $C(0, 7)$, furnishing higher-dimensional Clifford algebras. $C(10, 1)$ is constructed by successively applying (no matter in which order) (3) and (4) to $C(0, 7)$. For what concerns $C(0, 7)$, it must be previously known. Two inequivalent realizations of $C(0, 7)$ can be constructed. The first one is associative and admits a matrix realization. Without loss of generality (the associative irreducible representation of $C(0, 7)$ is unique) we can choose expressing it through

$$C(0, 7) \equiv \begin{aligned} & \tau_A \otimes \tau_1 \otimes \mathbf{1}_2, \\ & \tau_A \otimes \tau_2 \otimes \mathbf{1}_2, \\ & \mathbf{1}_2 \otimes \tau_A \otimes \tau_1, \\ & \mathbf{1}_2 \otimes \tau_A \otimes \tau_2, \\ & \tau_1 \otimes \mathbf{1}_2 \otimes \tau_A, \\ & \tau_2 \otimes \mathbf{1}_2 \otimes \tau_A, \\ & \tau_A \otimes \tau_A \otimes \tau_A. \end{aligned} \quad (6)$$

On the other hand another, inequivalent, realization is at disposal. It is based on the identification of the $C(0, 7)$ Clifford algebra generators with the seven imaginary octonions τ_i satisfying the algebraic relation

$$\tau_i \cdot \tau_j = -\delta_{ij} + C_{ijk}\tau_k, \quad (7)$$

for $i, j, k = 1, \dots, 7$ and C_{ijk} the totally antisymmetric octonionic structure constants given by

$$C_{123} = C_{147} = C_{165} = C_{246} = C_{257} = C_{354} = C_{367} = 1 \quad (8)$$

and vanishing otherwise. This octonionic realization of the seven-dimensional Euclidean Clifford algebra will be denoted as $C_{\mathbf{O}}(0, 7)$. Similarly, the octonionic realization $C_{\mathbf{O}}(10, 1)$, obtained through the lifting procedure, is realized in terms of 4×4 matrices with octonionic entries.

One should be aware of the properties of the non-associative realizations of Clifford algebras. In the octonionic case the commutators $\Sigma_{\mu\nu} = [\Gamma_{\mu}, \Gamma_{\nu}]$ are no longer the generators of the Lorentz group. They correspond instead to the generators of the coset $SO(p, q)/G_2$, being G_2 the 14-dimensional exceptional Lie algebra of automorphisms of the octonions. As an example, in the Euclidean 7-dimensional case, these commutators give rise to $7 = 21 - 14$ generators, isomorphic to the imaginary octonions. Indeed

$$[\tau_i, \tau_j] = 2C_{ijk}\tau_k. \quad (9)$$

The algebra (9) is not a Lie algebra, but a Malcev algebra (due to the alternativity property satisfied by the octonions, a weaker condition w.r.t. associativity, see Ref. 13). It can be regarded[14, 15] as the “quasi” Lorentz algebra of homogeneous transformations acting on the seven sphere S^7 .

3 The octonionic M -superalgebra

The octonionic M -superalgebra is introduced by assuming an octonionic structure for the spinors which, in the $D = 11$ Minkowskian spacetime, are octonionic-valued 4-component

vectors. The algebra replacing (1) is given by

$$\{Q_a, Q_b\} = \{Q_a^*, Q_b^*\} = 0, \quad \{Q_a, Q_b^*\} = Z_{ab}, \quad (10)$$

where $*$ denotes the principal conjugation in the octonionic division algebra and, as a result, the bosonic abelian algebra on the r.h.s. is constrained to be hermitian

$$Z_{ab} = Z_{ba}^*, \quad (11)$$

leaving only 52 independent components.

The Z_{ab} matrix can be represented either as the 11 + 41 bosonic generators entering

$$Z_{ab} = P^\mu (C\Gamma_\mu)_{ab} + Z_{\mathcal{O}}^{\mu\nu} (C\Gamma_{\mu\nu})_{ab}, \quad (12)$$

or as the 52 bosonic generators entering

$$Z_{ab} = Z_{\mathcal{O}}^{[\mu_1 \dots \mu_5]} (C\Gamma_{\mu_1 \dots \mu_5})_{ab}. \quad (13)$$

Due to the non-associativity of the octonions, unlike the real case, the sectors individuated by (12) and (13) are not independent. Furthermore, as we have already seen for $k = 2$, in the antisymmetric products of k octonionic-valued matrices, a certain number of them are redundant (for $k = 2$, due to the G_2 automorphisms, 14 such products have to be erased). In the general case[16] a table can be produced expressing the number of independent components in D odd-dimensional spacetime octonionic realizations of Clifford algebras, by taking into account that out of the D Gamma matrices, 7 of them are octonionic-valued, while the remaining $D - 7$ are purely real. We get the following table, with the columns labeled by k , the number of antisymmetrized Gamma matrices and the rows by D (up to $D = 13$)

$D \setminus k$	0	1	2	3	4	5	6	7	8	9	10	11	12	13
7	1	7	7	1	1	7	7	1						
9	1	9	22	22	10	10	22	22	9	1				
11	1	11	41	75	76	52	52	76	75	41	11	1		
13	1	13	64	168	267	279	232	232	279	267	168	64	13	1

(14)

For what concerns the octonionic equivalence of the different sectors, it can be symbolically expressed, in different odd space-time dimensions, according to the table

$D = 7$	$M0 \equiv M3$
$D = 9$	$M1 + M2 \equiv M4$
$D = 11$	$M1 + M2 \equiv M5$
$D = 13$	$M2 + M3 \equiv M6$
$D = 15$	$M3 + M4 \equiv M0 + M7$

(15)

In $D = 11$ dimensions the relation between $M1 + M2$ and $M5$ can be made explicit as follows. The 11 vectorial indices μ are split into the 4 real indices, labeled by a, b, c, \dots

and the 7 octonionic indices labeled by i, j, k, \dots . The 52 independent components are recovered from $52 = 4 + 2 \times 7 + 6 + 28$, according to

4	$M1_a$	$M5_{[aijkl]} \equiv M5_a$	(16)
7	$M1_i, M2_{[ij]} \equiv M2_i$	$M5_{[abcdi]} \equiv M5_i, M5_{[ijklm]} \equiv M5_i$	
6	$M1_{[ab]}$	$M5_{[abijk]} \equiv M5_{[ab]}$	
$4 \times 7 = 28$	$M2_{[ai]}$	$M5_{[abcij]} \equiv M5_{[ai]}$	

4 The octonionic superconformal M -algebra

The conformal algebra of the octonionic M-theory can be introduced[10] adapting to the eleven dimensions the procedure discussed in Ref. 5 for the 10 dimensional case. It requires the identification of the conformal algebra of the octonionic $D = 11$ M -algebra with the generalized Lorentz algebra in the $(11, 2)$ -dimensional space-time. In such a space-time the octonionic Clifford's Gamma-matrices are 8-dimensional. The basis of the hermitian generators is given by the 64 antisymmetric two-tensors $C\Gamma_{[\mu_1\mu_2]}Z^{\mu_1\mu_2}$ and the 168 antisymmetric three tensors $C\Gamma_{[\mu_1\mu_2\mu_3]}Z^{\mu_1\mu_2\mu_3}$ (or, equivalently, by the 232 antisymmetric six-tensors $C\Gamma_{[\mu_1\dots\mu_6]}Z^{\mu_1\dots\mu_6}$). This is already an indication that the total number of generators in the conformal algebra is 232. We will show that this is the case.

According to Ref. 8 the conformal algebra can be introduced as the algebra of transformations leaving invariant the inner product of Dirac's spinors. In $(11, 2)$ this is given by $\psi^\dagger C \eta$, where the matrix C , the analogous of the Γ^0 , given by the product of the two space-like Clifford's Gamma matrices, is real-valued and totally antisymmetric. Therefore, the conformal transformations are realized by the octonionic-valued 8-dimensional matrices \mathcal{M} leaving C invariant, i.e. satisfying

$$\mathcal{M}^\dagger C + C \mathcal{M} = 0. \quad (17)$$

This allows identifying the (quasi)-group of conformal transformations with the (quasi)-group of symplectic transformations. Indeed, under a simple change of variables, C can be recast in the form

$$\Omega = \begin{pmatrix} 0 & \mathbf{1}_4 \\ -\mathbf{1}_4 & 0 \end{pmatrix}. \quad (18)$$

The most general octonionic-valued matrix leaving invariant Ω can be expressed through

$$\mathbf{M} = \begin{pmatrix} D & B \\ C & -D^\dagger \end{pmatrix}, \quad (19)$$

where the 4×4 octonionic matrices B, C are hermitian

$$B = B^\dagger, \quad C = C^\dagger. \quad (20)$$

It is easily seen that the total number of independent components in (19) is precisely 232, as we expected from the previous considerations.

It is worth noticing that the set of matrices \mathbf{M} of (19) type forms a closed algebraic structure under the usual matrix commutation. Indeed $[\mathbf{M}, \mathbf{M}] \subset \mathbf{M}$ endows the structure of $Sp(8|\mathbf{O})$ to \mathbf{M} . For what concerns the supersymmetric extension of the superconformal algebra, we have to accommodate the 64 real components (or 8 octonionic) spinors of $(11, 2)$ into a supermatrix enlarging $Sp(8|\mathbf{O})$. This can be achieved as follows. The two 4-column octonionic spinors α and β can be accommodated into a supermatrix of the form

$$\left(\begin{array}{c|cc} 0 & -\beta^\dagger & \alpha^\dagger \\ \alpha & 0 & 0 \\ \beta & 0 & 0 \end{array} \right). \quad (21)$$

Under anticommutation, the lower bosonic diagonal block reduces to $Sp(8|\mathbf{O})$. On the other hand, extra seven generators, associated to the 1-dimensional antihermitian matrix A

$$A^\dagger = -A, \quad (22)$$

i.e. representing the seven imaginary octonions, are obtained in the upper bosonic diagonal block. Therefore, the generic bosonic element is of the form

$$\left(\begin{array}{c|cc} A & 0 & 0 \\ 0 & D & B \\ 0 & C & -D^\dagger \end{array} \right), \quad (23)$$

with A , B and C satisfying (22) and (20).

The closed superalgebraic structure, with (21) as generic fermionic element and (23) as generic bosonic element, will be denoted as $OSp(1, 8|\mathbf{O})$. It is the superconformal algebra of the M -theory and admits a total number of 239 bosonic generators.

References

- [1] R. Haag, J. Lopuszański and M. Sohnius, *Nucl. Phys.* **B 88**, (1975), 257.
- [2] T. Kugo and P. Townsend, *Nucl. Phys.* **B 221** (1983), 357.
- [3] R. D'Auria, S. Ferrara and S. Vaula, *Class. Quant. Grav.* **18** (2001), 3181; JHEP 0010 (2000), 013; S. Ferrara, M.A. Lledo, hep-th/0112177.
- [4] D.B. Fairlie and A.C. Manogue, *Phys. Rev.* **D 34**, 1(1986), 1832.
- [5] K.W. Chung and A. Sudbery, *Phys. Lett.* **B 198**, (1987), 161.
- [6] C.A. Barton and A. Sudbery, math.RA/0203010.
- [7] P. Ramond, "Algebraic Dreams", hep-th/0112261.
- [8] L. Smolin, hep-th/0104050.

- [9] J. Lukierski and F. Toppan, *Phys. Lett.* **B 539**, (2002), 266.
 - [10] J. Lukierski and F. Toppan, "Octonionic M -theory and $D = 11$ Generalized Conformal and Superconformal Algebras". Work in preparation.
 - [11] E. Bergshoeff, E. Sezgin, P. Townsend, *Ann. Phys.* **18**, 330 (1987).
 - [12] M.J. Duff, R.R. Khuri, J.X. Lu, *Phys. Rep.* **259**, 213 (1995).
 - [13] M. Günaydin and S.V. Ketov, *Nucl. Phys.* **B 467** (1996), 215.
 - [14] Z. Hasiewicz and J. Lukierski, *Phys. Rev.* **145 B** (1984), 65.
 - [15] M. Cederwall and C. Preitschopf, *Comm. Math. Phys.* **167** (1995), 373.
 - [16] H.L. Carrion, M. Rojas and F. Toppan, "Quaternionic and Octonionic Spinors. A classification". Work in preparation.
-

NOTAS DE FÍSICA é uma pré-publicação de trabalho original em Física.
Pedidos de cópias desta publicação devem ser enviados aos autores ou ao:

Centro Brasileiro de Pesquisas Físicas
Área de Publicações
Rua Dr. Xavier Sigaud, 150 – 4^o andar
22290-180 – Rio de Janeiro, RJ
Brasil
E-mail: socorro@cbpf.br/valeria@cbpf.br

NOTAS DE FÍSICA is a preprint of original unpublished works in Physics.
Requests for copies of these reports should be addressed to:

Centro Brasileiro de Pesquisas Físicas
Área de Publicações
Rua Dr. Xavier Sigaud, 150 – 4^o andar
22290-180 – Rio de Janeiro, RJ
Brazil
E-mail: socorro@cbpf.br/valeria@cbpf.br