Fermi National Accelerator Laboratory

FN-336
0402.000

On an Invariant of a System affected by Intra-Beam Scattering

Ken TAKAYAMA

Fermilab, March , 81

1. Introduction

A. Piwinski has shown[1] that there is an invariant that is useful in understanding the effects of intra-beam scattering on betatron oscillations and the energy spread. After some mathematical manipulations, the invariant can be written as

$$\frac{1}{n}\left(\frac{1}{\gamma^2}-\alpha\right)\left\langle\left(\frac{\Delta p}{p}\right)^2\right\rangle + \frac{1}{\beta_x}\langle\epsilon_x\rangle + \frac{1}{\beta_y}\langle\epsilon_y\rangle = \text{const} \qquad (1-1)$$

where $p, \Delta p$ are the momentum and momentum deviation, $\pi\epsilon_x$ and $\pi\epsilon_y$ are the betatron emittances for horizontal and vertical directions, respectively, β_x and β_y are the horizontal and vertical betatron amplitude functions, γ is the particle energy in units of its rest energy, α is the momentum compaction factor ($=\overline{\eta^2/\beta_x^2}$ which is considered by Piwinski as a good approximation), η is the momentum dispersion function, $\langle A \rangle$ is the mean value of a quantity A over all particles, $\bar{A}$ is the mean value of a quantity A in the orbit, and the integer n is 2 for an unbunched beam.

Operated by Universities Research Association Inc. under contract with the United States Department of Energy

If we asume that betatron oscillations and energy oscillations are harmonic oscillators and that intra-beam scattering is a local elastic collision, we can derive the expression for this invariant more easily with the aid of the energy conservation law.

2. System of N three-dimensional harmonic oscillators

According to the above assumption, we can regard a beam as an isolated system of N three-dimensional harmonic oscillators which experience many random collisions among themselves. Here the longitudinal oscillation is taken as a free motion because we consider the case of an unbunched beam.

Consider the behavior of a single particle before and after a collision. Its behavior can be described by the Hamiltonian

$$H(x, P_x; Y, P_Y; \psi, \delta; \theta) = \frac{\nu_x}{2}(P_x^2 + x^2) + \frac{\nu_Y}{2}(P_Y^2 + Y^2) - \lambda x \delta + \frac{\nu_s}{2}\delta^2 \qquad (2-1)$$

where ν_x, ν_Y, ν_s are the horizontal and vertical betatron tunes ,and longitudinal frequency, $x, P_x; Y, P_Y; \psi, \delta$ are the canonical variables for horizontal, vertical ,and longitudinal motions ($\delta \equiv \Delta p/p$), λ is a coupling coefficient, and θ = s/R where R is the mean machine radius and s is the distance along the orbit.

We can separate the horizontal excursion of Eq. (2-1) into two parts, the equilibrium orbit and the homogeneous harmonic oscillation around the equilibrium orbit. This homogeneous harmonic oscillation corresponds to a pure betatron oscillation and the equilibrium orbit corresponds to the well-known closed

orbit which varies linearly with δ , the longitudinal momentum deviation. The equation of motion obtained from the Hamiltonian (2-1) is then

$$\ddot{x} = -\nu_x^2 x + \lambda \nu_x \delta \qquad \left(\dot{} \equiv \frac{d}{d\theta} \right) \tag{2 - 2}$$

Therefore, the equilibrium orbit (x_{eq} ,P_{eq}) is written as

$$\begin{aligned} x_{eq} &= D\delta \qquad\qquad (D \equiv \lambda/\nu_x) \\ P_{eq} &= 0 \end{aligned} \tag{2 - 3}$$

Transforming into a new canonical variable (χ , P_χ),

$$\begin{aligned} \chi &= x - x_{eq} \\ P_\chi &= P_x - P_{eq} \end{aligned}$$

we find, from the generating function,

$$g(\chi, P_x; \theta) = -(x_{eq} + \chi) P_x + P_{eq} \cdot \chi \tag{2 - 4}$$

the new Hamiltonian

$$\begin{aligned} K &= H + \frac{\partial g}{\partial \theta} \\ &= \frac{\nu_x}{2}[P_x^2 + (\chi + D\delta)^2] - \lambda(\chi + D\delta)\delta + \frac{\nu_s}{2}\delta^2 + \frac{\nu_Y}{2}[P_Y^2 + Y^2] \\ &= \frac{\nu_x}{2}[P_\chi^2 + \chi^2] + \frac{\nu_Y}{2}[P_Y^2 + Y^2] + \frac{1}{2}(\nu_s - \nu_x D^2)\delta^2 \end{aligned} \tag{2 - 5}$$

In terms of action-angle variables, we can write the Hamiltonian in the form

$$G = \nu_x J_\chi + \nu_Y J_Y + \frac{1}{2}(\nu_s - \nu_x D^2) J_s \tag{2 - 6}$$

Next, consider the behavior of <u>the system</u> before and after a collision. This behavior is described by the Hamiltonian H_{total}

$$H_{total} = \sum_{i=1}^{N} [\nu_x J_\chi + \nu_Y J_Y + \frac{1}{2}(\nu_s - \nu_x D^2) J_s]_i \tag{2 - 7}$$

Since we have assumed that the Coulomb interaction between paticles is a local elastic collision, the Hamiltonian H_{total} must be an invariant of the motion. We therefore obtain easily the invariant expression

$$\nu_x \langle J_x \rangle + \nu_Y \langle J_Y \rangle + \frac{1}{2}(\nu_s - \nu_x D^2)\langle J_s \rangle = const \quad (2-8)$$

where $\langle A \rangle$ is the mean value of A over all particles.

Finally, we can rewrite this expression in terms of betatron parameters. Relations between the parameters used above and the orbit parameters of a real machine are

$D = \eta(s)/\sqrt{\beta(s)}$ because $x(s) = \sqrt{\beta_x(s)}\,X \longrightarrow x_{eq}(s) = \sqrt{\beta_x(s)}\,X_{eq} \longrightarrow \eta(s) = \sqrt{\beta_x(s)}\,D$ [2]

where $x(s), x_{eq}(s)$ are the horizontal excursion and the equilibrium orbit in a real ring,

$\eta(s)$ is the momentum dispersion function.

$\nu_{x,Y} = R/\bar{\beta}_{x,y}$ (smooth approximation)

$\nu_s = R/\gamma^2$ because in the rest frame (independent variable: s) the momentum deviation takes the term $\Delta p/\gamma$ (See Appendix) and in the θ frame (independent variable: θ) the longitudinal energy must be multiplied by R.

Thus Eq. (2-8) becomes

$$R\frac{\langle J_x \rangle}{\bar{\beta}_x} + R\frac{\langle J_y \rangle}{\bar{\beta}_y} + \frac{R}{2}\left(\frac{1}{\gamma^2} - \frac{\eta^2(s)}{\bar{\beta}_x\,\beta_x(s)}\right)\langle J_s \rangle = const \quad (2-9)$$

Using the approximate expression for the momentum compaction factor[2]

$$\alpha = \frac{\eta^2(s)}{\bar{\beta}_x\,\beta_x(s)}$$

we obtain, from Eq.(2-9),

$$\frac{\langle J_x \rangle}{\bar{\beta}_x} + \frac{\langle J_y \rangle}{\bar{\beta}_y} + \frac{1}{2}\left(\frac{1}{\gamma^2} - \alpha\right)\langle J_s \rangle = \text{const} \qquad (2-10)$$

which is identical with the original form, Eq.(1-1), derived by Piwinski.

3. Conclusion

Although particles in the beam undergo many collisions, the invariant Eq.(2-10) is valid if the interaction is elastic and this is simply a consequence of the conservation of energy.

Aknowledgements

The auther is grateful to S. Ohnuma, F. Cole and A. Ruggiero for valuable suggestions and comments.

References:

(1) A. Piwinski, " INTRA-BEAM SCATTERING ", Proc. 9th Int. Conf. on High Energy Accelerators, p. 405 (1974)

(2) M. Sands, " THE PHYSICS OF ELECTRON STORAGE RINGS-AN INTRODUCTION", SLAC-121, UC-283(ACC) , p.77 (1970)

Appendix



K is the laboratory system and K′ is the system at rest with the synchronous particle which has the velocity $v_0 (=c\beta_0)$ in K. The space-time coordinates of a test particle are (x,y,z,ict) in K and (x′,y′,z′,ict′) in K′. The four momenta $(p_x, p_y, p_z, iE/c)$ in K are

$$P_x = \frac{m_0 v_x}{\sqrt{1-\beta^2}}, \quad P_y = \frac{m_0 v_y}{\sqrt{1-\beta^2}}, \quad P_z = \frac{m_0 v_z}{\sqrt{1-\beta^2}}, \quad E = \frac{m_0 c^2}{\sqrt{1-\beta^2}} \qquad \text{(A - 1)}$$

with $\beta = v/c$ (v is the velocity in K) and $v_z \simeq v$. Similar relations hold for the four-momenta $(p_x', p_y', p_z', iE'/c)$ measured in K′.

In K the longitudinal momentum deviation from the design value is

$$(\Delta P_z) \equiv P_z - P_{z0}$$

$$= m_0 c \left[\frac{\beta}{\sqrt{1-\beta^2}} - \frac{\beta_0}{\sqrt{1-\beta_0^2}} \right]. \qquad \text{(A - 2)}$$

Transformed into K′, the momentum p_z' is written in the form

$$P_z' = \frac{P_z + i\beta_0 \cdot iE/c}{\sqrt{1-\beta_0^2}}$$

$$= \frac{m_0 c}{\sqrt{1-\beta_0^2}} \cdot \frac{\beta - \beta_0}{\sqrt{1-\beta^2}}.$$

Therefore the longitudinal momentum deviation in K′ is written as

$$(\Delta P_z)' = \frac{m_0 c}{\sqrt{1-\beta_0^2}} \cdot \frac{\beta - \beta_0}{\sqrt{1-\beta^2}} \qquad \text{(A - 3)}$$

We define a small parameter ϵ,

$$\epsilon \equiv \beta - \beta_0 \qquad \text{(A - 4)}$$

We expand the right-hand sides of Eq.(A-2) and (A-3) in ϵ and retain only the lowest order terms. We then get the expressions for the longitudinal momentum deviation

$$(\Delta P_z) = \frac{m_0 c \epsilon}{(1-\beta_0^2)^{3/2}} , \qquad \text{(A - 5)}$$

$$(\Delta P_z)' = \frac{m_0 c \epsilon}{(1-\beta_0^2)} , \qquad \text{(A - 6)}$$

so that

$$(\Delta P_z)' = \frac{(\Delta P_z)}{\gamma} . \qquad \text{(A - 7)}$$