TM-708
0420.042

Fermilab

MULTIPOLE FIELDS IN CIRCULAR CONCAVE POLES

S.C. Snowdon

January 4, 1977

Introduction

A paper by G. Bosi[1] provides an exact calculation of a quadrupole field in a circular concave pole geometry. His method is readily generalized to the case of 2N-poles.



Fig. 1 z-Plane

The 2N-pole case is shown in Fig. 1 for the case of N = 4 (octupole). The complex z-plane is measured in units of the radius of the poles. Thus $|z| = 1$ is the circle on which the concave poles are drawn. The problem of finding the complex potential $W = U + iV$ where V is the potential function and U the stream function will be solved by complex variable transformations of the z-plane to the t-plane

where the excitation is relatively clear.

Transformations

Transform the z-plane to the ω-plane using

$$\omega = \ln z \ . \tag{1}$$

One sector of Fig. 1 will appear as



Fig. 2 Map of One Sector of Fig. 1 in ω-plane.

The ω-plane can be generated by a Schwarz-Christoffel transformation from the t-plane according to

$$\omega = C_1 \int t^{-1}(1-t)^{\frac{1}{2}} \left(1-\frac{t}{a^2}\right)^{-\frac{1}{2}} dt + C_2 \tag{2}$$



Fig. 3 Map of Fig. 2 Boundary to Real Axis of t-plane

Integration of Eq. (2) gives

$$\omega = C_1 \left\{ a \ \ell n\left(\frac{1+\xi}{1-\xi}\right) - \ell n\left(\frac{1+a\xi}{1-a\xi}\right) \right\} + C_2, \tag{3}$$

where

$$\xi = \frac{1}{a}(1-t)^{\frac{1}{2}}\left(1-\frac{t}{a^2}\right)^{-\frac{1}{2}} . \tag{4}$$

Putting $t = 1$ for which $\omega = i\gamma$ gives $C_2 = i\gamma$. Also putting t equal to a negative real number gives a real positive ξ such that $1 < \xi < \frac{1}{a}$. Equation (3) assuming that C_1 is real gives ω = Real Number + $ia\pi C_1 + i\gamma$. Hence from Fig. (1) $\frac{\pi}{2N} = a\pi C_1 + \gamma$. For $|t|$ near zero $\omega \to C_1 \ \ell n\ t$ which gives $C_1 = \frac{1}{2N}$ since the imaginary part of ω for small negative t is $\pi/2N$. Thus

$$C_1 = \frac{1}{2N}, \qquad C_2 = i\gamma, \qquad a = 1 - \frac{2N}{\pi}\gamma . \tag{5}$$

Potential

Smythe[2] gives the solution of the potentials of a charged strip. For the origin shown in Fig. 3 and choosing the imaginary part of the complex potential W as the potential function one has from Sm 4.22 (3)

$$t - \frac{a^2}{2} = -\frac{a^2}{2} \sin\left[i \frac{\pi}{V_o}\left(W - i\frac{V_o}{2}\right)\right] \tag{6}$$

or

$$W = -\frac{iV_o}{\pi} \sin^{-1}\left(1-\frac{2t}{a^2}\right) + i\frac{V_o}{2} = i\frac{V_o}{\pi} \cos^{-1}\left(1-\frac{2t}{a^2}\right). \tag{7}$$

Using Pierce[3] (644-645) this may be written as

$$W = i\frac{2V_o}{\pi} \tan^{-1}\sqrt{\frac{\frac{t}{a^2}}{1-\frac{t}{a^2}}} = i\frac{2V_o}{\pi} \sin^{-1}\left(\frac{\sqrt{t}}{a}\right) . \tag{8}$$

Multipole Expansion

One desires to obtain a series of expansions of Eq. (8) in powers of z. First obtain an expansion of Eq. (8) in powers of t. From Gradsteyn and Ryzhik[4] (1.641) and (8.335) one has

$$W = i\frac{2V_o}{\pi} \cdot \frac{1}{\sqrt{\pi}} \sum_{n=o}^{\infty} \frac{a^{-(2n+1)}}{2n+1} \cdot \frac{\Gamma(n+\frac{1}{2})}{n!} t^{n+\frac{1}{2}} . \tag{9}$$

Next one needs t expanded in a power series in ω. First obtain ω as a power series in t. To this end expand $d\omega/dt$ from Eq. (2). Abramowitz and Stegun[5] (3.6.8) give

$$(1-t)^{\frac{1}{2}} = \sum_{n=\infty}^{\infty} (-1)^n \binom{\frac{1}{2}}{n} t^n \tag{10}$$

and

$$(1-\frac{t}{a^2})^{-\frac{1}{2}} = \sum_{m=o}^{\infty} (-1)^m a^{-2m} \binom{-\frac{1}{2}}{m} t^m . \tag{11}$$

Hence from Eq. (2) and Eq. (5) after juggling indices

$$\frac{d\omega}{dt} = \frac{1}{2N} \sum_{n=o}^{\infty} \sum_{m=o}^{n} (-1)^n a^{-2m} \binom{\frac{1}{2}}{n-m} \binom{-\frac{1}{2}}{m} t^{n-1} . \tag{12}$$

Abramowitz and Stegun[5] (6.1.21) and (6.1.17) also give

$$\binom{\frac{1}{2}}{n-m} = \frac{\Gamma(\frac{3}{2})}{\Gamma(n-m+1)\Gamma(\frac{3}{2}-n+m)} = \frac{-\frac{1}{2}\Gamma(\frac{1}{2})\Gamma(n-m-\frac{1}{2})(-1)^{n-m}}{(n-m)!\pi} \tag{13}$$

and

$$\binom{-\frac{1}{2}}{m} = \frac{\Gamma(\frac{1}{2})}{\Gamma(m+1)\Gamma(\frac{1}{2}-m)} = \frac{\Gamma(\frac{1}{2})\Gamma(m+\frac{1}{2})(-1)^m}{m!\pi} \tag{14}$$

or

$$\frac{d\omega}{dt} = -\frac{1}{4\pi N} \sum_{n=o}^{\infty} \sum_{m=o}^{n} a^{-2m} \frac{\Gamma(m+\frac{1}{2})\Gamma(n-m-\frac{1}{2})}{m!(n-m)!} t^{n-1} . \tag{15}$$

Integration yields

$$\omega = \omega_1 + \frac{1}{2N} \ell nt - \frac{1}{4\pi N} \sum_{n=1}^{\infty} \sum_{m=o}^{n} a^{-2m} \frac{\Gamma(m+\frac{1}{2})\Gamma(n-m-\frac{1}{2})}{m!(n-m)!} \cdot \frac{t^n}{n} , \tag{16}$$

where the constant of integration is found by comparing Eq. (16) with Eq. (3) for small $|t|$. Thus

$$\omega_1 = \frac{1}{2N} \left\{ a \ \ell n\left(\frac{1+a}{1-a}\right) - \ell n\left(\frac{4a^2}{1-a^2}\right) \right\}. \tag{17}$$

Since one expects z^N to be the first term in the multipole expansion of the potential W, first expand z^N in a power series in t. For $|t|$ small from Eq. (1) and (17) one has

$$z^N = e^{N\omega_1} \sqrt{t} . \tag{18}$$

Therefore one expects

$$e^{N(\omega-\omega_1)} = \sum_{n=o}^{\infty} \alpha_n \ t^{n+\frac{1}{2}} . \tag{19}$$

To find α_n first define for convenience

$$Q_n = -\frac{1}{4\pi} \cdot \frac{1}{n} \sum_{m=o}^{n} a^{-2m} \frac{\Gamma(m+\frac{1}{2})\Gamma(n-m-\frac{1}{2})}{m!(n-m)!} . \tag{20}$$

From Eqs. (16) and (20)

$$e^{N(\omega-\omega_1)} = t^{\frac{1}{2}} e^{\sum_{n=1}^{\infty} Q_n \ t^n} . \tag{21}$$

Comparing this with Eq. (19) one has

$$\sum_{n=o}^{\infty} \alpha_n t^n = e^{\sum_{k=1}^{\infty} Q_k t^k} . \tag{22}$$

But the coefficients α_n are given by Cauchy's integral formula as

$$\alpha_n = \frac{1}{2\pi i} \oint_c \frac{e^{\sum_{k=1}^{\infty} Q_k t^k}}{t^{n+1}} dt, \tag{23}$$

where C is any simple contour surrounding the origin. For t = o one sees from Eq. (22) that

$$\alpha_o = 1. \tag{24}$$

To find the other α_n consider $\sum_{j=1}^{n} j\alpha_{n-j} Q_j$. From Eq. (23) one has

$$\frac{1}{2\pi i} \oint_c \frac{1}{t_n} \sum_{j=1}^{n} jQ_j t^{j-1} e^{\sum_{k=1}^{\infty} Q_k t^k} dt = \sum_{j=1}^{n} j\alpha_{n-j} Q_j . \tag{25}$$

Again, for convenience let

$$P(t) = \sum_{j=1}^{\infty} Q_j t^j . \tag{26}$$

Then

$$P'(t) = \sum_{j=1}^{\infty} jQ_j t^{j-1} = \sum_{j=1}^{n} jQ_j t^{j-1} + R(t), \tag{27}$$

where the expansion of R(t) in powers of t to t^n as the lowest

order terms. Hence Eq. (25) becomes

$$\sum_{j=1}^{n} j\alpha_{n-j}\, Q_j = \frac{1}{2\pi i} \oint_c \frac{1}{t^n}[P'(t)-R(t)]e^{P(t)}\, dt. \tag{28}$$

The term in R(t) does not yield a simple pole and the term in P'(t) yields after integration by parts.

$$\sum_{j=1}^{n} j\alpha_{n-j}\, Q_j = \frac{1}{2\pi i} \left\{ \frac{e^{P(t)}}{t^n} \Big| + n \oint_c \frac{e^P}{t^{n+1}}\, dt \right\}. \tag{29}$$

The first term on the RHS is zero since the contour is closed. After using Eq. (23), one finally has

$$\alpha_n = \frac{1}{n} \sum_{j=1}^{n} j\, \alpha_{n-j}\, Q_j\ , \tag{30}$$

which is the recursion relation from which all the α_n may be found.

The next step requires the expansion of t in powers of z. From Eq. (18) one sees that the first term is

$$t^{\frac{1}{2}} = (ze^{-\omega_1})^N = e^{N(\omega-\omega_1)}. \tag{31}$$

However a 2N-pole expansion has terms in the sequence (Z^N, Z^{3N}, Z^{5N}, etc.). Hence, one expects

$$t^{\frac{1}{2}} = e^{N(\omega-\omega_1)} \sum_{k=o}^{\infty} A_k\, e^{2kN(\omega-\omega_1)}\ . \tag{32}$$

Since $t^{n+\frac{1}{2}}$ is desired raise Eq. (32) to the $n^{\underline{th}}$ power. Thus

$$t^{\frac{n}{2}} = e^{nN(\omega-\omega_1)} \sum_{k=o}^{\infty} A_k\, e^{2kN(\omega-\omega_1)^n}$$

$$= e^{nN(\omega-\omega_1)} \sum_{k_1=o}^{\infty} \sum_{k_2=o}^{\infty} \cdots \sum_{k_n=o}^{\infty} A_{k_1} A_{k_2} \cdots A_{k_n} e^{2N(\omega-\omega_1)(k_1+k_2+\cdots k_n)} . \tag{33}$$

Juggling the indices such that $k_1 + k_2 + \cdots k_n = m$, one may write Eq. (33) as

$$t^{\frac{n}{2}} = e^{nN(\omega-\omega_1)} \sum_{m=1}^{\infty} \beta_{m,n} e^{2mN(\omega-\omega_1)} . \tag{34}$$

Since Eq. (31) obtains for small $|t|$ it follows that

$$\beta_{o,1} = 1 . \tag{35}$$

Next, notice that the use of Eq. (34) in the identity, $t^{\frac{1}{2}(n+1)} = t^{\frac{n}{2}} t^{\frac{1}{2}}$, gives

$$\sum_{m=o}^{\infty} \beta_{m,n+1} e^{2mN(\omega-\omega_1)} = \sum_{k=o}^{\infty} \sum_{\ell=o}^{\infty} \beta_{k,n} \beta_{\ell,1} e^{2(k+\ell)N(\omega-\omega_1)} . \tag{36}$$

Let $k+\ell = m$ and equate coefficients of like powers

$$\beta_{m,n+1} = \sum_{k=o}^{\infty} \beta_{k,n} \beta_{m-k,1} . \tag{37}$$

Finally Eq. (34) gives for $t^{r+\frac{1}{2}}$

$$t^{r+\frac{1}{2}} = e^{(2r+1)N(\omega-\omega_1)} \sum_{m=o}^{\infty} \beta_{m,2r+1} e^{2mN(\omega-\omega_1)} . \tag{38}$$

Substituting this into Eq. (19) yields

$$e^{N(\omega-\omega_1)} = \sum_{r=o}^{\infty} \alpha_r \sum_{m=o}^{\infty} \beta_{m,2r+1} e^{[2(m+r)+1]N(\omega-\omega_1)} . \tag{39}$$

Rewriting the sum using m + r = n gives

$$e^{N(\omega-\omega_1)} = \sum_{n=o}^{\infty} \sum_{r=o}^{n} \alpha_r \, \beta_{n-r,\ 2r+1} \, e^{(2n+1)N(\omega-\omega_1)} . \tag{40}$$

Equating coefficients of like powers on each side gives

$$\alpha_o \, \beta_{o,1} = 1 \tag{41}$$

$$\sum_{r=o}^{n} \alpha_r \, \beta_{n-r,\ 2r+1} = 0 \quad (n = 1,2,3, \text{ etc.}). \tag{42}$$

Since $\alpha_o = 1$, Eq. (41) simply confirms Eq. (35). Equation (42) may be written as

$$\alpha_o \, \beta_{n,1} + \sum_{r=1}^{n} \alpha_r \, \beta_{n-r,\ 2r+1} = 0 \tag{43}$$

or

$$\beta_{n,1} = - \sum_{r=1}^{n} \alpha_r \, \beta_{n-r,\ 2r+1} . \tag{44}$$

Using Eq. (38) one may substitute into Eq. (9) to give

$$W = i\frac{2V_o}{\pi} \frac{1}{\sqrt{\pi}} \sum_{n=o}^{\infty} \frac{a^{-(2n+1)}}{2n+1} \cdot \frac{\Gamma(n+\frac{1}{2})}{n!} \cdot$$

$$\sum_{m=o}^{\infty} \beta_{m,2n+1} \, e^{[2(m+n)+1]N(\omega-\omega_1)} \tag{45}$$

Then, setting m + n = p and juggling the indices one has

$$W = i\frac{2V_o}{\pi} \frac{1}{\sqrt{\pi}} \sum_{p=o}^{\infty} \sum_{n=o}^{p} \frac{a^{-(2n+1)}}{2n+1} \cdot$$

$$\frac{\Gamma(n+\frac{1}{2})}{n!} \beta_{p-n,2n+1} \, e^{(2p+1)N(\omega-\omega_1)} \tag{46}$$

or

$$W = i\frac{2V_o}{\pi}\frac{1}{\sqrt{\pi}}\left\{a^{-1}\,\Gamma(\tfrac{1}{2})\,\beta_{o,1}\,e^{N(\omega-\omega_1)}\right.$$

$$+\left[a^{-1}\,\Gamma(\tfrac{1}{2})\,\beta_{1,1}+\frac{a^{-3}}{3}\,\Gamma(\tfrac{3}{2})\,\beta_{o,3}\right]e^{3N(\omega-\omega_1)}$$

$$+\left[a^{-1}\,\Gamma(\tfrac{1}{2})\,\beta_{2,1}+\frac{a^{-3}}{3}\,\Gamma(\tfrac{3}{2})\,\beta_{1,3}+\frac{a^{-5}}{5}\,\frac{\Gamma(\frac{5}{2})}{2!}\,\beta_{0,5}\right]e^{5N(\omega-\omega_1)}$$

$$\left.+\cdots\cdots\cdots\right\}. \tag{47}$$

From Eq. (37)

$$\beta_{o2} = \beta_{o1}\beta_{o1} \qquad \beta_{o3} = \beta_{o2}\beta_{o1}$$

$$\beta_{o4} = \beta_{o3}\beta_{o1} \qquad \beta_{o5} = \beta_{o4}\beta_{o1}\ . \tag{48}$$

Hence, since from Eq. (35) $\beta_{o1} = 1$, one has

$$\beta_{o1} = \beta_{o2} = \beta_{o3} = \beta_{o4} = \beta_{o5} = 1\ . \tag{49}$$

From Eq. (44)

$$\beta_{1,1} = -\,\alpha_1\beta_{o,3} = -\,\alpha_1 \tag{50}$$

and

$$\beta_{2,1} = -(\alpha_1\beta_{1,3} + \alpha_2\beta_{o,5})\ . \tag{51}$$

But, from Eq. (37)

$$\beta_{12} = \beta_{o1}\beta_{11} + \beta_{11}\beta_{o1}$$

$$\beta_{1,3} = \beta_{o,2}\beta_{1,1} + \beta_{1,2}\beta_{o1}\ . \tag{52}$$

Thus

$$\beta_{1,3} = -3\alpha_1 \tag{53}$$

and

$$\beta_{2,1} = 3\alpha_1^{\,2} - \alpha_2 \ . \tag{54}$$

From Eqs. (30) and (20)

$$\alpha_1 = Q_1 = \frac{1}{4} \cdot \frac{1-a^2}{a^2} \tag{55}$$

$$\alpha_2 = Q_2 + \frac{1}{2} Q_1^{\,2} = \frac{1}{32a^4} (4-4a^2) \quad . \tag{56}$$

Inserting the various β-coefficients into Eq. (47) gives for the multipole expansion

$$W = i\frac{2V_o}{\pi} \frac{1}{\sqrt{\pi}} \left\{ \frac{1}{a}\left(ze^{-\omega_1}\right)^N + \frac{1}{12a^3} (3a^2-1)\left(ze^{-\omega_1}\right)^{3N} + \frac{1}{80a^5} (1-10a^2 + 15a^4)\left(ze^{-\omega_1}\right)^{5N} + \cdots \right\}. \tag{57}$$

Note that for all multipoles it is possible to eliminate the next higher term in the series by choosing $a = 1/\sqrt{3}$ or from Eq. (5)

$$N\gamma = 38.04^{\circ} \ . \tag{58}$$

Acknowledgement

The paper by G. Bosi[1] was brought to my attention during discussions with Prof. M. Szilagyi while he was visiting at the Fermi Institute.

References

1. G. Bosi, Rev. Sci. Instrum. 45, 1260 (1974)

2. W.R. Smythe, "Static and Dynamic Electricity", Third Edition, McGraw-Hill Book Company, New York, 1968

3. B.O. Pierce, "A Short Table of Integrals", Ginn and Company, Boston, 1929

4. I.S. Gradshteyn and I.W. Ryzhik, "Table of Integrals Series and Products", Fourth Edition, Academic Press, New York, 1965

5. M. Abramowitz and I.A. Stegun, Editors, "Handbook of Mathematical Functions", U.S. Government Printing Office Washington, D.C., 1964