

# RECENT RESULTS FROM MINIBOONE

---

*E. D. Zimmerman*  
*University of Colorado*

RENCONTRES DE BLOIS  
CHÂTEAU DE BLOIS  
BLOIS, FRANCE  
31 MAY 2011

# Recent Results from MiniBooNE

---


- MiniBooNE
- Neutrino cross-sections
  - Quasielastic and elastic scattering
  - Hadron production channels
- Neutrino Oscillations
- Antineutrino Oscillations

# Motivating MiniBooNE: LSND

## Liquid Scintillator Neutrino Detector


- Stopped  $\pi^+$  beam at Los Alamos LAMPF produces  $\nu_e, \nu_\mu, \bar{\nu}_\mu$  but no  $\bar{\nu}_e$  (due to  $\pi^-$  capture).

Search for  $\bar{\nu}_e$  appearance via reaction:




- Look for delayed coincidence of positron and neutron capture.
- Major background non-beam (measured, subtracted)
- 3.8 standard dev. excess above background.
- Oscillation probability:

$$P(\bar{\nu}_\mu \rightarrow \bar{\nu}_e) = (2.5 \pm 0.6_{\text{stat}} \pm 0.4_{\text{syst}}) \times 10^{-3}$$



# LSND oscillation signal

- LSND “allowed region” shown as band
- KARMEN2 is a similar experiment with a slightly smaller L/E; they see no evidence for oscillations. Excluded region is to right of curve.



# The Overall Picture

LSND	$\Delta m^2 > 0.1 \text{eV}^2$	$\bar{\nu}_\mu \leftrightarrow \bar{\nu}_e$
Atmos.	$\Delta m^2 \approx 2 \times 10^{-3} \text{eV}^2$	$\nu_\mu \leftrightarrow \nu_\tau$
Solar	$\Delta m^2 \approx 10^{-4} \text{eV}^2$	$\nu_e \leftrightarrow \nu_\tau$

- With only 3 masses, can't construct 3  $\Delta m^2$  values of different orders of magnitude!
- Current ideas out there:
  - An experiment or two is wrong
  - Sterile neutrino sector: extra masses and mixing angles


# MiniBooNE: E898 at Fermilab

---

- Purpose is to test LSND with:
  - Higher energy
  - Different beam
  - Different oscillation signature
  - Different systematics
- $L=500$  meters,  $E=0.5-1$  GeV: same  $L/E$  as LSND.


# Oscillation Signature at MiniBooNE

- Oscillation signature is charged-current quasielastic scattering:



- Dominant backgrounds to oscillation:

- Intrinsic  $\nu_e$  in the beam




- Particle misidentification in detector

Neutral current resonance:




# MiniBooNE Beamline



- 8 GeV primary protons come from Booster accelerator at Fermilab
- Booster provides about 5 pulses per second,  $5 \times 10^{12}$  protons per  $1.6 \mu\text{s}$  pulse under optimum conditions
- Beryllium target, single 174 kA horn
- 50 m decay pipe, 91 cm radius, filled with stagnant air




# MiniBooNE neutrino detector




- Pure mineral oil
- 800 tons; 40 ft diameter
- Inner volume: 1280 8" PMTs
- Outer veto volume: 240 PMTs

# Cherenkov ring characteristics: muons



- Muons have sharp filled in Cherenkov rings.


# Cherenkov ring characteristics: electrons



- Electrons undergo more scattering and produce “fuzzy” rings.

# Cherenkov ring characteristics:

$\pi^0$



- $\pi^0$  decay to  $\gamma\gamma$  with 99% branching ratio.
- Photon conversions are nearly indistinguishable from electrons.

# MiniBooNE's track-based reconstruction

---

- A detailed analytic model of extended-track light production and propagation in the tank predicts the probability distribution for charge and time on each PMT for individual muon or electron/photon tracks.
- Prediction based on seven track parameters: vertex  $(x,y,z)$ , time, energy, and direction  $(\theta, \varphi) \Leftrightarrow (U_x, U_y, U_z)$ .
- Fitting routine varies parameters to determine 7-vector that best predicts the actual hits in a data event
- Particle identification comes from ratios of likelihoods from fits to different parent particle hypotheses

# Beam/Detector Operation

---

- Fall 2002 - Jan 2006: Neutrino mode (first oscillation analysis).
- Jan 2006 - 201?: Antineutrino mode
  - (Interrupted by short Fall 2007 - April 2008 neutrino running for SciBooNE)
- Present analyses use:
  - $\geq 5.7E20$  protons on target for neutrino analyses
  - $5.66E20$  protons on target for antineutrino analyses
  - Over one million neutrino interactions recorded: by far the largest data set in this energy range

# Neutrino scattering cross-sections

---


- To understand the flavor physics of neutrinos (*i.e.* oscillations), it is critical to understand the physics of neutrino interactions
- This is a real challenge for most neutrino experiments:
  - Broadband beams
  - Large backgrounds to most interaction channels
  - Nuclear effects (which complicate even the definition of the scattering processes!)

# Scattering cross-sections

## for $\nu_\mu$


The state of knowledge of  $\nu_\mu$  interactions before the current generation of experiments:

- Lowest energy (  $E < 500$  MeV ) is dominated by CCQE.
- Moderate energies (  $500$  MeV  $< E < 5$  GeV ) have lots of single pion production.
- High energies (  $E > 5$  GeV ) are completely dominated by deep inelastic scattering (DIS).
- Most data over 20 years old, and on light targets (deuterium).
- Current and future experiments use nuclear targets from C to Pb; almost no data available.







# Dominant interaction channels at MiniBooNE




Charged-current quasielastic


Charged-current  $\pi^+$  production




Charged-current  $\pi^0$  production




Neutral-current  $\pi^0$  production




Neutral-current elastic



# Dominant interaction channels at MiniBooNE





Charged-current quasielastic




Charged-current  $\pi^0$  production

Charged-current  $\pi^+$  production



Neutral-current  $\pi^0$  production



Neutral-current elastic

MiniBooNE has measured cross-sections for all of these exclusive channels, which add up to 89% of the total event rate

CCQE (44%)

CC  $\pi^0$  (4%)

CC multi- $\pi$  (3%)

NC  $\pi^0$  (5%)

Others (4.1%)

NC multi- $\pi$  (2%)

NC EL (17%)

DIS (0.4%)

CC  $\pi^+$  (19%)

CC  $\pi^0$  (19%)

# MiniBooNE cross-section measurements


---

- ~~NC  $\pi^0$~~
- CC  $\pi^0$
- CC  $\pi^+$
- CC Quasielastic
- ~~NC Elastic~~
- ~~CC Inclusive~~

Due to limited time, only discussing a few topics here.



# Charged-current $\pi^0$ production

- Least common interaction for which we do exclusive measurement
- Uniquely, proceeds only via resonance:  
 $\nu + n \rightarrow \mu + \Delta \rightarrow \mu + p + \pi^0$
- Challenging 15-parameter, 3-ring fit needed:
  - Event vertex:  $(x, y, z, t)$
  - Muon:  $(E, \theta, \varphi)$
  - 1st photon:  $(E, \theta, \varphi, s)$
  - 2nd photon:  $(E, \theta, \varphi, s)$
- Relatively high backgrounds (mostly  $CC\pi^+$  which we measure separately)





# Reconstructed CC $\pi^0$ signal candidates

- Two-photon invariant mass  $m_{\gamma\gamma}$  allows very effective identification of events with a  $\pi^0$
- Reconstruction of full event allows observation of  $\Delta$  resonance



NUANCE is the default MiniBooNE neutrino interaction generator

# Measured observable $CC\pi^0$ cross-section



Additionally, we measure differential cross-sections vs:

- $\theta_\mu$
- $\theta_\pi$
- $E_\mu$
- $E_\pi$

- The dominant error is  $\pi^+$  charge exchange and absorption in the detector.
- First-ever differential cross-sections on a nuclear target.
- The cross-section is larger than expectation for all energies.
- Phys.Rev.D83:052009,2011



# Charged-current $\pi^+$ production

---

- Second-largest interaction channel at MiniBooNE
- Can proceed via resonance  $\nu + N \rightarrow \mu + \Delta \rightarrow \mu + N' + \pi^+$  or by coherent nuclear scatter.
- Identified by observation of *two* stopped muon decays after primary event. Unique signature results in purest exclusive sample in MiniBooNE
- Pion reconstruction and  $\mu/\pi$  separation are challenging.

# Measured observable charged-current $\pi^+$ cross-sections


- Differential cross sections (flux averaged):
  - $d\sigma/dQ^2$ ,  $d\sigma/dE_\mu$ ,  $d\sigma/d\cos\theta_\mu$ ,  
 $d\sigma/d(E_\pi)$ ,  $d\sigma/d\cos\theta_\pi$ :
- Double Differential Cross Sections
  - $d^2\sigma/dE_\mu d\cos\theta_\mu$ ,  $d^2\sigma/dE_\pi d\cos\theta_\pi$
- Data  $Q^2$  shape differs from the model
- Phys.Rev.D83:052007,2011.





# Charged-current quasielastic scattering (CCQE)


- Lepton vertex well understood
- Nucleon vertex parametrized with 2 vector form factors  $F_{1,2}$  and one axial vector form factor  $F_A$
- Use relativistic Fermi gas model of nucleus;  $F_{1,2}$  come from electron scattering measurements
- Generally assume dipole form of  $F_A$ ; only parameter is axial mass  $m_A$  extracted from neutrino-deuteron scattering experiments: 2002 average  $M_A = 1.026 \pm 0.021$  GeV




$$F_A(Q^2) = -\frac{g_A}{\left(1 + \frac{Q^2}{M_A^2}\right)^2}$$

# CCQE fit results: $Q^2$ dependence

- Data are compared (absolutely) with CCQE (RFG) model with various parameter values
- We prefer larger  $m_A$  compared to  $D_2$  data
- Our CCQE cross-section is 30% above the world-averaged CCQE model (red).
- Model with CCQE parameters extracted from shape-only fit agrees well with overall event rate (to within normalization error).




# Comparisons to other experiments (carbon targets)



- Our data (and SciBooNE) appear to prefer higher  $M_A$  than NOMAD, but the disagreement is not very significant.
- Note that:
  - Our errors are systematic-dominated and grow at highest energies
  - NOMAD allowed maximum of two tracks in event: in principle, different processes may contribute to the two experiments' samples
- Possible explanation for what appears to be higher  $M_A$ : two-nucleon correlations: Martini *et al.*, PRC 80, 065501 (2009)

# Neutrino Oscillations: 2007 result

- Search for  $\nu_e$  appearance in the detector using quasielastic scattering candidates
- Sensitivity to LSND-type oscillations is strongest in  $475 \text{ MeV} < E < 1250 \text{ MeV}$  range
- Data consistent with background in oscillation fit range
- Significant excess at lower energies: source unknown, consistent experimentally with either  $\nu_e$  or single photon production




# Antineutrino Oscillations

---

- LSND was primarily an antineutrino oscillation search; need to verify with antineutrinos as well due to potential *CP*-violating explanations
- Analysis has same number of protons on target in antineutrino vs. neutrino mode, but...
  - Antineutrino oscillation search suffers from lower statistics than in neutrino mode due to lower production and interaction cross-sections
  - Also, considerable neutrino contamination ( $20 \pm 5$ )% in antineutrino event sample

# Oscillation Fit Method


- Simultaneous maximum likelihood fit to
  - $\bar{\nu}_e$  CCQE sample
  - High statistics  $\bar{\nu}_\mu$  CCQE sample
- $\nu_\mu$  CCQE sample constrains many of the uncertainties:
  - $\bar{\nu}_e$  and  $\bar{\nu}_\mu$  flux uncertainties:



- Cross section uncertainties (assume lepton universality)
- Background modes -- estimate before constraint from  $\bar{\nu}_\mu$  data (constraint changes background by about 1%)
- Systematic error on background  $\approx 10.5\%$  (energy dependent)


# Data in antineutrino oscillation search

- $475 \text{ MeV} < E < 1250 \text{ MeV}$ :
  - $99.1 \pm 9.8(\text{syst})$  expected after fit constraints
  - 120 observed
  - Raw “one-bin” counting excess significance is  $1.5\sigma$
- Also see small excess at low energy, consistent with neutrino mode excess if attributed to neutrino contamination in  $\bar{\nu}$  beam



# Electron antineutrino appearance oscillation results


- Results for **5.66E20 POT**
  - Maximum likelihood fit for *simple two-neutrino model*
  - Oscillation hypothesis preferred to background-only at 99.4% confidence level.
  - $E > 475$  avoids question of low-energy excess in neutrino mode.
  - Signal bins only:
    - $P_{\chi^2}(\text{null}) = 0.5\%$
    - $P_{\chi^2}(\text{best fit}) = \sim 10\%$
- *Phys. Rev. Lett.* 105, 181801 (2010)






# Future sensitivity in $\bar{\nu}$ data

- MiniBooNE has requested a total of  $1.5 \times 10^{21}$  POT in antineutrino mode
- Potential  $3\sigma+$  significance assuming best fit signal
- Systematics limit approaches above  $2 \times 10^{21}$  POT
- Run is underway: total  $0.86 \times 10^{21}$  collected as of May 2011




# New result: disappearance with SciBooNE as near detector



- SciBooNE: Scintillating bar detector (originally from K2K) was in the BooNE beamline in 2007-08 to measure cross-sections
- Can also be used as a near detector for MiniBooNE
- New result this month:  $\nu_\mu$  and  $\bar{\nu}_\mu$  disappearance searches using both detectors
- Mean baseline: 76m (SciBooNE), 520m (MiniBooNE): oscillation probabilities differ significantly for  $0.5 < \Delta m^2 < 30 \text{ eV}^2$


# SciBooNE constraint reduces error at MiniBooNE



- Flux errors become 1-2% level: negligible for this analysis
- Cross-section errors reduced, but still significant due to different kinematic acceptance.

# SciBooNE-MiniBooNE $\nu_\mu$ disappearance result

- No evidence for oscillations
- Limit is better than other experiments in 10-30  $\text{eV}^2$  region
- Analysis of antineutrino mode is underway



# Conclusions

---

- Cross-sections:
  - MiniBooNE has most precise measurements of top five interaction modes on carbon; only differential and double-differential cross-sections in some modes
  - Some disagreements with most common nuclear models
- Oscillation searches
  - Significant  $\nu_e$  and  $\bar{\nu}_e$  excesses above background are emerging in both neutrino mode and antineutrino mode in MiniBooNE
  - The two modes do not appear to be consistent with a simple two-flavor neutrino model
  - Antineutrino results still heavily statistics-limited; MiniBooNE plans to accumulate more data until the goal of  $1.5 \times 10^{21}$  protons on target is reached

