## Enhancement of Cosmic Rays from the Vela Region at E>10<sup>12</sup> eV

## The BASJE Collaboration

F. Kakimoto<sup>1</sup>, S. Ogio<sup>1</sup>, Y. Tsunesada<sup>1</sup>, H. Yoshii<sup>2</sup>, T. Kaneko<sup>3</sup>, N. Tajima<sup>4</sup>, K. Nishi<sup>4</sup>, Y. Tanimoto<sup>2</sup>, Y. Nagaso<sup>2</sup>, S. Nakamitsu<sup>2</sup>, Y. Kuwata<sup>2</sup>, K. Okamoto<sup>2</sup>, K. Murakami<sup>5</sup>, Y. Matsubara<sup>6</sup>, Y. Toyada<sup>7</sup>, Y. Mizumoto<sup>8</sup>, Y. Yamada<sup>4</sup>, S. Shimoda<sup>4</sup>, H. Nakatani<sup>4</sup>, E. Gotoh<sup>4</sup>, N. Gironda<sup>9</sup>, A. Velarde<sup>9</sup>, W. Verasquez<sup>9</sup>, P. Miranda<sup>9</sup>, O. Burgoa<sup>9</sup>, R. Bustos<sup>9</sup>, K. Choque<sup>9</sup>, and J.C. Perez<sup>9</sup>

Department of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan
Department of Physics, Ehime University, Matsuyama 790-8577, Japan
Department of Physics, Okayama University, Okayama 700-8530, Japan
The Institute of Physical and Chemical Research, Saitama 351-0198, Japan
Nagoya University of Foreign Studies, Aichi 470-0197, Japan
Solar-Terrestrial Environment Laboratory, Nagoya University, Aichi 442-8507, Japan
Faculty of General Education, Fukui University of Technology, Fukui 910-8505, Japan
National Astronomical Observatory, Tokyo 181-8588, Japan
Institute de Investigaciones Fisicas, Universidad Mayor de San Andres, La Paz, Bolivia

## **Abstract**

At 25<sup>th</sup> ICRC we reported enhancement of cosmic rays arriving from the Vela region observed with SAS array at Mt. Chacaltaya in Bolivia. The significance level of this enhancement is 4.3. We, however, could not conclude that this enhancement is caused by primary gamma rays, since the observed air showers not limited with less muons show the same enhancement. To confirm these results with improved statistics and to investigate the energy dependence of this enhancement, we have installed a new air shower array: MAS array. The threshold energy of MAS array is  $6 \times 10^{12}$  eV while that of SAS array is  $3 \times 10^{13}$  eV. The observation with MAS array was started at October 1998. In this report, we will present the preliminary result including new data. Furthermore, we will calculate the degree of enhancement contributed by cosmic rays originated at the Vela SNR to compare with the observed result.