



**Fermi National Accelerator Laboratory**

FERMILAB-PUB-86/73-T

May, 1986

**Studies of the Hydrodynamic Evolution  
of Matter Produced in Fluctuations in  $p - \bar{p}$  Collisions and  
in Ultra-relativistic Nuclear Collisions II:  
Transverse Momentum Distributions**

**L. D. McLerran**

Fermi National Accelerator Laboratory  
Theoretical Physics Department  
P. O. Box 500  
Batavia IL 60510

**M. Kataja and P. V. Ruuskanen**

University of Jyväskylä  
Jyväskylä, Finland

**E. von Gersdorff**

Department of Physics and Astronomy  
University of Minnesota, Minneapolis, MN 55455

May 20, 1986

Abstract: We study solutions to the hydrodynamic equations appropriate for ultra-relativistic nuclear collisions. We find that the matter produced in such collisions spends times  $t > 30$  fm/c at temperatures larger than 150 Mev. The transverse momentum of protons, kaons and pions is computed in the central region of ultra-relativistic nuclear collisions. Assuming Bjorken's initial conditions for the hydrodynamic equations, and a bag model equation of state, we show that the transverse momentum distribution as a function of  $\frac{dN}{dy}$  does reflect properties of the equation of state. We demonstrate that such a distribution approximately scales as a function of  $\frac{1}{A} \frac{dN}{dy}$ . The relation between  $p_t$  and  $\frac{dN}{dy}$  is shown to be significantly altered under different assumptions about the equation of state. The transverse momentum distribution of heavy hadrons is shown to be much enhanced relative to that of light pions. These distributions are little changed by differences in the assumptions about the initial transverse density and velocity profile. We are unable to fit the observed correlation between  $p_t$  and  $\frac{dE}{dy}$  observed in the JACEE experimental collaboration.

## Section 1: Introduction

In this second paper of a series of two, we use the methodology explained in the previous paper in an attempt to compute the transverse momentum distributions of hadrons as they might be produced in ultra-relativistic nuclear collisions. Arguments have been put forth by Shuryak<sup>(1)</sup>, Shuryak and Zhirov<sup>(2)</sup>, and by van Hove<sup>(3)</sup> that the transverse momentum as a function of multiplicity reflects general properties of the equation of state of high temperature hadronic matter. These arguments, when substantiated by detailed computation, might allow an experimental determination of the properties of high temperature hadronic matter.

The argument originally proposed by Shuryak is quite simple.<sup>(1-2)</sup> In the expansion of a fluid, the matter in the fluid does work on particles as they are pushed into the vacuum. This work is caused by a pressure difference between the vacuum and the matter in the fluid. As the energy density increases, the pressure increases, and therefore the transverse momentum of secondary particles increases. For example, the multiplicity of particles reflects the energy density achieved in a collision. Therefore, the degree to which the average transverse momentum increases as a function of multiplicity reflects properties of the equation of state.

The correlation between multiplicity and transverse momentum most dramatically reflects the equation of state in the case that there is a first order phase transition. As has been emphasized by van Hove,<sup>(3)</sup> the transverse momentum may not rise so steeply for energy densities appropriate for a first order phase transition. During a first order transition, the energy density increases during the mixed phase, while the pressure remains constant. For equations of state appropriate to describe quark-gluon matter, this region of constant pressure may occur for energy densities which vary by an order of magnitude. The transverse momentum distribution does not change much for this range of energy densities since the work done by the fluid on particles while expanding into the vacuum remains roughly constant, since the pressure

remains constant.

There are of course many reservations concerning the above argument. In fact for a quark gluon plasma, there is a slight increase in the energy per degree of freedom relative to that of an ideal pion gas. Probably the greatest gap in this argument however, and the gap which is hardest to fill without the detailed computations which we present, is the following. As the energy density increases, the system spends more time in a mixed phase than as a low pressure pion gas. Therefore, as the energy density increases, we expect a continued increase in the average transverse momentum due to a larger time averaged value of the pressure.

The correlation between transverse momentum and multiplicity may be most easily seen by considering the example of the expansion of a spherical droplet of fluid, initially at rest.<sup>(4-5)</sup> Such an initial condition might arise as a fluctuation in  $\bar{p}p$  collision. For such a system, initially in a volume  $V$ , the total energy and total entropy are conserved. Therefore, the quantities  $E/S$ , the energy per degree of freedom, and  $E/V$ , where  $V$  is the initial volume are conserved by the equation of motion. Up to a constant which may be determined from geometry and by relating entropy to multiplicity,  $E/S$  is the average transverse momentum. The total energy of the secondaries may also be measured, and if the initial volume is known, the initial energy density may be experimentally determined. Notice that the total energy is easily related to the total multiplicity once the energy per particle is known. The relation between  $E/S$  and  $E/V$  may also be theoretically determined. The result of such a computation for a bag model is shown in Fig. 1 At low densities,  $E/S \sim T$  since the energy per degree of freedom is that of a pion gas. At high densities,  $E/S$  is again  $\sim T$  with roughly the same proportionality constant, since the energy per degree of freedom is almost independent of the constituents for a massless ideal gas. At energy densities appropriate to the phase transition, the energy per degree of freedom remains approximately constant as the hadron gas melts into a plasma.

For ultra-relativistic nuclear collisions, the geometry is not so simple as was the case for  $p\text{-}\bar{p}$  collisions. For systems of infinite transverse extent, the solutions of Bjorken<sup>(6)</sup> determine the space-time evolution of matter in the central region. These equations may also be generalized to the fragmentation region.<sup>(7-8)</sup> Unfortunately, the issue of measuring the properties of the equation of state by studying the transverse momentum distribution of secondaries is not possible without allowing for transverse expansion. The formalism for treating transverse expansion of a quark-gluon plasma has been developed by Baym et. al.<sup>(9)</sup> and by Bialas and Czyz.<sup>(10)</sup> Their method is applicable for the central region of head on nuclear collisions of equal  $A$  nuclei. They have made explicit computations for the case of an ideal gas expansion.

The generalization of their results to the case where a fluid transversely expands and undergoes a phase transition is not entirely straight forward. As matter transversely rarefies into the phase transition region, a transverse rarefaction shock develops.<sup>(11-12)</sup> In order to treat this case, in a previous paper<sup>(13)</sup> we presented an algorithm originally due to Boris and Book<sup>(14)</sup> which has been applied in the past to relativistic nuclear collisions by workers at Frankfurt.<sup>(15-16)</sup> We also discussed the general initial conditions, the method of solution of hydrodynamic equations, and decoupling. In this paper we shall make extensive use of the results presented in the first paper, which we now refer to as I.

This second paper presents results of explicit solutions to the hydrodynamic equation appropriate for the conditions in the central region of ultra-relativistic nuclear collisions. In the second section, we discuss qualitative and quantitative features of our scenario for initial conditions and subsequent expansion of matter. We discuss the interplay between transverse and longitudinal expansion. We describe the shock wave which appears in the transverse rarefaction, and will show in the third section that our numerical solution properly reproduces the known features of this shock as gleaned from analytic studies. We also qualitatively analyze the correlation between transverse momentum and multiplicity for various nuclear sizes  $A$ .

In the third section, we present the results of our computations for the hydrodynamic evolution of matter as it might be produced in ultra-relativistic nuclear collisions. We explicitly compute the  $p_t$  distributions of hadrons. We argue that the correlation between  $p_t$  and  $\frac{dN}{dy}$  is fairly independent of the details of the assumptions concerning the matter distributions and the initial conditions. We show that this correlation does in fact measure properties of the equation of state. We show that the correlation approximately scales as a function of  $\frac{1}{A} \frac{dN}{dy}$ . We also compute the effects of a quark-gluon plasma upon the kaon and proton spectrum. We present the actual  $p_t$  distribution for pions for a variety of values of  $\frac{dN}{dy}$  and for pions, kaons and nucleons together for a typical value of the multiplicity density in head on U-U collisions.

In the fourth section, we attempt to compare our results with the experimental results from the JACEE emulsion chamber experiment. We find that our results are unable to fit the rapid rise in  $p_t$  seen in this experiments. We attempt to critically assess this discrepancy.

In the last section, we briefly summarize our results.

## Section 2: Initial Conditions, and Qualitative Features of the Flow

We shall give in this section a qualitative and semi-quantitative discussion of how the matter flows in ultra-relativistic nuclear collisions, and of qualitative features of the  $p_t$  spectrum of hadrons as a function of  $\frac{dN}{dy}$ . We also discuss the expansion time of the matter produced in these collisions and the shock wave which appears in the transverse rarefaction of the matter. In addition, the interplay between transverse and longitudinal expansion, as well as general features of the correlation between multiplicity and average transverse momentum are described.

We will consider a situation where the hot hadronic matter expands

longitudinally according to the scaling solution but is initially at rest in the transverse direction. Because of pressure gradients, transverse flow will develop, the details depending on the initial conditions, the equation of state and on other assumed dynamical properties like the nucleation rate if a first order phase transition is possible. Here we shall assume the bag model equation of state and a 'fast nucleation scenario' which means that the phase transition is assumed to proceed through the equilibrium mixed phase where pressure stays constant. This has a profound effect on the evolution of transverse flow and consequently the resulting  $p_t$ -distribution is not related in a simple way to the initial temperature  $T_i$  or the initial energy density  $\epsilon_i$ .

During the expansion energy is lost from any slice of matter in a given rapidity interval  $dy$  through the work which is done in the longitudinal expansion. Since for an ideal massless relativistic gas, entropy conservation implies gas particle number conservation in this slice, the energy per particle will decrease. In principle mass effects in the pion gas phase could lead to decrease in particle number but the processes to achieve this are probably too slow when the pion mass is important.<sup>(17)</sup> This might be also compensated by the entropy production during various stages of the expansion. In terms of  $\langle p_t \rangle$ , the coupling to the longitudinal expansion means that the final value is usually less than what one would expect on the basis of initial temperature.

The effect of the phase mixture on the flow is two-fold. On one hand it slows down the transverse expansion as compared to the ordinary transverse rarefaction wave in simple relativistic fluid, leading to longer decoupling times. On the other hand when the initial energy density is in the mixed phase interval, it provides a mechanism to drive the transverse expansion through the formation of a rarefaction shock wave at the interface of mixed phase and hadron gas. Since the strength of the shock depends on the energy density of the mixed phase, the flow velocity at decoupling increases with increasing energy density even though the initial temperature is fixed to  $T_c$  in this interval. As a result the  $p_t$ -distribution broadens and  $\langle p_t \rangle$  increases

throughout the whole phase transition region contrary to the naive expectations.

This conclusion should hold even in the case without homogeneous mixed phase when the initial matter would consist of plasma droplets embedded in hadron gas. As the droplets deflagrate the resulting pions will get the vacuum energy of bag constant in addition to the thermal energy and consequently their average energy will be higher than that of the pions originally in hadron phase. It should be noted that even though  $\langle p_{\perp} \rangle$  grows throughout the phase transition region  $T_i = T_c$ , the growth is much slower than in the case of no phase transition when the initial matter would consist of pions.

When  $T_i > T_c$  the temperature drops to  $T_c$  in time  $\tau_Q = \tau_i (T_i/T_c)^3$  due to the longitudinal expansion alone. If  $\tau_Q \ll 3^{1/2} R_A$  the ordinary rarefaction wave has little effect and from  $\tau_Q$  on the system developed essentially as if the initial energy density had been  $\epsilon_Q$ . Consequently the growth of  $\langle p_{\perp} \rangle$  is slow until  $\tau_Q$  becomes comparable with  $3^{1/2} R_A$ . Then the ordinary rarefaction wave has time to evolve and a strong transverse flow builds up. Depending on the details of the decoupling this may even lead to shorter decoupling times for some parts of the matter when initial temperature increases.

The above arguments about the decrease of average energy per particle in the final state due to longitudinal expansion hold only for pions. For the heavier particles the situation may be different if they are able to stay in local thermal equilibrium and follow the collective flow. This is because their thermal motion even in the high temperature initial state may, due to their mass, be small compared to the velocity of final flow which is dominated by the (almost) massless pions. As a result the  $p_{\perp}$ -distribution of heavy particles should broaden more than that of the pions when going from hadron collisions to heavy ion collisions. This expectation is verified by numerical results.

### Section 3: Results and Their Dependence on the Parameters



In this chapter we present the numerical results on the flow and the  $p_t$ -distributions and discuss the details of their dependence on the initial conditions and the main parameters of the model, the critical temperature,  $T_c$ , the decoupling temperature,  $T_{dec}$ , and the radius of the colliding nuclei,  $R_A$ . The number of quark flavors is taken to be 2.5 to account roughly for the mass of strange quark. With pions alone in the hadron phase, the ratio of the number of degrees of freedom in the plasma and hadron gas is 14.1. Most results are for  $T_c = 200$  MeV which gives  $B = 0.8$  GeV/fm<sup>3</sup> for the bag constant. Initial time  $\tau_i$  and temperature  $T_i$  are related through  $\tau_i = C/T_i$  where  $C = 250$  MeV-fm/c as explained in I.

It actually turns out that for the flow and the hadronic  $p_t$ -distributions the values of  $\tau_i$  and  $T_i$  separately are not very important when  $T_i > T_c$ . The primary quantity is  $\tau_i T_i^3$  which is a measure of total entropy  $\frac{dS}{dy}$  or, because of the approximate entropy conservation during the expansion, of the total multiplicity  $\frac{dN}{dy}$ . Incidentally,  $\tau_i T_i^3$  fixes also the value of  $\tau_Q$ , which is the essential timescale in the build up of the transverse flow. Since in most of the cases we consider,  $\tau_Q$  is much less than the transverse size of the system, there is little transverse expansion before forming a mixed phase, and therefore little dependence in the flow upon  $T_i$  or  $\tau_i$ . The value of  $T_i$  is on the other hand essential from the point of view of dilepton production and possible other signals which depend on the whole space-time history of the collision.<sup>(18)</sup> Here the discussion of dependence of different quantities on initial conditions can always be taken to refer to the total multiplicity.

If the initial energy density is less than  $\epsilon_H$ , the critical energy density of the hadron gas, transverse flow will be weak. All that happens is that longitudinal expansion cools the matter and the  $p_t$ -distribution is essentially the thermal distribution at  $T_{dec}$  with  $\langle p_t \rangle = 2.12 T_{dec}$ . When  $\epsilon_i$  grows across the phase transition region from  $\epsilon_H = 0.2$  GeV/fm<sup>3</sup> to  $\epsilon_Q = 3.8$  GeV/fm<sup>3</sup>, the rarefaction shock gets stronger, the flow velocity of the outgoing hadrons increases and the  $p_t$ -distribution broadens. Figs. 2a and b show the contour plots of the

energy density and the radial velocity for the flow when  $\epsilon_i = \epsilon_Q$ . The shock front follows closely the  $\epsilon = \epsilon_H$  contour and is clearly seen as a jump in radial velocity. As the longitudinal expansion dilutes the energy density the velocity of the shock front increases<sup>(11)</sup> and this is seen as a bending of the shock region towards the time axis.

In Figs. 3a and b the same plots are shown for  $T_i = 350$  MeV. One can see how the ordinary rarefaction wave starts from the surface but since  $\tau_Q = 3.8$  fm/c, it can not reach the inner parts of the matter before turning into the slowly propagating rarefaction shock. As a result the transverse flow is not dramatically increased from that in Fig. 2 and the growth of  $\langle p_t \rangle$  is moderate as will be seen soon. This is possible because the decoupling time increases and the matter can lose a large part of its initial thermal energy as work in the longitudinal expansion. From here on the ordinary rarefaction becomes more important and  $\langle p_t \rangle$  starts to grow faster with multiplicity.

To exhibit how the phase transition affects the flow, we show in Fig. 4 the flow for the pion gas with the same multiplicity as for the flow in Fig. 2. In the pion gas, the transverse expansion is faster and a much stronger flow builds up than is the case for a mixed phase. This shows up quantitatively in the values of the average transverse momentum which for the pions in these two cases are 495 and 690 Mev with and without a phase transition.

From the flow quantities the  $p_t$ -distributions for the pions can be calculated as explained in I and for the massless pions the results as function of multiplicity are shown in Fig. 5. Qualitatively the distributions for largest multiplicities differ from thermal distributions in the sense that they do not exhibit exponential falloff as the thermal distributions would do in the shown  $p_t$ -range.

The effect of the phase transition on the dependence of the  $p_t$  spectra is illustrated in Fig. 6 where the average value of  $p_t$  is shown both with and without phase transition as a function of multiplicity for iron-iron collisions with  $R_A = 4.2$  fm. The rapid growth in the case

If no phase transition reflects the fact that with the pion degrees of freedom available only, high initial temperatures are needed to raise the multiplicity. The highest point corresponds to temperature  $T_i = 550$  MeV at  $\tau = 1$  fm/c. According to the present ideas about the reasonable values of  $T_c$ , this is probably not a realistic temperature for the hadron phase. We return to this question later when discussing the SACCIE data.<sup>(19)</sup> The fact that in the case of a phase transition the growth of  $\langle p_t \rangle$  is slowest around  $\epsilon_i = \epsilon_Q$  shows how effective the rarefaction shock is in delivering energy into the transverse flow. When the shock attains its full strength at  $\epsilon_i = \epsilon_Q$  (the place is indicated by  $\epsilon_Q$  in Fig. 6) the growth of  $\langle p_t \rangle$  almost stops until the ordinary rarefaction wave in the plasma has enough time to develop.

Next we consider the effect of the size of the colliding nuclei on  $\langle p_t \rangle$ . It turns out that  $\langle p_t \rangle$  as a function of  $\frac{1}{A} \left( \frac{dN}{dy} \right)$  exhibits approximate scaling. Qualitatively this can be understood as follows: For large  $A$ , a fixed value of  $\frac{1}{A} \left( \frac{dN}{dy} \right)$  implies larger initial temperatures than for small  $A$ . On the other hand the transverse flow develops faster and decoupling is reached sooner for small  $R_A$ , so that longitudinal expansion has less of a diluting effect upon the transverse momentum distribution. The cancellation of these opposite effects then leads to the aforesaid scaling which is illustrated in Fig. 7.

It should be emphasized that in a scenario where the matter undergoes scaling expansion in the longitudinal direction, this expansion has a crucial effect on the final values of  $\langle p_t \rangle$ . In the case of no transverse motion the final  $p_t$ -distribution is always the thermal distribution at  $T = T_{dec}$  and the role of the longitudinal motion is to cool the matter to this temperature through the work in the expansion. Even with transverse motion taking place some part of the initial thermal energy is always lost into the longitudinal work. As a result  $\langle p_t \rangle$  is, in general, reduced during the hydrodynamic stage from its initial value, which for massless quanta is  $\langle p_t \rangle = 2.12 T_i$ . The effect of the longitudinal work in our explicit calculations is shown in Fig. 8 where the final values of  $\langle p_t \rangle$  are plotted together with  $\langle p_t \rangle$  of the

initial thermal distribution. As explained earlier, there are uncertainties in the initial time and consequently in the initial temperature, which would be reflected as an uncertainty in the initial  $\langle p_t \rangle$ . Still since we are looking  $\langle p_t \rangle$  as a function of multiplicity which means fixing of  $\tau_i T_i^3$ , a factor of two uncertainty in  $\tau_i$  would cause a 20 % uncertainty in  $T_i$ , only.

In the actual nuclear collisions at high but finite energies the longitudinal work should result as an energy transfer from central region to the fragmentation regions<sup>(8)</sup> and should be studied through the systematics of the  $A$  ( $R_A$ ) and  $\frac{dN}{dy}$  dependence of the shapes of both  $p_t$  and  $y$ -distributions. Note that in spherical expansion  $\langle p_t \rangle$  is conserved as a consequence of the symmetry throughout the expansion. In that situation  $\langle p_t \rangle$  directly reflects  $T_i$ . Such situations might occur due to fluctuations in the initial conditions. On the other hand the possible high  $T_i$  fluctuations with small or zero velocity (longitudinal) gradient may not be spherically symmetric. Instead they might exhibit Landau type of initial conditions which would still lead to faster expansion in the longitudinal direction. Since, especially for large nuclei, one would expect the spherical fluctuations to be embedded in longitudinally expanding matter and interacting with it, it may be difficult to understand large  $\langle p_t \rangle$  values from the point of view of hydrodynamics.

In this study we assume that the density of heavier particles is low enough so that their effect on hydrodynamic evolution of the matter can be neglected. This seems reasonable because they affect only the hadron gas stage of the hydrodynamic expansion and because of the mass suppression one would expect the effect to be less than 10 % to the equation of state. (The fraction of kaons between  $T_c$  and  $T_{dec}$  is of the order of 30 % or less and for that part the pressure could be lowered roughly by 30 % giving a combined effect of  $.3 \times .3 = .1$  or 10 %). If we further assume that the heavy particles follow the overall flow, we obtain the  $p_t$ -distributions of kaons and nucleons by simply calculating the flux of  $n_i u^\mu$  through the decoupling surface as explained in detail in I.  $n_i$ ,  $i=K,N$ , are the ideal gas densities with zero chemical

potential for the kaons and nucleons at the decoupling temperature and  $u^\mu$  is the four velocity of flow.

The results are shown in Fig. 9 together with the  $p_t$ -distribution of massless pions for U-U collisions with  $\frac{1}{A} \left( \frac{dN}{dy} \right) = 13.5$ . The normalization corresponds to one internal degree of freedom for each particle species. The spectra differ qualitatively from each other showing clearly the strong boost given by the transverse flow to the heavy particles. The dependence of  $\langle p_t \rangle$  on the (total) multiplicity is shown in Fig. 10 for the three particle species exhibiting the stronger growth of  $\langle p_t \rangle$  for the heavier particles. These features of the dependence of  $p_t$  on the mass of the particle and the multiplicity of events provide an experimental test for the existence of the collective flow in the final state of heavy ion collisions.

In our computations, the kaon and nucleon distributions which are used to compute the emission of these particles through the decoupling surface were taken to be ideal gas distribution functions with zero chemical potential. We might have allowed for a constant chemical potential which would change the relative normalization of the presented distributions. It would however change neither the shape of the distributions nor the average transverse momentum. If, as is expected for kaons, the chemical potential for strangeness is time dependent, then this algorithm is not strictly correct. The qualitative and semi-quantitative nature of the enhancement of heavy particles at large  $p_t$  is quite general, and we expect that our approximate treatment reflects this effect

We next consider how the results depend on the different basic parameters, which up to now have been kept fixed.

(i) Decoupling temperature  $T_{dec}$ :

When  $T_{dec}$  is lowered to 100 MeV there is a small increase (10 MeV/c or less) in  $\langle p_t \rangle$  except for the lowest initial energy density  $\epsilon_i = \epsilon_f$  where the longitudinal cooling still dominates over the transverse

cooling. Compared to the other uncertainties, those coming from the uncertainty of the value of  $T_{dec}$  can be considered unimportant.

(ii) Critical temperature  $T_c$ :

The value of critical temperature affects the  $p_t$ -distributions in two ways, it changes the timescales and contributes to the initial energy density (for fixed multiplicity) through the bag constant. Table I shows the situation in the case of  $T_i=500\text{MeV}$ . These values correspond to the same initial entropy and essentially the same  $\frac{1}{A} \left(\frac{dN}{dy}\right)$ . (The final values of  $\frac{dN}{dy}$  differ a little because the details of the flow differ and this results in small differences in entropy production associated with the shock wave rarefaction.)

One can see from these results that  $\langle p_t \rangle$  is not very sensitive to the precise value of  $T_c$ . In order to get significant increase in  $\langle p_t \rangle$  the value of  $T_c$  must be given values which are quite high at least from the point of view of using the simple bag model equation of state. E.g.  $T_c = 400\text{MeV}$  leads to  $B = 14 \text{ GeV}/\text{fm}^3$ , and at such high temperatures pions alone would not give a good approximation of the (degrees of freedom of) hadron gas.

(iii) Initial time  $\tau_i$  and temperature  $T_i$ :

As has been already discussed, the changes of  $\tau_i$  and  $T_i$  are not important as long as  $\frac{dN}{dy} \sim \tau_i T_i^3$  is kept fixed. At the initial time  $v_r=0$  and since  $\tau_0$  is now fixed, the differences for different  $\tau_i$  arise from the evolution of the flow in the time interval  $\delta\tau_i$  which one would expect to be of the order of 1 Fm or less. This is a short time interval from the perspective of the total expansion time.

(iv) Shape of the initial densities:

If the initial temperature distribution is not a step function, there will be a transverse pressure gradient (except where  $T = T_c$ ) and the transverse acceleration takes place also in the matter. This will

strengthen the flow and shorten the decoupling time. Taking the initial temperature distribution as

$$T(r) = T_0(1-(r/R_A)^2)^\kappa \quad (1)$$

We have considered three cases with  $\kappa = 0, 1/6$  and  $1/3$ . Roughly speaking for  $\kappa = 1/6$ , the production of final state matter would be proportional to the number and for  $\kappa = 1/3$  to the square of the number of colliding pairs of nucleons at the distance  $r$  from the collision axis. For  $R_A = 6.8$  Fm and  $\frac{1}{A} \frac{dN}{dy} = 27.5$ ,  $T_0 = 500, 572$ , and  $630$  Mev and the resulting values of  $\langle p_t \rangle$  are  $630, 657$  and  $697$  Mev/c respectively. The effect is quite moderate.

(v) The effect of initial transverse velocity:

In all the results which have been discussed above  $v_r = 0$ , initially. It is clear that  $\langle p_t \rangle$  can be arbitrarily enhanced if strong initial transverse flow is assumed. In the head on collisions this, however, does not sound physically justified; one would rather expect that with the exception of the outer surface, the produced matter would be transversally at rest. A calculation with  $\kappa = 1/6$ ,  $R_A = 6.8$  Fm,  $\frac{1}{A} \frac{dN}{dy} = 27.5$  and initial  $v_r$  growing linearly from 0 at  $r = 0$  to  $0.3$  at the surface gives  $\langle p_t \rangle = 765$  Mev as compared to  $660$  MeV for  $v_r = 0$  at  $r = r_1$ . While it is conceivable that the initial flow might be stronger close to the surface but at small and intermediate values of  $r$ , it is difficult to understand how in head on collisions, a significant collective motion in the transverse direction could be present initially. Calculations with different multiplicities show that the effect grows with multiplicity. The change in numbers is less than 20% for the above numbers with rather high multiplicity.

(vi) The effects due to uncertainties in decoupling:

Fig. 11 shows the  $\langle p_t \rangle$  from three different decoupling calculation. In flow decoupling the contribution from the collective motion to  $\langle p_t \rangle$  is included, only. As a result it gives smaller values than the

particle decoupling. The dependence on increasing initial temperature is, however similar in all cases and the flow decoupling gives a nice independent check of the calculations. The third curve in the Fig. 11 corresponds to pions with mass. Strictly speaking it is not consistent to include the mass in the decoupling calculation since the hydrodynamics is calculated with massless pions. However, the effect of the mass to the equation of state and the hydrodynamic calculation is small but is not that negligible to  $\langle p_t \rangle$  (of the order of 15% to the right direction from the point of view of JACEE data). Most of this effect is due however to the fact that for massive pions, the relationship between multiplicity and entropy is altered. This effect is presumably less important if pion number is treated as conserved when the effect of pion mass is important.

In summary, we have shown that there is a strong correlation between  $\langle p_t \rangle$  and  $\frac{dN}{dy}$  which reflects properties of the equation of state of hadronic matter. We have shown that these correlations may be computed up to a fair number of theoretical uncertainties. These uncertainties are all at the 10%-20% level or less, and hopefully the sum of their contributions is equally small. In the future, computations may allow a more precise determination of these uncertainties, and a hopefully more reliable estimate of the correlation. Given the stability of our numerical results to such changes, we believe that this correlation will survive closer inspection.

#### Section 4: Comparison with JACEE Data and Discussion

The data taken by the JACEE cosmic ray collaboration for  $p_t$  vs  $dE_t/dy$  is shown in Fig. 12. We have here only included the data for nucleus-nucleus collisions. In this plot, the energy density  $\langle \epsilon \rangle$  at initial time  $\tau_0 = 1\text{fm}/c$  is defined as

$$\langle \epsilon \rangle = \frac{3}{4} \pi \langle m_t \rangle \frac{dN}{dy} |_{ch} A_{\min}^{-2/3} \quad (2)$$

according to the formula of Bjorken.<sup>(6)</sup> All quantities in this



relationship are experimentally measurable parameters. In order to compare our hydrodynamic results with these data, we have calculated from the numerical results of  $\langle p_t \rangle$  and  $\frac{dN}{dy}$  for massless pions with different initial conditions an 'initial energy density' at  $\tau_0 = 1$  fm/c according to the above formula. It should be noted that as compared to the hydrodynamic calculation, this result always underestimates the initial energy density because it has no reference to the work during the expansion. Since the data is mainly for light nuclei, we use  $R_A = 4.2$  fm.

At a value of  $\langle \epsilon \rangle$  of the order of 1-10 Gev/fm<sup>3</sup>, there is a sharp break in the  $p_t$  distribution. This break occurs in a plot which involves a variety of nuclei, and in fact primarily light nuclei. The interpretation of the lower axis as an energy density is therefore not on a very sound footing. Nevertheless, the occurrence of this break at an 'energy density' where one expects a quark-gluon plasma, and the above arguments for enhancements in the  $p_t$  spectrum due to hydrodynamic flow make it tempting to interpret this plot as evidence for the existence of a quark-gluon plasma.

In Ref. 4, it was argued that such a curve might be fit if it was assumed that the plasma was formed as a result of spherically symmetric fluctuations in a volume of radius  $1/p_t$ . While such a model does fit the data on  $p_t$  vs  $\langle \epsilon \rangle$ , it suffers from lack of evidence that the distribution appears in any sense to correspond to a spherical fluctuation. Also even for light nuclei, it is not clear why small volume (as compared to the nuclear size) fluctuations would dominate the whole event. We shall therefore in this paper, try to analyze the JACEE data assuming a boost invariant matter distribution, as is expected to be the case for typical collisions of ultra-relativistic heavy nuclei.

The result of such an attempted comparison is shown in Fig. 12. As can be seen by the plot, the JACEE data rises too sharply to for such interpretation. Inclusion of the effects from pion mass and initial transverse velocity would enhance the calculated curve, but the shape

of the curve would be almost unaffected. It is obviously the steep rise of  $\langle p_t \rangle$  in a very narrow range of  $\epsilon$  which is very difficult to obtain in scaling hydrodynamics with phase transition. There is some possibility that the data might be fit by an ideal pion gas, with no plasma, as shown in the figure. At the energy densities of interest, the use of an ideal pion gas equation of state is probably not valid, since the pions most likely strongly interact, and contributions from strange particles and resonances are important.

There are several reasons why such a comparison might not be meaningful. First, in the absence of more detailed data which would establish whether or not the matter produced in such collisions was fluctuation dominated or not, we do not know if for these small nuclei, we are in fact using a correct space-time model to describe the collisions. Also, it has not been established that for such small nuclei, there is any evidence for hydrodynamic flow.

Another possibility is that the physics which produces this large  $p_t$  enhancement is not that of a quark-gluon plasma. For example, mini-jets may be important. To check this possibility, the azimuthal angular distribution of matter produced in such processes must be studied in detail, and systematic comparison of model computations which include jets must be made.

It is tempting to assume that perhaps the particles which generate the  $p_t$  enhancement are particles heavier than pions, and therefore acquire a larger  $p_t$  enhancement. Unfortunately, the experiment measures the distribution of photons to compute  $dE_t/dy$ . These photons should arise primarily from the decay of  $\pi^0$ s. If there was another substantial source of photons, it might be possible to explain the data.

Another possibility which we do not consider is that the JACEE data is incorrect.

To summarize, using a conventional space-time picture of heavy ion

collisions, we are unable to explain the observed correlation in the AACEE data between  $p_t$  and  $\frac{dN}{dy}$ .

## Section 5 Conclusions

In this paper, we have applied a conventional space-time picture of heavy ion collisions together with a hydrodynamic computation of the matter produced in such collisions, to describe the transverse momentum of particles, and to semi-quantitatively determine the lifetime of this matter. We have found that the matter probably lives long enough, at least for heavy nuclei, to justify such a treatment. We have also shown that the transverse momentum distributions do in fact reflect properties of the equation of state. It is of course premature to claim that these computations present signals for a quark-gluon plasma because there are many unforeseen possibilities. On the other hand, if there is any hope to obtain information on the equation of state from heavy ion collisions, collective phenomenon must be observed experimentally. This makes the systematic study of the dependence of the  $p_t$  spectra on multiplicity and the size of the colliding nuclei a very important experimental issue. Differences in the behaviour of the spectra between pions and heavier particles should also show up if the collective flow exists. In order to achieve the above goals, a good understanding of background processes is necessary.

Our conclusions on these points leave some room for optimism, but it would be useful to have further theoretical studies in several areas to make the conclusions firmer. For example, to understand the possibilities for finding evidence of collective flow, a truly 3+1 dimensional code should be employed to study the correlations between the central region and the fragmentation regions or between the transverse momentum and rapidity distributions. Also an equation of state which allows for massive pions is needed because the issue of decoupling is not clearly defined. More realistic equations of state should be used to further estimate the uncertainties in these computations. There should also be cascade analysis. Such an analysis might determine the degree to which the plasma evolution is fluctuation

dominated or a smooth hydrodynamic evolution. The effect of density fluctuations due to phase separation in the mixed phase could be determined.

References:

- (1) E. V. Shuryak, Phys. Rep. 61, 71 (1980).
- (2) E. V. Shuryak and O. Zhironov, Phys. Lett. 89B, 352 (1978); Sov. J. Nucl. Phys. 28, 247 (1978).
- (3) L. van Hove, Phys. Lett. 118B, 138 (1982).
- (4) E. von Gersdorff, J. Kapusta, L. McLerran and S. Pratt, Phys. Lett. 163B, 253 (1985).
- (5) K. Redlich and H. Satz, University of Bielefeld Preprint (1985).
- (6) J. D. Bjorken, Phys. Rev. D27, 140 (1982).
- (7) K. Kajantie and L. McLerran, Phys. Lett. 119B, 203 (1982); Nucl. Phys. B214, 261 (1983).
- (8) K. Kajantie, R. Raitio, and P. V. Ruuskanen, Nucl. Phys. B222, 52 (1983).
- (9) G. Baym, B. Friman, J.-P. Blaizot, M. Soyeur and W. Czyz, Nucl. Phys. A407, 541 (1983).
- (10) A. Bialas and W. Czyz, Acta. Phys. Pol. B15, 229, (1984).
- (11) B. Friman, K. Kajantie, and P. V. Ruuskanen, Nucl. Phys. B266, 68 (1986).
- (12) S. Pratt, University of Minnesota Preprint June 1985.
- (13) E. von Gersdorff, M. Kataja, P. V. Ruuskanen and L. McLerran, Fermilab Preprint Fermilab-Pub-86/13-T (1986).
- (14) J. P. Boris and D. L. Book, Journal of Comp. Phys. 11, 38, (1973); 20, 397 (1976); D. L. Book, J. P. Boris and K. Hain, Journal of Comp. Phys. 18, 248 (1975).
- (15) G. Graebner, Frankfurt University PhD Thesis.
- (16) G. Buchwald, G. Graebner, J. Theis, S. Maruhn, W. Greiner and H. Stoecker, Phys. Rev. Lett. 52, 1594 (1984); Nucl. Phys. A418, 625, (1984).
- (17) S. Gavin, Nucl. Phys. A435, 826 (1985).
- (18) K. Kajantie, M. Kataja, L. McLerran, and P.V. Ruuskanen, Helsinki University Preprint HU-TFT-86-6 (1986).
- (19) T. H. Burnett et al. Phys. Rev. Lett. 50, 2062, (1983).

Figures :

Fig. 1 E/S vs E/V for the bag model equation of state with the parameters given in Section 3.

Fig. 2 Curves of (a) constant energy density (GeV/fm<sup>3</sup>) and (b) constant radial velocity for  $T_i = T_c = 200\text{MeV}$  and  $\epsilon_i = \epsilon_Q = 3.8\text{ GeV/fm}^3$ . The velocity curves near the light cone are an artifact of numerical approximations and have no physical significance.

Fig. 3 As Fig. 2 but for  $T_i = 350\text{ MeV}$  and  $v_i = 0.71\text{ fm/c}$  corresponding to  $\epsilon_i = 28\text{ GeV/fm}^3$ .

Fig. 4 As in Fig. 3 but without phase transition. To achieve the same multiplicity  $\frac{1}{A} \frac{dN}{dy} = 4.2$  with the pion gas, the initial temperature is  $T_i = (14.1)^{2.5} T_c = 480\text{ MeV}$ .

Fig. 5  $p_t$  distributions for (massless) pions in U-U collisions ( $R_A = 6.8\text{ fm}$ ) for different values of multiplicity  $\frac{1}{A} \left(\frac{dN}{dy}\right)$ .

Fig. 6 Average  $p_t$  with and without phase transition. In the latter case the matter is assumed to be produced as hot pion gas. At the point indicated by  $\epsilon = \epsilon_Q$ , pure plasma state at  $T_c$  is reached at  $v_i$ .

Fig. 7 Average  $p_t$  as a function of normalized multiplicity  $\left(\frac{dN}{dy}\right)$  for U-U ( $R_A = 6.8\text{ fm}$ ), Fe-Fe ( $R_A = 4.2\text{ fm}$ ), and O-O ( $R_A = 2.8\text{ fm}$ ) collisions.

Fig. 8 Comparison of average  $p_t$  at initial time, i.e. the earliest time with thermal equilibrium, with no transverse flow and at the decoupling time when the flow is fully developed but the matter has cooled down to dec.

Fig. 9 Comparison of the shapes of the  $p_t$  spectra of (massless) pions, kaons, and nucleons in U-U collisions for  $\frac{1}{A} \left(\frac{dN}{dy}\right) = 13.5$ . For

ormalization, see the text.

Fig. 10 Average  $p_t$  for pions, kaons, and nucleons as function of normalized multiplicity  $\frac{1}{A} \left( \frac{dN}{dy} \right)$  in U-U collisions ( $R_A = 6.8$  fm).

Fig. 11 Comparison of the results for average  $p_t$  from different decoupling algorithms.

Fig. 12 Comparison of the hydrodynamic results for average  $p_t$  with the data of JACEE<sup>(19)</sup> group.

Table I

Average transverse momentum of pions for different values of critical temperature. Multiplicity is fixed to  $\frac{1}{A} \frac{dN}{dy} = 27.5$  and  $r_0 = 6.8$  Fm.

$T_c$	160	200	300	400
$\langle p_t \rangle$	592	632	817	980






Fig. 1







Fig. 2



(a)



(b)

Fig. 3





Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8






Fig. 9




Fig.10




Fig.11




Fig. 12