

New Limits on $K_{L,S} \rightarrow \pi^0 e^+ e^-$

L. K. Gibbons, V. Papadimitriou, J. R. Patterson, Y.W. Wah, B. Winstein,
R. Winston, M. Woods ^(a), and H. Yamamoto
The Enrico Fermi Institute and the Department of Physics,
The University of Chicago, Chicago, IL 60637

E. C. Swallow
Department of Physics, Elmhurst College, Elmhurst, IL 60126 and
The Enrico Fermi Institute, The University of Chicago, Chicago, IL 60637

G. J. Bock, R. Coleman, Y. B. Hsiung, K. Stanfield, R. Stefanski,
and T. Yamanaka
Fermi National Accelerator Laboratory, Batavia, IL 60510

G. Blair ^(b), G. D. Gollin, M. Karlsson, and J. K. Okamitsu
Department of Physics, Princeton University

P. Debu, B. Peyaud, R. Turlay, and B. Vallage
Department de Physique des Particules Elementaires, Centre d'Etudes
Nucleaires de Saclay, F-91191 Gif-sur-Yvette Cedex, France

ABSTRACT

Data taken in a Fermilab experiment designed to measure the CP violation parameter ϵ'/ϵ from a study of $K \rightarrow 2\pi$ decays were used to look for the as yet unseen decay modes $K_{L,S} \rightarrow \pi^0 e^+ e^-$. The detector was optimized for the detection of kaon decays with four electromagnetic showers in the final state. The results (90 % confidence) are branching ratios $< 4.2 \times 10^{-8}$ and $< 4.5 \times 10^{-5}$ for $K_L \rightarrow \pi^0 e^+ e^-$ and $K_S \rightarrow \pi^0 e^+ e^-$ respectively .

The $\pi^0 e^+ e^-$ decay of the long-lived neutral kaon (K_L) is an attractive avenue for the observation of CP violation in a decay amplitude, particularly should detailed studies of the 2π decays of the neutral kaon (ϵ'/ϵ) prove inconclusive. The CP violating amplitude is expected¹ to be comparable to or larger than the conserving one. The leading CP conserving amplitude proceeds through two photon exchange while the violating one may proceed via one photon exchange. Within the framework of the Standard Model where CP violation comes from the phase δ in the Kobayashi-Maskawa matrix², $K_L \rightarrow \pi^0 e^+ e^-$ may have a sizable $\Delta S=1$ CP violating effect. Theoretical estimates¹ of the branching ratio are in the 10^{-11} range while the current experimental limit³ is $< 2.3 \times 10^{-6}$ at 90% confidence.

Fermilab experiment E-731 which was performed in the Meson Center beam line at the Tevatron, had as its primary goal the determination of the $K \rightarrow 2\pi$ CP violation parameter ϵ'/ϵ . The present search is based upon the analysis of a special data set in which $K_{L,S} \rightarrow \pi^0 \pi^0$ and $K_{L,S} \rightarrow \pi^+ \pi^-$ were recorded simultaneously.

Two neutral K_L beams ($1/2 \times 1/2$ mrad²) were created at 4.8 mrad by 800 GeV protons striking a Be target. A regenerator which moved alternately between the beams every proton spill was used to provide K_S . The detector was employed in an earlier test run and it is shown schematically in Fig.1; it has been described elsewhere^{4,5} in detail. Charged particles were measured and momentum analyzed with a 2000-wire drift-chamber spectrometer which consisted of eight x planes and eight y planes with 0.635 cm maximum drift distance. These planes had a position resolution of about 110 μm and were 98 % efficient. Energies and positions of photons and electrons were measured with an 804 block lead-glass array stacked circularly. Each block measured 5.82 cm (H) x 5.82 cm (W) x 60 cm (L), giving a depth of 20 radiation lengths. There were two holes (11.6 cm x 11.6 cm) separated

vertically by 11.6 cm at the center of the array for the beams to pass. A common pulsed light source illuminated every block once every second to provide short term gain tracking. The pulse heights were digitized with effectively 15 bit ADCs using a 150 ns gate.

Several improvements have been made to the detector since the previous data taking period. The most important one was the instrumentation of each of the lead-glass phototube outputs with a 60 MHz flash ADC. These were the front-end electronics for a two dimensional cluster finding trigger processor, and they also served to suppress out-of-time photons. A cluster was defined as a 'neighbor-connected' island of lead-glass blocks each with more than 1 GeV. The trigger processor contributed less than 2 % dead-time and a FASTBUS-based data acquisition system was implemented to increase the data taking capability as well.

There were two triggers relevant to the data set used in this search. The first ('four-cluster') required exactly four clusters, 30 GeV or more energy deposited in the lead-glass, and no hit in the trigger plane (see Fig. 1). Hence this trigger recorded $\pi^0 e^+ e^-$ candidates in the downstream decay region as well as $\pi^0 \pi^0$ candidates from both upstream and downstream decay regions. The second trigger ('two-track') required one or more hits at the trigger plane and two tracks in the spectrometer and hence was sensitive to $\pi^0 e^+ e^-$ decays from the upstream decay region; however, this trigger was prescaled by a factor of 8.

Because the trigger processor used signals from the lead-glass directly, the relative gains of all lead-glass blocks were monitored and adjusted to within 5 % over the entire data taking period. Calibration data with $e^+ e^-$ pairs produced in a thin upstream foil were taken periodically to provide high statistics calibration for the lead-glass; the resolution was $\sigma/E \simeq 1.5\% + 5\%/\sqrt{E}$. Chamber alignment data were also

recorded daily.

The momenta of the e^+ and e^- and the decay vertex of $K_L \rightarrow \pi^0 e^+ e^-$ candidates were determined by the drift chamber spectrometer. The e^+ and e^- were identified by matching the tracks with the clusters, and requiring $0.85 < E/P < 1.15$, where E is the cluster energy and P is the momentum. Figure 2(a) shows the E/P distribution for electrons from calibration data. From a study of $K_L \rightarrow \pi^+ \pi^- \pi^0$ decays, the π^0 mass resolution was determined to be about $4 \text{ MeV}/c^2$ (see Fig.2(b)). The $\gamma\gamma$ mass was required to be within $10 \text{ MeV}/c^2$ of the nominal π^0 value. By then constraining the $\gamma\gamma$ mass to the nominal value, the reconstructed kaon mass ($M_{\pi ee}$) would have a resolution of about $4.5 \text{ MeV}/c^2$. The square of the transverse momentum (P_t^2) of the $\pi^0 e^+ e^-$ system with respect to the line connecting the decay vertex and the production target had a resolution of about $50 \text{ MeV}^2/c^2$. The candidates are displayed in a two dimensional $M_{\pi ee}$ vs. P_t^2 plot as shown in Fig. 3(b). A candidate is defined to have $P_t^2 < 200 \text{ MeV}^2 / c^2$ and $489 < m_K < 507 \text{ MeV}/c^2$; these cuts would include about 95% of the signal. No candidate is found in the signal region. Figure 3(a) shows the equivalent region for $K_L \rightarrow \pi^+ \pi^- \pi^0$ decays. Given the timing and energy resolution of the detector, possible backgrounds such as radiative K_{e3} with an accidental photon or $K \rightarrow 2\pi^0$ with Dalitz decays are insignificant.

Figure 4 shows the relative acceptance for Monte Carlo generated $K_L \rightarrow \pi^0 e^+ e^-$ decays as a function of the $e^+ e^-$ effective mass. For the 'four-cluster' trigger, the acceptance is 9.5 % for a fiducial downstream decay volume of 22.2 meters and for the 'two-track' trigger, the acceptance is about 10 % for an upstream decay volume of 14 meters for kaon energy between 30 and 150 GeV, assuming a uniform three-body phase space distribution.

The upper limit $B.R.(K_L \rightarrow \pi^0 e^+ e^-) < 4.7 \times 10^{-8}$ (90 % confidence) is obtained

by normalizing to a sample of 58.8×10^3 $K_L \rightarrow 2\pi^0$ decays observed simultaneously in the 'four-cluster' trigger. The normalization data do not require track reconstruction, however the relative branching ratios of $\pi^+\pi^-$ to $\pi^0\pi^0$ from both K_L and K_S were also determined from the same data set and they agree with the published values to within 5 %. Using 71.3×10^3 $K_L \rightarrow \pi^+\pi^-$ and 2.24×10^5 $K_S \rightarrow \pi^+\pi^-$ decays from the 'two-track' trigger as normalization, the 90 % confidence limits from that trigger are $< 4.1 \times 10^{-7}$ and $< 4.5 \times 10^{-5}$ respectively.

Combining the above, the results are $B.R.(K_L \rightarrow \pi^0 e^+ e^-) < 4.2 \times 10^{-8}$ and $B.R.(K_S \rightarrow \pi^0 e^+ e^-) < 4.5 \times 10^{-5}$ (90 % confidence). This is the first limit of any significance for the K_S decay. The K_L limit is an improvement of more than a factor of 50 over the previous limit³; while still far from the level predicted by the Standard Model, it serves to constrain the parameters of light scalar particles coupling to e^+e^- .

We are happy to acknowledge important contributions by H.Sanders, J. Ting and E. Weatherhead from The University of Chicago; G. Grazer from Princeton University; J.C. Brisson, R. Daudin and P. Jarry from Saclay; and D. Gielow and T. Kowalczyk from Elmhurst College. The support of the Fermilab staff is gratefully acknowledged. This work was supported in part by the National Science Foundation, the Department of Energy, and the French Atomic Energy Commission.

REFERENCES

- (a) Present address : SLAC, Stanford, CA 94305.
- (b) Present address : Department of Nuclear Physics, Oxford University, Oxford, United Kingdom.
1. J. F. Donoghue, B. R. Holstein, and G. Valencia, Phys. Rev. D35, 2769 (1987) and references within; L. M. Sehgal, Phys. Rev. D38, 808 (1988); G. Ecker, A. Pich, and E. deRafael, Nucl. Phys. B303, 665 (1988).
 2. M. Kobayashi and T. Maskawa, Prog. Theor. Phys. 49, 652 (1973).
 3. A. S. Carroll et. al., Phys. Rev. Lett. 46, 525 (1980).
 4. M. Woods et. al., Phys. Rev. Lett. 60, 1695 (1988).
 5. For a complete description of the detector see P. Jarry, Ph.D. thesis, Universite de Paris-Sud, 1987 (unpublished); M. Woods, Ph.D. thesis, University of Chicago, 1988 (unpublished).

FIGURE CAPTIONS

Figure 1. Detector schematic, elevation view.

Figure 2. (a) Distribution of E/P in the lead-glass from the electron calibration data.

The resolution is about 4% rms; (b) distribution of the π^0 mass reconstructed from $K_L \rightarrow \pi^+\pi^-\pi^0$ decays.

Figure 3. Reconstructed kaon mass vs. the square of the transverse momentum for (a)

$K_L \rightarrow \pi^+\pi^-\pi^0$ and (b) $K_L \rightarrow \pi^0e^+e^-$. There are 24 events above the top of the plot in (b). The events in the plots were selected with a π^0 mass cut of 2.5σ and the boxes represent the signal region.

Figure 4. The relative acceptance vs the e^+e^- invariant mass of the $K_L \rightarrow \pi^0e^+e^-$ for the 'four-cluster' trigger.







