

Fermi National Accelerator Laboratory

FERMILAB-Pub-92/236-E

**Inclusive J/ψ , ψ (2S) and b-quark Production in
 $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV**

**F. Abe et al
The CDF Collaboration**

Fermi National Accelerator Laboratory, Batavia, Illinois 60510

August 1992

Submitted to *Physical Review Letters*



Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Inclusive J/ψ , $\psi(2S)$ and b -quark Production in $\bar{p}p$ Collisions at $\sqrt{s} = 1.8$ TeV

F. Abe,⁽¹¹⁾ D. Amidei,⁽¹⁴⁾ C. Anway-Wiese,⁽⁸⁾ G. Apollinari,⁽²⁰⁾ M. Atac,⁽⁶⁾ P. Auchincloss,⁽¹⁹⁾
P. Assi,⁽¹⁵⁾ A. R. Baden,⁽⁸⁾ N. Bacchetta,⁽¹⁵⁾ W. Badgett,⁽¹⁴⁾ M. W. Bailey,⁽¹⁸⁾ A. Bamberger,^(9,*)
P. de Barbaro,⁽¹⁹⁾ A. Barbaro-Galtieri,⁽¹²⁾ V. E. Barnes,⁽¹⁶⁾ B. A. Barnett,⁽¹⁰⁾ G. Bauer,⁽¹³⁾
T. Baumann,⁽⁸⁾ F. Bedeschi,⁽¹⁷⁾ S. Behrends,⁽²⁾ S. Belforte,⁽¹⁷⁾ G. Bellettini,⁽¹⁷⁾ J. Bellinger,⁽²⁵⁾
D. Benjamin,⁽²⁴⁾ J. Benloch,^(6,*) J. Bensinger,⁽²⁾ A. Beretvas,⁽⁶⁾ J. P. Berge,⁽⁶⁾ S. Bertolucci,⁽⁷⁾
K. Biery,^(16,*) S. Bhadra,⁽⁹⁾ M. Binkley,⁽⁶⁾ D. Bisello,⁽¹⁵⁾ R. Blair,⁽¹⁾ C. Blocker,⁽²⁾ A. Bodek,⁽¹⁹⁾
V. Bolognesi,⁽¹⁷⁾ A. W. Booth,⁽⁶⁾ C. Boswell,⁽¹⁰⁾ G. Brandenburg,⁽⁸⁾ D. Brown,⁽⁸⁾
E. Buckley-Geer,⁽²¹⁾ H. S. Budd,⁽¹⁹⁾ G. Busetto,⁽¹⁵⁾ A. Byon-Wagner,⁽⁶⁾ K. L. Byrum,⁽²⁵⁾
C. Campagnari,⁽⁶⁾ M. Campbell,⁽¹⁴⁾ A. Caner,⁽⁹⁾ R. Carey,⁽⁸⁾ W. Carithers,⁽¹²⁾ D. Carlsmith,⁽²⁵⁾
J. T. Carroll,⁽⁶⁾ R. Cashmore,^(9,*) A. Castro,⁽¹⁵⁾ F. Cervelli,⁽¹⁷⁾ K. Chadwick,⁽⁹⁾ J. Chapman,⁽¹⁴⁾
G. Chiarelli,⁽⁷⁾ W. Chinowsky,⁽¹²⁾ S. Cihangir,⁽⁶⁾ A. G. Clark,⁽⁶⁾ M. Cobal,⁽¹⁷⁾ D. Connor,⁽¹⁶⁾
M. Contreras,⁽⁴⁾ J. Cooper,⁽⁶⁾ M. Cordelli,⁽⁷⁾ D. Crane,⁽⁶⁾ J. D. Cunningham,⁽²⁾ C. Day,⁽⁶⁾
F. DeJongh,⁽⁶⁾ S. Dell'Agnello,⁽¹⁷⁾ M. Dell'Orso,⁽¹⁷⁾ L. Demortier,⁽²⁾ B. Denby,⁽⁶⁾
P. F. Derwent,⁽¹⁴⁾ T. Devlin,⁽²¹⁾ D. DiBitonto,⁽²²⁾ M. Dickson,⁽²⁰⁾ R. B. Drucker,⁽¹²⁾
K. Einsweiler,⁽¹²⁾ J. E. Elias,⁽⁶⁾ R. Ely,⁽¹²⁾ S. Eno,⁽⁴⁾ S. Errede,⁽⁹⁾ A. Etchegoyen,^(9,*)
B. Farhat,⁽¹²⁾ G. Feldman,⁽⁶⁾ B. Flaughner,⁽⁶⁾ G. W. Foster,⁽⁶⁾ M. Franklin,⁽⁸⁾ J. Freeman,⁽⁶⁾
D. Frei,^(6,*) H. Frisch,⁽⁴⁾ T. Fuess,⁽⁶⁾ Y. Fukui,⁽¹¹⁾ A. F. Garfinkel,⁽¹⁶⁾ A. Gauthier,⁽⁹⁾ S. Geer,⁽⁶⁾
D. W. Gerdes,⁽⁴⁾ P. Giannetti,⁽¹⁷⁾ N. Giokaris,⁽²⁰⁾ P. Giromini,⁽⁷⁾ L. Gladney,⁽¹⁶⁾ M. Gold,⁽¹²⁾
K. Goulianos,⁽²⁰⁾ H. Grassmann,⁽¹⁵⁾ G. M. Grieco,⁽¹⁷⁾ R. Grindley,^(16,*) C. Grosso-Pilcher,⁽⁴⁾
C. Haber,⁽¹²⁾ S. R. Hahn,⁽⁶⁾ R. Handler,⁽²⁵⁾ K. Hara,⁽²³⁾ B. Harral,⁽¹⁶⁾ R. M. Harris,⁽⁶⁾
S. A. Hauger,⁽⁸⁾ J. Hauser,⁽⁸⁾ C. Hawk,⁽²¹⁾ T. Hessing,⁽²²⁾ R. Hollebeck,⁽¹⁶⁾ L. Holloway,⁽⁹⁾
S. Hong,⁽¹⁴⁾ P. Hu,⁽²¹⁾ B. Hubbard,⁽¹²⁾ B. T. Huffman,⁽¹⁶⁾ R. Hughes,⁽¹⁶⁾ P. Hurst,⁽⁶⁾ J. Huth,⁽⁶⁾
J. Hylen,⁽⁶⁾ M. Incagli,⁽¹⁷⁾ T. Ino,⁽²³⁾ H. Iso,⁽²³⁾ H. Jensen,⁽⁶⁾ C. P. Jessop,⁽⁶⁾ R. P. Johnson,⁽⁶⁾
U. Joshi,⁽⁶⁾ R. W. Kadel,⁽¹²⁾ T. Kamon,⁽²²⁾ S. Kanda,⁽²³⁾ D. A. Kardelis,⁽²³⁾ I. Karliner,⁽⁹⁾
E. Kearns,⁽⁸⁾ L. Keeble,⁽²²⁾ R. Kephart,⁽⁶⁾ P. Kesten,⁽²⁾ R. M. Keup,⁽⁹⁾ H. Keutelian,⁽⁶⁾ D. Kim,⁽⁹⁾
S. B. Kim,⁽¹⁴⁾ S. H. Kim,⁽²³⁾ Y. K. Kim,⁽¹²⁾ L. Kirsch,⁽²⁾ K. Kondo,⁽²³⁾ J. Konigsberg,⁽⁶⁾
K. Kordas,^(16,*) E. Kovacs,⁽⁶⁾ M. Krasberg,⁽¹⁴⁾ S. E. Kuhlmann,⁽¹⁾ E. Kuns,⁽²¹⁾
A. T. Laasanen,⁽¹⁹⁾ S. Lammel,⁽⁸⁾ J. I. Lamoureux,⁽²⁵⁾ S. Leone,⁽¹⁷⁾ J. D. Lewis,⁽⁶⁾ W. Li,⁽¹⁾
P. Limon,⁽⁶⁾ M. Lindgren,⁽⁸⁾ T. M. Liss,⁽⁹⁾ N. Lockyer,⁽¹⁶⁾ M. Loretì,⁽¹⁸⁾ E. H. Low,⁽¹⁶⁾
D. Lucchesi,⁽¹⁷⁾ C. B. Luchini,⁽⁹⁾ P. Lukens,⁽⁶⁾ P. Maas,⁽²³⁾ K. Maeshima,⁽⁶⁾ M. Mangano,⁽¹⁷⁾
J. P. Marriner,⁽⁶⁾ M. Mariotti,⁽¹⁷⁾ R. Markeloff,⁽²⁵⁾ L. A. Markosky,⁽²⁵⁾ R. Mattingly,⁽²⁾
P. McIntyre,⁽²²⁾ A. Mensione,⁽¹⁷⁾ E. Meschi,⁽¹⁷⁾ T. Meyer,⁽²²⁾ S. Mikamo,⁽¹¹⁾ M. Miller,⁽⁴⁾
T. Mimashi,⁽²³⁾ S. Miscetti,⁽⁷⁾ M. Mishina,⁽¹¹⁾ S. Miyashita,⁽²³⁾ Y. Morita,⁽²³⁾ S. Moulding,⁽²⁾ J.
Mueller,⁽²¹⁾ A. Mukherjee,⁽⁶⁾ T. Muller,⁽³⁾ L. F. Nakae,⁽²⁾ I. Nakano,⁽²³⁾ C. Nelson,⁽⁶⁾
D. Neuberger,⁽³⁾ C. Newman-Holmes,⁽⁶⁾ J. S. T. Ng,⁽⁸⁾ M. Ninomiya,⁽²³⁾ L. Nodulman,⁽¹⁾
S. Ogawa,⁽²³⁾ R. Paoletti,⁽¹⁷⁾ V. Papadimitriou,⁽⁶⁾ A. Para,⁽⁶⁾ E. Pare,⁽⁸⁾ S. Park,⁽⁶⁾ J. Patrick,⁽⁶⁾
G. Pauletta,⁽¹⁷⁾ L. Pescara,⁽¹⁸⁾ T. J. Phillips,⁽⁸⁾ F. Ptohos,⁽⁶⁾ R. Plunkett,⁽⁶⁾ L. Pondrom,⁽²⁵⁾
J. Proudfoot,⁽¹⁾ G. Punsì,⁽¹⁷⁾ D. Quarrie,⁽⁶⁾ K. Ragan,^(16,*) G. Redlinger,⁽⁴⁾ J. Rhoades,⁽²⁵⁾
M. Roach,⁽²⁴⁾ F. Rimondi,^(6,*) L. Ristori,⁽¹⁷⁾ W. J. Robertson,⁽⁶⁾ T. Rodrigo,⁽⁶⁾ T. Rohaly,⁽¹⁶⁾
A. Roodman,⁽⁴⁾ W. K. Sakumoto,⁽¹⁹⁾ A. Sansoni,⁽⁷⁾ R. D. Sard,⁽⁹⁾ A. Savoy-Navarro,⁽⁹⁾
V. Scarpine,⁽⁹⁾ P. Schlabach,⁽⁸⁾ E. E. Schmidt,⁽⁹⁾ O. Schneider,⁽¹²⁾ M. H. Schub,⁽¹⁶⁾
R. Schwitters,⁽⁸⁾ A. Scribano,⁽¹⁷⁾ S. Segler,⁽⁶⁾ Y. Seiya,⁽²³⁾ G. Sganos,^(16,*) M. Shapiro,⁽¹²⁾
N. M. Shaw,⁽¹⁶⁾ M. Sheaff,⁽²⁵⁾ M. Shochet,⁽⁴⁾ J. Siegrist,⁽¹²⁾ P. Sinervo,^(16,*) J. Skarha,⁽¹⁰⁾
K. Sliwa,⁽²⁴⁾ A. Spies,⁽¹⁰⁾ D. A. Smith,⁽¹⁷⁾ F. D. Snider,⁽¹⁰⁾ L. Song,⁽⁶⁾ T. Song,⁽¹⁴⁾ M. Spahn,⁽¹²⁾
P. Sphicas,⁽¹²⁾ R. St. Denis,⁽⁸⁾ L. Stanco,^(6,*) A. Stefanini,⁽¹⁷⁾ G. Sullivan,⁽⁴⁾ K. Sumorok,⁽¹³⁾
R. L. Swartz, Jr.,⁽⁹⁾ M. Takano,⁽²³⁾ K. Takikawa,⁽²³⁾ S. Tarem,⁽²⁾ F. Tartarelli,⁽¹⁷⁾ S. Tether,⁽¹³⁾
D. Theriot,⁽⁶⁾ M. Timko,⁽²⁴⁾ P. Tipton,⁽¹⁹⁾ S. Tkaczyk,⁽⁶⁾ A. Tollestrup,⁽⁶⁾ J. Tonnison,⁽¹⁶⁾
W. Trischuk,⁽⁸⁾ Y. Tsay,⁽⁴⁾ J. Tseng,⁽¹⁰⁾ N. Turini,⁽¹⁷⁾ F. Ukegawa,⁽²³⁾ D. Underwood,⁽¹⁾ S. Vejckik,

III,⁽¹⁰⁾ R. G. Wagner,⁽¹⁾ R. L. Wagner,⁽⁶⁾ N. Wainer,⁽⁶⁾ R. C. Walker,⁽¹⁹⁾ J. Walsh,⁽¹⁶⁾
 G. Watts,⁽¹⁹⁾ T. Watts,⁽²¹⁾ R. Webb,⁽²²⁾ C. Wendt,⁽²⁵⁾ H. Wenzel,⁽¹⁷⁾ W. C. Wester, III,⁽¹²⁾
 T. Westhusing,⁽⁹⁾ S. N. White,⁽²⁰⁾ A. B. Wicklund,⁽¹⁾ E. Wicklund,⁽⁶⁾ H. H. Williams,⁽¹⁶⁾
 B. L. Winer,⁽¹⁹⁾ J. Wolinski,⁽²²⁾ D. Wu,⁽¹⁴⁾ J. Wyss,⁽¹⁵⁾ A. Yagil,⁽⁶⁾ K. Yasuoka,⁽²³⁾ Y. Ye,^(16,a)
 G. P. Yeh,⁽⁶⁾ J. Yoh,⁽⁶⁾ M. Yokoyama,⁽²³⁾ J. C. Yun,⁽⁶⁾ A. Zanetti,⁽¹⁷⁾ F. Zetti,⁽¹⁷⁾ S. Zhang,⁽¹⁴⁾
 S. Zucchelli,^(6,a)

The CDF Collaboration

- (1) Argonne National Laboratory, Argonne, Illinois 60439
 (2) Brandeis University, Waltham, Massachusetts 02254
 (3) University of California at Los Angeles, Los Angeles, California 90024
 (4) University of Chicago, Chicago, Illinois 60637
 (5) Duke University, Durham, North Carolina 27706
 (6) Fermi National Accelerator Laboratory, Batavia, Illinois 60510
 (7) Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, Frascati, Italy
 (8) Harvard University, Cambridge, Massachusetts 02138
 (9) University of Illinois, Urbana, Illinois 61801
 (10) The Johns Hopkins University, Baltimore, Maryland 21218
 (11) National Laboratory for High Energy Physics (KEK), Japan
 (12) Lawrence Berkeley Laboratory, Berkeley, California 94720
 (13) Massachusetts Institute of Technology, Cambridge, Massachusetts 02139
 (14) University of Michigan, Ann Arbor, Michigan 48109
 (15) Università di Padova, Istituto Nazionale di Fisica Nucleare, Sezione di Padova, I-35131 Padova, Italy
 (16) University of Pennsylvania, Philadelphia, Pennsylvania 19104
 (17) Istituto Nazionale di Fisica Nucleare, University and Scuola Normale Superiore of Pisa, I-56100 Pisa, Italy
 (18) Purdue University, West Lafayette, Indiana 47907
 (19) University of Rochester, Rochester, New York 14627
 (20) Rockefeller University, New York, New York 10021
 (21) Rutgers University, Piscataway, New Jersey 08854
 (22) Texas A&M University, College Station, Texas 77843
 (23) University of Tsukuba, Tsukuba, Ibaraki 305, Japan
 (24) Tufts University, Medford, Massachusetts 02155
 (25) University of Wisconsin, Madison, Wisconsin 53706

Abstract

Inclusive J/ψ and $\psi(2S)$ production has been studied in $\bar{p}p$ collisions at $\sqrt{s} = 1.8$ TeV using 2.6 ± 0.2 pb⁻¹ of data taken with the Collider Detector at Fermilab. The products of production cross section times branching fraction were measured as functions of P_T for $J/\psi \rightarrow \mu^+\mu^-$ and $\psi(2S) \rightarrow \mu^+\mu^-$. In the kinematic range $P_T > 6$ GeV/c and $|\eta| \leq 0.5$ we get $\sigma(\bar{p}p \rightarrow J/\psi X) \times B(J/\psi \rightarrow \mu^+\mu^-) = 6.88 \pm 0.23(stat) \begin{smallmatrix} +0.93 \\ -1.08 \end{smallmatrix} (syst)$ nb, and $\sigma(\bar{p}p \rightarrow \psi(2S)X) \times B(\psi(2S) \rightarrow \mu^+\mu^-) = 0.232 \pm 0.051(stat) \begin{smallmatrix} +0.029 \\ -0.032 \end{smallmatrix} (syst)$ nb. From these values we calculate the inclusive b -quark production cross section.

PACS numbers: 13.85.Ni, 14.40.Jz, 14.80.Dq

We report a study of the reactions $\bar{p}p \rightarrow J/\psi(\psi(2S)) X \rightarrow \mu^+\mu^-X$ at $\sqrt{s} = 1.8$ TeV. This study yields the P_T dependence of the products of production cross section times branching fraction over the kinematic range $P_T > 6$ GeV/c and $|\eta| \leq 0.5$. The products of integrated cross section times branching fraction are also presented. These measurements are important for the investigation of charmonium production mechanisms in $\bar{p}p$ collisions [1]. Both J/ψ 's and $\psi(2S)$'s can be produced either by direct gluon fusion or in the decay of B mesons. J/ψ 's also occur as daughters of directly-produced χ 's. The resulting differences in P_T distributions for J/ψ and $\psi(2S)$ can help to understand these mechanisms. We also emphasize the importance of the J/ψ and $\psi(2S)$ signals to study the production of b quarks at low P_T [2, 3]. We have compared J/ψ and $\psi(2S)$ production with a Monte Carlo model for b -quark production and B -meson decay in order to obtain the inclusive cross section for production of b -quarks. This technique was first used by UA1 [2].

This is the first measurement of J/ψ and $\psi(2S)$ cross sections at Tevatron energies. The data were taken with the Collider Detector at Fermilab (CDF) which has been described in detail elsewhere[4]. Here we give a brief description of the components relevant to this analysis. The central tracking chamber (CTC) is in a 1.4116-T axial magnetic field and has a resolution of $\delta P_T/P_T = \sqrt{(0.0011P_T)^2 + (0.0066)^2}$ for beam constrained tracks, where P_T is the momentum transverse to the beam direction. The central muon chambers, at a radius of 3.5 m from the beam axis, provide muon identification in the region of pseudorapidity $|\eta^\mu| < 0.61$, where $\eta^\mu = -\ln(\tan \frac{\theta}{2})$ and θ is the polar angle with respect to the beam axis. The measurements reported here are based on a data sample with a total integrated luminosity of 2.6 ± 0.2 pb $^{-1}$, collected with a multi-level central dimuon trigger. The Level 1 trigger required a track segment in the central muon chambers with a threshold set at a nominal transverse momentum, $P_T^\mu > 3$ GeV/c. The Level 2 trigger required two

muon chamber track segments that each satisfied the Level 1 trigger, that were separated from each other by at least one full muon module (15° wide in ϕ) and that each matched, within a muon module, to a track in the CTC. From events passing this trigger, a sample of opposite sign dimuons was selected with the following cuts: $P_T^\mu > 3.0$ GeV/ c for each muon; less than a 3σ difference in position between each muon chamber track and its associated, extrapolated CTC track, where σ is the calculated uncertainty due to multiple scattering, energy loss, and measurement uncertainties; a common vertex along the beam axis for the two muons; $|\eta| \leq 0.5$ and $6.0 < P_T < 14.0$ GeV/ c for the muon pair. The resulting mass distributions after all cuts are shown in Fig. 1. The J/ψ and $\psi(2S)$ peaks were each fit to a Gaussian line shape plus a linear background. The number of J/ψ candidates above background within a $\pm 2.5\sigma$ mass signal region, $3.05 < m_{\mu^+\mu^-} < 3.15$ GeV/ c^2 , is 889 ± 30 and the resulting J/ψ mass is (3.0965 ± 0.0007) GeV/ c^2 with $\sigma = (18.5 \pm 0.6)$ MeV/ c^2 . The number of $\psi(2S)$ candidates above background within a $\pm 2.5\sigma$ mass signal region, $3.63 < m_{\mu^+\mu^-} < 3.73$ GeV/ c^2 , is 35 ± 8 and the $\psi(2S)$ mass is (3.683 ± 0.005) GeV/ c^2 with $\sigma = (20 \pm 4)$ MeV/ c^2 .

The geometric and kinematic acceptances for $J/\psi \rightarrow \mu^+\mu^-$ and $\psi(2S) \rightarrow \mu^+\mu^-$ were evaluated by a dimuon event generator that produces J/ψ 's and $\psi(2S)$'s with flat P_T , η , and ϕ distributions and with a tunable polarization distribution. The generated events were processed with the full detector simulation and the same reconstruction used for the data. The acceptance is P_T -dependent, increasing from 3% at 6 GeV/ c to plateaux of 15.5% for J/ψ and 19% for $\psi(2S)$. To verify that this is independent of the kinematic distribution of generated events, the acceptance calculation was repeated for J/ψ using the ISAJET generator[5]. This yielded good agreement between the two acceptance calculations.

Efficiency corrections required for the J/ψ and $\psi(2S)$ cross section calculations are the

following: the Level 1 trigger efficiency for each muon is an increasing function of P_T , rising from $68 \pm 5\%$ at $3 \text{ GeV}/c$ to $91 \pm 3\%$ for $P_T > 6 \text{ GeV}/c$. The CTC track finding efficiency in the Level 2 trigger is $99 \pm 1\%$ for each muon. The total trigger efficiency for each dimuon event is taken to be the product of the Level 1 and Level 2 efficiencies for each muon. The offline CTC track reconstruction efficiency is $97 \pm 2\%$. The muon reconstruction efficiency is $98 \pm 1\%$. The matching cut on the difference between the muon chamber track and the extrapolated CTC track is $97 \pm 1\%$ efficient. The cuts on the J/ψ and $\psi(2S)$ mass window are $97 \pm 2\%$ efficient.

Polarized J/ψ 's ($\psi(2S)$'s) from B meson decay would produce muons with an angular distribution of $\frac{d\sigma}{d\Omega} \sim 1 + \alpha \times \cos^2\theta$, where α is a parameter that depends on the $J/\psi(\psi(2S))$ polarization and θ is the angle in the J/ψ ($\psi(2S)$) rest frame between one of the emerging muons and the J/ψ ($\psi(2S)$) direction in its parent's rest frame. The uncertainty introduced in the quarkonium cross sections by the unknown polarization was evaluated by calculating the acceptance with the two extreme values, $\alpha = +1$ and $\alpha = -1$ with the conservative assumption that all J/ψ 's ($\psi(2S)$'s) originate from B mesons. Two sets of J/ψ 's ($\psi(2S)$'s), one of each polarization, were generated in the B meson rest frame with a momentum spectrum given by ARGUS [6] for J/ψ and by CLEO [7] for $\psi(2S)$. The B mesons were produced from b quarks with the transverse momentum spectrum given by Nason, Dawson, & Ellis (NDE) [8] and with the Peterson fragmentation function [9], using $\epsilon_P = 0.006 \pm 0.002$. The central values of the $J/\psi(\psi(2S))$ production cross sections are given for zero polarization,

and the systematic uncertainties due to unknown polarizations are

$+6.5 (4.4)\%$.
$-11.2(7.3)\%$.

The other major systematic uncertainty in both the J/ψ and $\psi(2S)$ production cross sections is due to the trigger efficiency and was estimated to be $\pm 9\%$.

The J/ψ and $\psi(2S)$ differential cross sections as functions of P_T are displayed in Fig. 2. The vertical error bars are from statistical fluctuations in the number of counts (background fluctuations included) and the P_T -dependent systematic uncertainties added in quadrature. Theoretical predictions for the two types of processes expected to dominate J/ψ and $\psi(2S)$ production are also plotted.

The solid curve in Fig. 2a (2b) is a next-to-leading-order calculation by NDE of the production of b -quarks [8] leading to B -mesons and subsequent decay to J/ψ ($\psi(2S)$) as discussed above. We refer to this overall calculation as B-production model (BPM). Uncertainties in this prediction arise because the b mass is comparable to the P_T of this experiment and in a range where neglected higher order terms may be significant. Further, the strongest leading-order processes for b production are gluon fusion at low x where the structure functions are not well known. For these reasons, “the prediction of bottom production at collider energies is subject to considerable uncertainty.”[10]

The dashed curve in Fig. 2a (2b) corresponds to J/ψ 's ($\psi(2S)$'s) from direct charmonium production [1, 11]. We refer to this overall calculation as the charmonium production model (CPM). The direct $\psi(2S)$ production was estimated [11] with the same parameters and structure functions as used for J/ψ direct charmonium production and is found to be very small, ~ 25 times smaller in magnitude than the data.

The sum of these two contributions (BPM and CPM) to J/ψ production is plotted in Fig. 2a. We also fit the theory to the data by summing the two theoretical contributions with independent normalization factors. This assumes that there are no other significant contributions to J/ψ production. With no normalization constraints a good fit ($\chi^2/d.f. = 9.1/14$) was obtained with the CPM factor at ~ 4.4 ($\sim 69\%$ of J/ψ production) and the BPM factor at ~ 2.2 ($\sim 31\%$). The fit suppresses the BPM contribution because of the

difference in slope between the BPM curve and the data. However, a previous CDF study [3] of $B^\pm \rightarrow J/\psi K^\pm$ showed that the BPM calculation underestimates the b -quark cross section by a factor of 5.5 ± 2.8 which indicates that $(75_{-40}^{+25})\%$ of our J/ψ 's come from B decays. This datum was added to the fit with a Poisson probability distribution yielding $\chi^2/d.f. = 14.9/15$ and an BPM factor of 2.9 ± 0.5 corresponding to $\sim 42\%$ J/ψ 's from B -production. The 90% C.L. upper limit on the BPM contribution is $\sim 60\%$. If future measurements exceed this value, then one must conclude that not only the normalization of BPM, but also the P_T -dependence of at least one of the models is wrong.

Figure 2b shows a similar set of curves for $\psi(2S)$ production. While the normalization and slope of the BPM contribution seem to differ from those of the data in the same way as those for J/ψ , the statistical significance here does not allow any conclusion.

The products of the inclusive production cross section times branching fraction in the kinematic range $P_T > 6$ GeV/ c and $|\eta| \leq 0.5$ are:

$$\sigma(\bar{p}p \rightarrow J/\psi X) \times B(J/\psi \rightarrow \mu^+ \mu^-) = 6.88 \pm 0.23(stat) \begin{matrix} +0.93 \\ -1.08 \end{matrix} (syst) \text{ nb and}$$

$$\sigma(\bar{p}p \rightarrow \psi(2S)X) \times B(\psi(2S) \rightarrow \mu^+ \mu^-) = 0.232 \pm 0.051(stat) \begin{matrix} +0.029 \\ -0.032 \end{matrix} (syst) \text{ nb,}$$

where an extrapolation of the cross sections for values of $P_T > 14$ GeV/ c was carried out.

We assign a 2% uncertainty in the above values due to this extrapolation.

In order to determine the b -quark cross section, we use the measurement of the J/ψ ($\psi(2S)$) inclusive production cross sections, the ratio of J/ψ ($\psi(2S)$) to b -quark cross sections as determined by a Monte Carlo technique [12, 8, 9, 6, 7], the combined branching ratios $B(B \rightarrow J/\psi(\psi(2S)) X) \times B(J/\psi(\psi(2S)) \rightarrow \mu^+ \mu^-)$, and the fraction f_B of $J/\psi(\psi(2S))$'s

from B meson decays:

$$\sigma_{exp}^b(P_T^b > P_T^{min}, |y^b| < 1) = \frac{B(J/\psi(\psi(2S)) \rightarrow \mu^+\mu^-) \times \sigma_{exp}^c(P_T^c > 6\text{GeV}/c, |\eta^c| < 0.5) \times R \times f_B}{2 \times B(B \rightarrow J/\psi(\psi(2S))X) \times B(J/\psi(\psi(2S)) \rightarrow \mu^+\mu^-)}, \quad (1)$$

where

$$R = \frac{\sigma_{BPM}^b(P_T^b > P_T^{min}, |y^b| < 1.0)}{\sigma_{BPM}^c(P_T^c > 6\text{GeV}/c, |\eta^c| < 0.5)}$$

and the index ‘‘c’’ stands for ‘‘ $J/\psi(\psi(2S))$ ’’. The rapidity of the b -quark is $y^b = \frac{1}{2} \ln\left(\frac{E^b + P_{||}^b}{E^b - P_{||}^b}\right)$. The Monte Carlo program is used to determine P_T^{min} which is chosen such that approximately 90% of the produced J/ψ ($\psi(2S)$) have $P_T^b > P_T^{min}$; we have set $P_T^{min} = 8.5$ GeV/c for this analysis. The branching ratios are taken as unity for the calculation of σ_{BPM}^c in the Monte Carlo. The combined branching ratios we use are: $B(B \rightarrow J/\psi X) \times B(J/\psi \rightarrow \mu^+\mu^-) = (7.7 \pm 1.3) \times 10^{-4}$ [13, 14] and $B(B \rightarrow \psi(2S)X) \times B(\psi(2S) \rightarrow \mu^+\mu^-) = (3.6 \pm 1.4) \times 10^{-5}$ [13, 15]. The ‘2’ in the denominator of Eq. 1 arises since $J/\psi(\psi(2S))$ ’s can either be produced by b or \bar{b} quarks and we are interested only in the b -quark cross section. We assign an 8% systematic error to the b -quark cross section due to the uncertainty in the shape of the ARGUS(CLEO) momentum spectra for J/ψ ’s ($\psi(2S)$ ’s). Although we are sensitive to b -quarks with $|y^b| < 0.6$, we are quoting the b -quark cross section in the rapidity range $|y^b| < 1.0$ for comparison with theory[8]. We assign a 4% systematic error to the b -quark cross section due to the uncertainty in the rapidity spectrum in the range $0.6 < |y^b| < 1.0$. We also assign a 5% systematic error to the b -quark cross section due to variations in the shape of the b -quark P_T spectrum and a 6% systematic error due to the uncertainty in ϵ_P . Assuming the fraction f_B to be unity, we get:

$$\sigma^b(P_T^b > 8.5\text{GeV}/c, |y^b| < 1) = 18.9^{+4.7}_{-5.0} \mu\text{b} \text{ using } J/\psi \text{ 's.}$$

and

$$\sigma^b(P_T^b > 8.5\text{GeV}/c, |y^b| < 1) = 10.5^{+5.0}_{-5.1} \mu\text{b} \text{ using } \psi(2S) \text{ 's.}$$

The fraction is believed to be close to one for $\psi(2S)$'s [16, 1, 11] but not for the J/ψ . The b -quark cross section we get using $\psi(2S)$'s is approximately 1.5 standard deviations higher than the theoretical calculation [8], in reasonable agreement with Ref. [3].

We thank the Fermilab staff and the technical and support staffs of the participating institutions for their vital contributions. We also acknowledge Dr. N. Glover for useful discussions. This work was supported by the Department of Energy, the National Science Foundation, the Istituto Nazionale di Fisica Nucleare, the Ministry of Science, Culture and Education of Japan, and the A. P. Sloan Foundation.

References



^a Visitor

- [1] E.W.N.Glover, A.D.Martin, W.J.Stirling, *Z. Phys.* **C38**, 473 (1988).
- [2] C. Albajar et al., *Physics Letters* **B213**, 405 (1988); C. Albajar et al., *Physics Letters* **B256**, 121 (1991).
- [3] F. Abe et al., *Phys. Rev. Lett.* **B68**, 3403 (1992).
- [4] F. Abe et al., *Nucl. Inst. Meth.* **A271**, 387 (1988); F. Bedeschi et al., *Nucl. Inst. Meth.* **A268**, 50 (1988); G. Ascoli et al., *Nucl. Inst. Meth.* **A268**, 33 (1988).
- [5] F. Paige and S. D. Protopopescu, ISAJET Monte Carlo version 6.36, BNL Report No. BNL38034, 1986 (unpublished).

- [6] H. Schroder, personal communication.
- [7] W. Chen, Ph.D Thesis, Purdue University, May, 1990.
- [8] P. Nason, S. Dawson, and R.K. Ellis, Nucl. Phys. **B303**, 607 (1988), Nucl. Phys. **B327**, 49 (1989), Nucl. Phys. **B335**, 260 (1990). The present curves are for their choice of parameters: $m_B = 4.75 \text{ GeV}/c^2$, $\Lambda_4 = 270 \text{ MeV}$ and the DFLM structure functions.
- [9] C. Peterson et al., Phys. Rev. **D27**, 105 (1983).
- [10] R. K. Ellis, Proceedings of the 17th SLAC Summer Institute of Particle Physics, SLAC Report 361, Stanford Linear Accelerator Center, Stanford, CA, July, 1989.
- [11] E.W.N.Glover, personal communication. The present calculation was performed using the method of Ref. [1] and the theoretical uncertainty is estimated to be $\pm 50\%$ or more. It used $Q^2 = (M^2 + P_T^2)/4$ and the D_0 structure functions of A. D. Martin, W. J. Stirling and R. G. Roberts, RAL-92-021, DTP/92/16, April, 1992.
- [12] N. Ellis and A. Kernan, Physics Reports **195**, 23 (1990).
- [13] H. Albrecht et al., Physics Letters **B199**, 451 (1987).
- [14] M. S. Alam et al., Phys. Rev. **D34**, 3279 (1986).
- [15] Particle Data Group, M. Aguillar-Benitez et al., Rev. Mod. Phys. **56**, S1 (1984).
- [16] S. D. Ellis, M. B. Einhorn, C. Quigg, Phys. Rev. Letters **36**, 1263 (1976), C. E. Carlson, R. Suaya, Phys. Review **D15**, 1416 (1977).

Figure Captions

- 1) The mass distribution of $\mu^+\mu^-$ for (a) J/ψ and (b) $\psi(2S)$. The histogram corresponds to the data and the solid curve is a fit to a Gaussian plus a linear background.
- 2) The product $B \times \left(\frac{d\sigma}{dP_T}\right)$ vs. P_T for (a) $J/\psi \rightarrow \mu^+\mu^-$ and (b) $\psi(2S) \rightarrow \mu^+\mu^-$. The circles correspond to the data. The solid curve corresponds to $J/\psi(\psi(2S))$'s produced from B meson decays by using Ref.[8, 9, 6, 7]. The dashed curve corresponds to $J/\psi(\psi(2S))$'s from direct charmonium production from the model of Ref.[1]. The dot-dashed curve is their sum. Uncertainties in the theoretical curves are discussed in the text.



$M(\mu^+, \mu^-)$ (GeV/c^2)

