



**Fermi National Accelerator Laboratory**

**FERMILAB-Conf-93/162-E  
E789 and P865**

## **E789 and P865: High-Rate Fixed-Target Studies of Charm and Beauty**

**Daniel M. Kaplan**

*Physics Department, Northern Illinois University,  
DeKalb, Illinois 60115*

**for the E789 and P865 Collaborations**

E789 Collaboration: J. Boissevain<sup>1</sup>, C.N. Brown<sup>2</sup>, G. Brown<sup>3</sup>, T.A. Carey<sup>1</sup>, Y.C. Chen<sup>4,5</sup>, R.L. Childers<sup>6</sup>, W.E. Cooper<sup>2</sup>, C.W. Darden<sup>6</sup>, G. Gidal<sup>7</sup>, H.D. Glass<sup>2</sup>, K.N. Gounder<sup>2</sup>, P.M. Ho<sup>7</sup>, L.D. Isenhower<sup>3</sup>, D.M. Jansen<sup>1</sup>, R. Jeppesen<sup>1</sup>, D.M. Kaplan<sup>8</sup>, J.S. Kapustinsky<sup>1</sup>, G.C. Kiang<sup>4</sup>, M.S. Kowitt<sup>10</sup>, D.W. Lane<sup>1</sup>, L.M. Lederman<sup>9</sup>, M.J. Leitch<sup>1</sup>, J.W. Lillberg<sup>1</sup>, W. Luebke<sup>8</sup>, K.B. Luk<sup>7,10</sup>, V.M. Martin<sup>8</sup>, P.L. McGaughey<sup>1</sup>, C.S. Mishra<sup>2</sup>, J.M. Moss<sup>1</sup>, R.S. Preston<sup>8</sup>, D. Pripstein<sup>10</sup>, J. Sa<sup>8</sup>, M.E. Sadler<sup>3</sup>, R. Schnathorst<sup>3</sup>, M.H. Schub<sup>9</sup>, R. Schwindt<sup>3</sup>, D. Snodgrass<sup>6</sup>, W.E. Sondheim<sup>1</sup>, V. Tanikella<sup>8</sup>, P.K. Teng<sup>4</sup>, J.R. Wilson<sup>6</sup>

P865 Collaboration: M. Atac<sup>2,11</sup>, C.N. Brown<sup>2</sup>, R.C. Chaney<sup>1,2</sup>, G. Charpak<sup>13</sup>, R.L. Childers<sup>6</sup>, R. Chipaux<sup>14</sup>, D. Chrisman<sup>11</sup>, D. Cline<sup>11</sup>, W.E. Cooper<sup>2</sup>, H.B. Crawley<sup>15</sup>, C.W. Darden<sup>6</sup>, J. Derré<sup>14</sup>, L. Diaz-Cruz<sup>16</sup>, E.J. Fenyves<sup>12</sup>, A. Firestone<sup>15</sup>, J.R. Friedrich<sup>12</sup>, Y. Giomataris<sup>17</sup>, H. Hammack<sup>12</sup>, L.D. Isenhower<sup>3</sup>, C. Joseph<sup>17</sup>, D.M. Kaplan<sup>8</sup>, C. Kochowski<sup>14</sup>, J.W. Lamsa<sup>15</sup>, L.M. Lederman<sup>18</sup>, Y. Lemoigne<sup>14</sup>, S. Loucatos<sup>14</sup>, W.T. Meyer<sup>15</sup>, C. Morel<sup>17</sup>, J. Orgeron<sup>12</sup>, J. Park<sup>11</sup>, M. Perez<sup>16</sup>, J.P. Perroud<sup>17</sup>, Ph. Rebougeard<sup>14</sup>, J. Rhoades<sup>11</sup>, E.I. Rosenberg<sup>15</sup>, M.E. Sadler<sup>3</sup>, J. Toscano<sup>16</sup>, M.T. Tran<sup>17</sup>, J.R. Wilson<sup>6</sup>

<sup>1</sup>Los Alamos National Laboratory    <sup>2</sup>Fermilab    <sup>3</sup>Abilene Christian University    <sup>4</sup>Academia Sinica, Taiwan  
<sup>5</sup>National Cheng Kung University, Taiwan    <sup>6</sup>University of South Carolina    <sup>7</sup>Lawrence Berkeley Laboratory  
<sup>8</sup>Northern Illinois University    <sup>9</sup>University of Chicago    <sup>10</sup>University of California at Berkeley  
<sup>11</sup>University of California at Los Angeles    <sup>12</sup>University of Texas at Dallas    <sup>13</sup>CERN    <sup>14</sup>CE Saclay  
<sup>15</sup>Iowa State University    <sup>16</sup>CINVESTAV-IPN, Mexico    <sup>17</sup>Université de Lausanne    <sup>18</sup>Illinois Institute of Technology

June 1993

Presented at *Heavy Quarks at Fixed Target (HQ93)*,  
Laboratori Nazionali di Frascati, Frascati, Italy, May 31-June 2, 1993



## **Disclaimer**

*This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.*

# E789 and P865: High-Rate Fixed-Target Studies of Charm and Beauty

Daniel M. Kaplan

*Physics Department, Northern Illinois University,  
DeKalb, Illinois 60115, USA*

for the E789\* and P865† collaborations

June 1993

\*E789 Collaboration: J. Boissevain,<sup>1</sup> C. N. Brown,<sup>2</sup> G. Brown,<sup>3</sup> T. A. Carey,<sup>1</sup> Y. C. Chen,<sup>4,5</sup> R. L. Childers,<sup>6</sup> W. E. Cooper,<sup>2</sup> C. W. Darden,<sup>6</sup> G. Gidal,<sup>7</sup> H. D. Glass,<sup>2</sup> K. N. Gounder,<sup>2</sup> P. M. Ho,<sup>7</sup> L. D. Isenhower,<sup>3</sup> D. M. Jansen,<sup>1</sup> R. Jeppesen,<sup>1</sup> D. M. Kaplan,<sup>8</sup> J. S. Kapustinsky,<sup>1</sup> G. C. Kiang,<sup>4</sup> M. S. Kowitt,<sup>10</sup> D. W. Lane,<sup>1</sup> L. M. Lederman,<sup>9</sup> M. J. Leitch,<sup>1</sup> J. W. Lillberg,<sup>1</sup> W. Luebke,<sup>8</sup> K. B. Luk,<sup>7,10</sup> V. M. Martin,<sup>8</sup> P. L. McGaughey,<sup>1</sup> C. S. Mishra,<sup>2</sup> J. M. Moss,<sup>1</sup> R. S. Preston,<sup>8</sup> D. Pripstein,<sup>10</sup> J. Sa,<sup>8</sup> M. E. Sadler,<sup>3</sup> R. Schnathorst,<sup>3</sup> M. H. Schub,<sup>9</sup> R. Schwindt,<sup>3</sup> D. Snodgrass,<sup>6</sup> W. E. Sondhein,<sup>1</sup> V. Tanikella,<sup>8</sup> P. K. Teng,<sup>4</sup> J. R. Wilson<sup>6</sup>

†P865 Collaboration: M. Atac,<sup>2,11</sup> C. N. Brown,<sup>2</sup> R. C. Chaney,<sup>12</sup> G. Charpak,<sup>13</sup> R. L. Childers,<sup>6</sup> R. Chipaux,<sup>14</sup> D. Chrisman,<sup>11</sup> D. Cline,<sup>11</sup> W. E. Cooper,<sup>2</sup> H. B. Crawley,<sup>15</sup> C. W. Darden,<sup>6</sup> J. Derré,<sup>14</sup> L. Diaz-Crus,<sup>16</sup> E. J. Fenyves,<sup>12</sup> A. Firestone,<sup>15</sup> J. R. Friedrich,<sup>12</sup> Y. Giomataris,<sup>17</sup> H. Hammack,<sup>12</sup> L. D. Isenhower,<sup>3</sup> C. Joseph,<sup>17</sup> D. M. Kaplan,<sup>8</sup> C. Kochowski,<sup>14</sup> J. W. Lamsa,<sup>15</sup> L. M. Lederman,<sup>18</sup> Y. Lemoigne,<sup>14</sup> S. Loucatos,<sup>14</sup> W. T. Meyer,<sup>15</sup> C. Morel,<sup>17</sup> J. Orgeron,<sup>12</sup> J. Park,<sup>11</sup> M. Perez,<sup>16</sup> J.-P. Perroud,<sup>17</sup> Ph. Rebourgeard,<sup>14</sup> J. Rhoades,<sup>11</sup> E. I. Rosenberg,<sup>15</sup> M. E. Sadler,<sup>3</sup> J. Toscano,<sup>16</sup> M. T. Tran,<sup>17</sup> J. R. Wilson<sup>6</sup>

<sup>1</sup> Los Alamos National Laboratory    <sup>2</sup> Fermilab    <sup>3</sup> Abilene Christian University

<sup>4</sup> Academia Sinica, Taiwan    <sup>5</sup> National Cheng Kung University, Taiwan

<sup>6</sup> University of South Carolina    <sup>7</sup> Lawrence Berkeley Laboratory

<sup>8</sup> Northern Illinois University    <sup>9</sup> University of Chicago

<sup>10</sup> University of California at Berkeley    <sup>11</sup> University of California at Los Angeles

<sup>12</sup> University of Texas at Dallas    <sup>13</sup> CERN    <sup>14</sup> CE Saclay    <sup>15</sup> Iowa State University

<sup>16</sup> CINVESTAV-IPN, Mexico    <sup>17</sup> Université de Lausanne

<sup>18</sup> Illinois Institute of Technology

Presented at Heavy Quarks at Fixed Target (HQ93), Laboratori Nazionali di Frascati, Frascati, Italy, May 31 – June 2, 1993

# **E789 AND P865: HIGH-RATE FIXED-TARGET STUDIES OF CHARM AND BEAUTY**

**Daniel M. Kaplan**

*Northern Illinois University, DeKalb, Illinois, USA*

for the E789 and P865 collaborations

## **Abstract**

Experiment 789 at Fermilab used the high-rate E605/E772 spectrometer to study low-multiplicity charm and beauty decays. Preliminary results on charm and beauty production are presented based on analysis of  $\approx 100\%$  of the charm data and  $\approx 50\%$  of the beauty data. A new experiment is proposed to improve charm and beauty sensitivity by several orders of magnitude.

## **1. Description of E789**

Experiment 789 studied charm and beauty production in 800-GeV  $pN$  collisions using the high-rate E605/E772 spectrometer (Figure 1) in Fermilab's Meson-East laboratory, upgraded with a 16-plane silicon-microstrip array and a vertex trigger processor [1]. Beam of intensity  $\approx (1 - 6) \times 10^{10}$  protons per 20s spill was incident on thin wire targets 0.1 – 0.2 mm high and 0.8 – 3 mm thick. For the charm data the SM12 analyzing magnet was operated at 900 or 1000 amperes, and Au and Be targets were used to study the  $A$ -dependence of  $D$  production. For the beauty data a 3-mm-thick Au target was used and the SM12 current was 1500 amperes, optimizing acceptance for the  $J/\psi$  while simultaneously accepting direct two-prong  $B^0$  decays. (Insufficient running time was available for a dedicated  $B^0 \rightarrow$  dihadron run.) In the 1991 run,  $4 \times 10^8$  ( $9 \times 10^8$ ) events were recorded on 240 (770) 8-mm tapes during the charm (beauty) running periods. Typical interaction rates were  $\approx 2$  MHz in the charm mode and  $\approx 50$  MHz in the beauty mode.

The silicon detectors were arrayed from 40 – 115 cm downstream of the target in two arms covering the angular ranges  $\pm(20 - 60)$  mr above and below the beam axis. The detectors, type "B" from Micron Semiconductor, were of  $5 \times 5$  cm<sup>2</sup> area, 300  $\mu$ m thickness, and 50  $\mu$ m pitch. In each arm, planes measuring the bend ( $y$ - $z$ ) view alternated with planes at  $\pm 5^\circ$  stereo angles. Signals from 8,544 strips were individually read out via Fermilab 128-channel amplifier cards [2] and LBL discriminators [3] synchronized to the accelerator RF. The discriminated signals were transmitted through  $\approx 400$  ns of multiconductor cable to coincidence registers.




Figure 1: E789 apparatus (plan view).

The vertex processor [1], which found tracks in the bend-view drift chambers and silicon detectors and selected events with candidate decay vertices downstream of the target, was used in the charm running and provided a rejection factor of  $\approx 10$  against non-charm events with charm efficiency of  $\approx (50-60)\%$  (caused primarily by silicon-plane inefficiencies) for decays outside the target. The thin target localized the primary interaction vertex, simplifying the design of the processor, which needed only to determine the position of the decay vertex.

## 2. Preliminary Results from E789

Figure 2 shows dihadron invariant mass spectra from the charm data sample for various lifetime cuts. Cherenkov  $\pi/K$  particle identification has not been used in this analysis, as the RICH was not optimized for the low momenta characteristic of charm decays. Rather, each plot contains two entries per event, one for the  $\pi^+ K^-$  and the other for the  $\pi^- K^+$  particle assignments. The correct particle assignments give a sharp peak ( $\sigma \approx 6$  MeV), corresponding to either the  $D^0$  or  $\bar{D}^0$ . The wrong particle assignments give a much broader peak ( $\sigma \approx 50$  MeV). A limit on possible  $D - \bar{D}$  production-rate asymmetry can be derived from the numbers of  $D^0$  and  $\bar{D}^0$  observed. At the 10% level, no significant asymmetry is observed (or expected) at the  $x_F \approx 0.05$  characterizing these data. Our preliminary  $D$  cross section is consistent with previous measurements reported by E653 [4] and E743 [5]. The systematics of the  $A$ -dependence and normalization are still under study.

Figure 3a shows the invariant mass spectrum from a preliminary analysis of  $\approx 50\%$  of the beauty dimuon data sample. Figure 3b shows the spectrum of events in which both tracks have impact parameters at the target center greater than  $150 \mu\text{m}$  and the pair vertex satisfies  $0.7 < z_{\text{vertex}} < 5$  cm, indicating a decay downstream of the target. In the  $J/\psi$  mass region, 22 events satisfy these requirements. To




Figure 2: Dihadron invariant mass spectra from the charm data sample for three lifetime cuts: a)  $\tau/\sigma_\tau > 3.9$ , b)  $\tau/\sigma_\tau > 6$ , c)  $\tau/\sigma_\tau > 7.9$ .

estimate background due to tails of the target distribution (arising from possible vertex tracking errors or large Coulomb scatters in the silicon planes), we also study events with reconstructed vertices upstream of the target (Figure 3c). No upstream events are observed near the  $J/\psi$  region, supporting the likely beauty origin of the downstream  $J/\psi$  events. There is also a significant excess of continuum events in the downstream sample, attributed to semileptonic decays of  $b\bar{b}$  pairs.




Figure 3: Dimuon invariant mass spectra from the beauty data sample: a) all events, b)  $0.7 < z_{\text{vertex}} < 5$  cm, c)  $-5 < z_{\text{vertex}} < -0.7$  cm.

### 3. Proposal 865

The sensitivity of E789 is limited primarily by its small acceptance ( $\lesssim 10\%$  per prong). We have proposed a new apparatus [6] (Figure 4) which retains the high interaction-rate capability of E789 while achieving acceptance of  $\approx 70\%$  per prong. This allows studies of charm at the level of  $10^{11}$  produced and  $10^7 - 10^8$  fully reconstructed events, and of beauty at the level of  $10^8$  produced and  $10^4$  fully reconstructed events, giving access to such physics as  $b \rightarrow s\gamma$  transitions,  $B_s$  mixing, beauty semileptonic form factors, doubly-Cabibbo-suppressed charm decays, semileptonic  $D_s$  and charmed-baryon decays, and  $D^0$  mixing and  $CP$  violation.




Figure 4: P865 apparatus (elevation).

A significant sensitivity limit is posed by radiation damage in the silicon detectors. To configure detectors which can survive at the desired sensitivity, we choose suitable maximum and (in one view) minimum angles for the instrumented aperture, arranging the silicon detectors along the beam axis with a small gap through which pass the uninteracted beam and secondaries below the minimum angle (Figure 5). Thus the rate is spread approximately equally over several silicon planes, with large-angle secondaries measured close to the target and small-angle secondaries farther downstream. Along the beam axis the spacing of detectors increases approximately geometrically, making the lever arm for vertex reconstruction independent of production angle. The instrumented angular range is  $|\theta_x| \leq 200 \text{ mr}$ ,  $8 \leq \theta_y \leq 150 \text{ mr}$ , corresponding to the rapidity range  $|y| \lesssim 1.5$ . To maximize the rate capability of the spectrometer, the tracking is performed entirely with silicon and scintillating-fiber planes.

The scintillating fibers are read out using the solid-state “visible-light photon counters” (VLPCs) under development by a collaboration among Fermilab, UCLA,




Figure 5: Detail of P865 vertex region.

and Rockwell International Science Center [7]. VLPCs are also key to the proposed Cherenkov-based optical impact-parameter trigger [8] and fast ring-imaging Cherenkov counter [9]. VLPCs are highly suitable for these applications due to their high quantum efficiency, low noise, and high speed. Quantum efficiency as high as 85% for green light has been achieved [10], and 30-MHz rate capability has been demonstrated, with single-electron noise rates of several kHz [11].

In addition to the Level-1 optical trigger, other triggering strategies appear promising. These include calorimetric  $E_t$  and high- $p_t$ -lepton requirements and a trigger processor to reconstruct vertices and invariant masses. At each trigger level, just enough rejection should be provided to meet the input bandwidth of the succeeding level, so as to trigger on charm and beauty as efficiently as possible without restricting attention only to modes with final-state leptons. High-bandwidth data-recording technology and inexpensive workstations now or soon-to-be available should make feasible a recorded sample of  $\sim 10^{11}$  events, key to the desired charm sensitivity.

P865 seeks to study the physics described above within the next few years, then upgrade the apparatus for a sensitive study of  $B$   $CP$  violation by decade's end. This might be accomplished by increasing the interaction rate a factor  $\approx 10$ , employing (for example) radiation-hardened silicon pixel detectors to cope with the increased event complexity and radiation dose. Alternatively, the beauty production cross section might be increased by moving to a higher- $\sqrt{s}$  interaction region. A promising idea under study [12] is to arrange collisions between a high-energy proton beam in the Tevatron, SPS, or HERA proton ring and a lower-energy beam, for example by storing protons in the HERA electron ring or constructing a small, inexpensive storage ring adjacent to the Tevatron or SPS. At  $\sqrt{s} \approx (150 - 200)$  GeV, this scheme can provide a factor  $\approx 100$  increase in beauty cross section while retaining sufficient Lorentz boost to ease triggering, particle identification, and coverage of a large acceptance.

## 4. Summary

Experiment 789 explores the feasibility of studying beauty and charm physics in a high-rate fixed-target environment. Over 3000 neutral  $D \rightarrow \pi K$  events have been observed from Au and Be targets. A preliminary analysis of  $\approx 50\%$  of the dimuon beauty data yields 22  $J/\psi$  events with decay vertices more than 7 mm downstream of the target; these are candidate  $b \rightarrow J/\psi + X$  decays. Analysis is underway to determine  $\sigma(b\bar{b})$  at 800 GeV. Analysis of the dihadron beauty data and  $J/\psi \rightarrow e^+e^-$  data is also in progress. A new experiment, P865, has been proposed as the next step in charm and beauty sensitivity, with the ultimate goal of studying  $CP$  violation in beauty decays.

## References

- [1] C. Lee *et al.*, IEEE Trans. Nucl. Sci. **38** (1989) 461.
- [2] D. Christian *et al.*, IEEE Trans. Nucl. Sci. **36** (1989) 507.
- [3] B. T. Turko *et al.*, IEEE Trans. Nucl. Sci. **39** (1992) 758 .
- [4] K. Kodama *et al.*, Phys. Lett. B **263** (1991) 573 .
- [5] R. Ammar *et al.*, Phys. Rev. Lett. **61** (1988) 2185 .
- [6] L. D. Isenhower *et al.*, "P865: Revised Letter of Intent for a High-Sensitivity Study of Charm and Beauty Decays," D. M. Kaplan, spokesperson, April 2, 1993.
- [7] M. D. Petroff and M. Atac, IEEE Trans. Nucl. Sci. **36** (1989) 163;  
M. Atac *et al.*, Nucl. Instr. & Meth. **A314** (1992) 56;  
M. Atac *et al.*, Nucl. Instr. & Meth. **A320** (1992) 155.
- [8] G. Charpak, L. M. Lederman, and Y. Giomataris, Nucl. Instr. & Meth. **A306** (1991) 439 ;  
D. M. Kaplan *et al.*, Nucl. Instr. & Meth. **A330** (1993) 33;  
G. Charpak *et al.*, "Experimental Study of an Impact-Parameter Optical Discriminator", CERN-PPE/93-14, to appear in Nucl. Instr. & Meth. (1993).
- [9] D. M. Kaplan *et al.*, "A Fast Ring-Imaging Cherenkov Counter for a Fixed-Target Heavy-Quark Experiment," FERMILAB-Conf-93/148, to appear in *Proceedings of the First Workshop on RICH Detectors*, Bari, Italy, June 1993.
- [10] M. G. Stapelbroek, M. D. Petroff, and R. Bharat, "Visible Light Photon Counters for Astronomy," to appear in *Proceedings of an ESA Symposium on Photon Detectors for Space Instrumentation*, ESA/ESTEC, Noordwijk, The Netherlands, 10-12 November 1992.
- [11] M. D. Petroff and W. G. Stapelbroek, IEEE Trans. Nucl. Sci. **NS-36** (1989) 158.
- [12] Y. Giomataris, "Perspectives on an Asymmetric Proton Collider with Small Crossing Angle for B Physics," this Workshop.