



CONSTRAINTS ON THE STRENGTH OF A PRIMORDIAL MAGNETIC FIELD FROM BIG BANG NUCLEOSYNTHESIS *

B. Cheng¹, D.N. Schramm^{2,3} and J.W. Truran²

¹Department of Astronomy, University of Illinois at Urban-Champaign
1002 W. Green Street, Urbana, IL 61801 USA

²Department of Astronomy & Astrophysics, The University of Chicago
5640 S. Ellis Ave., Chicago IL 60637 USA

³NASA/Fermilab Astrophysics Center
Box 500, Batavia, IL 60510 USA

Abstract

The effects of magnetic fields on Big Bang Nucleosynthesis (BBN) have been calculated, and the impact on the abundances of the light elements have been investigated numerically. An upper limit on the strength of primordial magnetic fields compatible with observations of light element abundances has been thus obtained. In the framework of standard BBN theory, the maximum strength of the primordial magnetic fields, on scales greater than 10^4 cm but smaller than the event horizon at the BBN epoch (~ 1 min., $\sim 2 \times 10^{12}$ cm), is $\leq 10^{11}$ Gauss. This limit is shown to allow magnetic fields at the time of recombination no stronger than ~ 0.1 gauss on scales $\geq 10^{11}$ cm. Our results also strongly indicate that, at the BBN epoch, and for field strengths $B \leq 10^{13}$ gauss, the effects of magnetic fields on the primordial abundances of light elements are dominated by effects from reaction rates in the presence of primeval magnetic fields rather than by magnetic density effects on the expansion rate.

* submitted to *Physical Review D*



1. Introduction

It is recognized that primordial nucleosynthesis provides a unique quantitative window on the early universe^[1,2]. Since the synthesis of the light elements is determined by events occurring in the epoch from $\sim 1s$ to $\sim 1000s$ in the history of the universe, when the temperatures varied from $\sim 10^{10}K$ ($\geq 1MeV$) to $\sim 10^9K$ ($\leq 0.1MeV$), the observed abundances constitute a probe of the universe at epochs far earlier than those directly probed by the cosmic microwave background radiation (CMBR) ($t \sim 10^5yr$; $T \sim 10^4K$ ($\sim 1eV$)). Thus, through a detailed comparison of the predicted abundances with the observational data, proposed cosmological models can be tested and their controlling parameters can be constrained. For example, big bang nucleosynthesis (BBN) was found to constrain the number of families of light neutrinos^[3] prior to accurate accelerator measurements.

In this paper, we reexplore how the strength of certain primordial magnetic fields can be constrained by BBN. If magnetic fields of sufficient strength existed in the early universe, particularly at or just before the epoch of primordial nucleosynthesis, they could have had direct influences on both the expansion rate of the universe and the nuclear reaction rates.^[4,5] These influences could, of course, affect the abundances of the light elements produced in this environment. In addition, if the scale of the primeval magnetic field were greater than the event horizon, the geometry of the universe would also be affected and an anisotropic universe might result. An analysis of nucleosynthesis in anisotropic Euclidean Universes, in which the dependence of the primordial abundances of 4He , 3He , D on the isotropy parameters was specified more precisely, has been presented by Thorne^[4] and by Hawking and Tayler.^[6] If a significant degree of anisotropy had persisted up to times ≥ 20000 years, the primordial 4He abundance would have been reduced to a few per cent. On the contrary, if the anisotropy is important only during the early stages of the expansion, the 4He abundance is about 30% while there is no (negligible) D or 3He production, and one might hope to eventually reach agreement with the observational values by refining this model. This had been done by Juskiewicz et al,^[7] who studied the influence of the anisotropic momentum distribution of neutrinos neglected by Thorne. The resultant limit on the magnetic field at the BBN time, set by the condition of small anisotropy for $t > 1s$ is about $B < 4.1 \times 10^{12}Gauss$.^[8]

On the other hand, if the primeval magnetic field were sufficiently spread over distances small compared with the event horizon at that epoch, the geometry of the universe would not be affected^[9] and it would still be described by a Robertson-Walker metric. For this situation, it has been qualitatively pointed out by a number of authors^[9,10] that, in the presence of a very intense magnetic field ($B \geq 10^{13}G$), the neutron would decay more rapidly than in the field-free case, this could obviously affect light element synthesis in a dramatic way.

However, in previous studies, only the effects of a very strong primordial magnetic field ($B > 10^{13}$ G) on abundances ${}^4\text{He}$, D , and ${}^3\text{He}$, have been studied, not vice-versa, and no critical limit on the primeval magnetic field was explicitly derived. The questions we address in this paper are: (1) What is the limit on the primordial magnetic field? (2) How does the magnetic field influence the emerging abundances of other light elements ($A > 9$), such as lithium, boron, etc.? We find that there are still constraints to be explored on the strength and coherence scale of primordial magnetic fields, using observational abundances and BBN.

2. The Direct Effects of the Primeval Field on BBN

In the early expansion of the universe, the existence of a large scale primordial magnetic field may have both direct and indirect effects on BBN. In this regard, the two most sensitive and competing effects are: (1) the effects on reaction rates and (2) the effects on the expansion rate. These two effects will further alter the resulting abundances of the elements.

For simplicity, we assume that our universe is fully filled by randomly oriented and distributed thin-wall magnetic domains (or bubbles)^[11]. The size of each domain is large enough so that the field inside the domain can be seen as a uniform field, but it is still small compared with the event horizon. Thus, the magnetic field will have similar effects on the motion of particles in each domain. Here, we will neglect the boundary effects since the wall is assumed to be thin.

As to the effects on reaction rates, we have recently derived the reaction rates as a function of a uniform magnetic field \vec{B} in the presence of an arbitrary degeneracy and polarization^[12]. As an application, if we assume that the magnetic field is nearly uniform in each domain, we can use our derived results (Ref.[12]) as a first order approximation for our purpose here. However, we would like to point out that, if the magnetic field is not uniform through the whole universe but varies with spacial variables, or if the scale of the magnetic field (or magnetic bubbles) is much smaller than the horizon scale and the magnetic domains are disconnected from each other, the nuclear reaction rates will become inhomogeneous; i.e., the reaction rates will differ from region to region even though the geometry of the universe is still not affected. This would require that we introduce reaction rate fluctuations into the standard big bang code (similar to the introduction of the density fluctuations associated with the first order QCD phase transition^[13]), and perform multizone calculations. Such exploration is beyond the scope of the present paper and will be addressed in future work.

Now let us explore the effects of the magnetic fields on the expansion rate of the universe. According to our assumptions, the globally chaotic (but locally orderly) magnetic

fields will have no effect on the geometry of the universe. The geometry of the universe is still described by a Robertson-Walker metric. For this metric, the work-energy equations can be expressed as

$$\frac{d}{dt}(\rho R^3) + \frac{p}{c^2} \frac{d}{dt}(R^3) = 0, \quad (2.1)$$

where $R(t)$ is the distance measure, ρ is the total mass-energy density, and p is the total pressure.

In general, we consider that the universe consists of three types of matter during the epoch of interest. These are: (1) the strongly and electromagnetically interacting particles (e.g., nucleons, electrons, photons, etc.), which can be described as a perfect fluid; (2) the weakly interacting particles, which nevertheless affect the n-p ratio (e.g, electron neutrinos, etc.); and (3) the effectively noninteracting particles (e.g., ν_μ and so on) which only contribute to the energy density but do not enter into specific reactions. The total mass-energy density ρ and pressure p can be expressed as

$$\rho = \rho_\gamma + \rho_e + \rho_\nu + \rho_b + \rho_B, \quad p = p_\gamma + p_e + p_\nu + p_b + p_B, \quad (2.2)$$

where

$$\begin{aligned} \rho_e &= \rho_{e^-} + \rho_{e^+}, & p_e &= p_{e^-} + p_{e^+}, \\ \rho_\nu &= \rho_{\nu_{e^-}} + \rho_{\nu_{e^+}} + \rho_{\nu_\mu} + \rho_{\nu_\tau} + \rho_{\bar{\nu}_e} + \rho_{\bar{\nu}_\mu} + \rho_{\bar{\nu}_\tau}, \\ p_\nu &= p_{\nu_{e^-}} + p_{\nu_{e^+}} + p_{\nu_\mu} + p_{\nu_\tau} + p_{\bar{\nu}_e} + p_{\bar{\nu}_\mu} + p_{\bar{\nu}_\tau}, \end{aligned}$$

and the subscripts γ , e , ν_e , ν_μ , ν_τ , b , and B stand, respectively, for photons, electrons, e-neutrinos, μ -neutrinos, τ -neutrinos, baryons, and magnetic field. Expressions for these thermodynamic quantities are given below, for the case of nondegenerate neutrinos^[14]

$$\begin{aligned} \rho_\gamma &= 8.42 T_9^4 \text{ gm cm}^{-3}, & p_\gamma &= \frac{1}{3} \rho_\gamma c^2, & \rho_\nu &= 6 \rho_{\nu_i} = \frac{21}{8} \rho_\gamma \left(\frac{T_\nu}{T}\right)^4, \\ p_\nu &= \frac{1}{3} \rho_\nu c^2, & \rho_e &= \frac{7}{4} \rho_\gamma (T_9 \gg 6), & p_e &= \frac{1}{3} \rho_e c^2 (T_9 \gg 6), \\ \rho_b &= 7 \times 10^{-6} T_9^3 \text{ gm cm}^{-3}, & p_b &= n_b k T \sum_i Y_i, & Y_i &= \frac{X_i}{A_i}, & \rho_B &= \frac{B^2}{8\pi} \end{aligned}, \quad (2.3)$$

where $T_9 = T/(10^9 K)$, ρ_{ν_i} ($\nu_i = \nu_e, \bar{\nu}_e, \nu_\mu, \bar{\nu}_\mu, \nu_\tau, \bar{\nu}_\tau$) $= \frac{7}{16} \rho_\gamma \left(\frac{T_\nu}{T}\right)^4$ is the mass density of each type of neutrino and antineutrino, T_ν is the neutrino temperature, n_b is the number density of baryons, and Y_i , A_i , and Z_i designate the mass fraction, mass number and atomic number of the i th nucleus.

From the assumptions of flux conservation and the presence of a conducting medium (as appropriate for the universe prior to recombination), we can obtain a simple temperature dependence for B :

$$B \propto R^{-2} \propto T^2.$$

Therefore, the energy density of the magnetic field has the same temperature dependence as the energy density of the leptons and that of the radiation field.

We now define

$$\begin{aligned}\rho &= \rho_1 + \rho_2, & p &= p_1 + p_2, \\ \rho_1 &\equiv \rho_\gamma + \rho_{e^-} + \rho_{e^+} + \rho_B, & p_1 &\equiv p_\gamma + p_{e^-} + p_{e^+} + p_B, & \rho_2 &\equiv \rho_\nu + \rho_b, & p_2 &= p_\nu + p_b, \\ \chi &\equiv \frac{\rho_B}{\rho_\gamma + \rho_e + \rho_\nu + \rho_b} \equiv \frac{\rho_B}{\rho_0}, & \rho_0 &\equiv \rho(B=0).\end{aligned}\tag{2.4}$$

The relation between the magnetic field and radiation is thus

$$\rho_B/\rho_\gamma \simeq \frac{43}{8} \chi.\tag{2.5}$$

Substituting these into equations (2.2) and (2.1), noticing the following approximations

$$\rho_b \ll \rho_\nu \rightarrow \rho_2 \simeq \rho_\nu, \quad p_b \ll p_\nu \rightarrow p_2 \simeq p_\nu,\tag{2.6}$$

and using the fact that $\rho_\nu \propto R^{-4}$, we obtain

$$\frac{dR}{dT} = \frac{-R}{3[\rho_1(T) + \frac{p_1}{c^2}(T)]} \frac{d\rho_1}{dT}.\tag{2.7}$$

By using the expansion rate

$$\frac{1}{R} \frac{dR}{dt} = \pm \left(\frac{8\pi G}{3} \rho \right)^{1/2},\tag{2.8}$$

where G is the gravitational constant, the relation between the photon temperature and the time is found to be

$$\frac{dT}{dt} = \mp \left(\frac{8\pi G}{3} \rho \right)^{1/2} [\rho_1(T) + \frac{p_1}{c^2}(T)] \left[\frac{d\rho_1}{dT} \right]^{-1}.\tag{2.9}$$

At high temperatures, $T_\nu = T \propto R^{-1}$, and

$$\rho = \rho_0(1 + \chi) \simeq \frac{43}{8} \rho_\gamma(1 + \chi),\tag{2.10}$$

$$\rho_1 = \frac{11}{4} \rho_\gamma + \rho_B = \frac{11}{4} \rho_\gamma \left(1 + \frac{43}{22} \chi \right) \equiv \rho_{10} \left(1 + \frac{43}{22} \chi \right), \quad \rho_{10} \equiv \rho_1(B=0),\tag{2.11}$$

$$\frac{d\rho_1}{dT} = \left(1 + \frac{43}{22} \chi \right) \frac{d\rho_{10}}{dT} + \frac{43}{22} \rho_{10} \frac{d\chi}{dT},\tag{2.12}$$

$$\frac{d\rho_{10}}{dT} \simeq \beta T_9^3, \quad \beta = 9.262 \times 10^{-8}.\tag{2.13}$$

Incorporating Eq.(2.5) into Eq.(2.8), we obtain a final expression for the relationship between photon temperature and time

$$\frac{dT}{dt} = \mp \frac{\left(\frac{8\pi G}{3}\rho_0\right)^{1/2}(1+\chi)^{1/2}\rho_{10}\left[1 + \frac{43}{22}\chi + \frac{1}{3} + \frac{p_B}{\rho_{10}c^2}\right]}{\left(1 + \frac{43}{22}\chi\right)\beta T_9^3 + \frac{43}{22}\rho_{10}\frac{d\chi}{dT}}. \quad (2.14)$$

Moreover, considering the fact that, $B \propto T^2$, $\rho_B \propto T^4$, and $\rho_0 \simeq \frac{43}{8}\rho_\gamma \propto T^4$, we can introduce a convenient invariant measure of magnetic field strength and assume that the ratio of the magnetic energy density (ρ_B) to the total other energy density (ρ_0) is nearly a constant during BBN. This gives $\frac{d\chi}{dT} = 0$. We now consider two cases:

1. *A global zero magnetic pressure* ($p_B = 0$, but $B \neq 0$):

Physically, this corresponds to a situation where there exists a non-zero local uniform magnetic field (inside each bubble) but a zero total magnetic pressure (pressure free) due to the random distribution of the tangled magnetic bubbles. In this case, Eq.(2.14) becomes

$$\frac{dT}{dt} = \mp \frac{\left(\frac{8\pi G}{3}\rho_0\right)^{1/2}(1+\chi)^{1/2}\left(1 + \frac{129}{88}\chi\right)}{3\left(1 + \frac{43}{22}\chi\right)} T_9. \quad (2.15)$$

Integrating, we obtain

$$T_9 = \kappa \frac{\left(1 + \frac{43}{22}\chi\right)^{1/2}}{\left(1 + \chi\right)^{1/4}\left(1 + \frac{129}{88}\chi\right)^{1/2}} t^{-1/2}, \quad (2.16)$$

where

$$\kappa = \left(\frac{12\pi G a g_{eff}}{2c^2}\right)^{-1/4} \simeq \begin{cases} 10.4, & \text{if } N_\nu = 2 \text{ and } g_{eff} = 9; \\ 4.7, & \text{if } N_\nu = 3 \text{ and } g_{eff} = \frac{43}{4}, \end{cases}$$

in c.g.s. units, g_{eff} is the "effective" number of relativistic degrees of freedom (helicity states), and a is the Stefan black-body constant.

2. *A non-zero magnetic pressure* ($p_B = \rho_B c^2$):

If the magnetic field inside each bubble and the distribution of the magnetic bubbles are not so chaotic, for example, if each magnetic bubble is dipole-like, we will have an averaged magnetic pressure $p_B \propto \rho_B c^2$. In this instance, Eq.(2.14) becomes

$$\frac{dT}{dt} = \mp \frac{\left(\frac{8\pi G}{3}\rho_0\right)^{1/2}(1+\chi)^{1/2}\left(\frac{1}{3} + \frac{43}{44}\chi\right)}{4\left(1 + \frac{43}{22}\chi\right)} T_9 \quad (2.17)$$

and integration yields

$$T_9 = \kappa \frac{\left(1 + \frac{43}{22}\chi\right)^{1/2}}{\left(1 + \chi\right)^{1/4}\left(1 + \frac{129}{44}\chi\right)^{1/2}} t^{-1/2}, \quad (2.18)$$

Note that in the limit when the magnetic fields are absent or very weak ($\rho_B = 0$; $\chi = 0$, or $\chi \ll 1$), Eqs.(2.16) and (2.18) both reduce to

$$T_9 \simeq \kappa t^{-1/2}. \quad (2.19)$$

This is just the formula used in standard BBN calculations.^[14,15]

If the magnetic field is very strong, $\chi \gg 1$, then equation (2.16) and equation (2.18) become, respectively,

$$T_9 = \sqrt{\frac{4}{3}}\chi^{-1/4} \kappa t^{-1/2}, \quad \text{and} \quad T_9 = \sqrt{\frac{2}{3}}\chi^{-1/4} \kappa t^{-1/2}, \quad (2.20)$$

The dependences of the temperature on the time t and the magnetic parameter χ , in the presence of a strong magnetic field, are shown in Figures 1a and 1b. These relations clearly indicate that the effect of the presence of a strong magnetic field on the expansion rate of the universe are indeed significant.

Now we introduce another magnetic parameter $\gamma = B/(2B_c)$, where $B_c = \frac{m_e^2 c^3}{e\hbar} = 4.414 \times 10^{13}$ gauss is the field strength where quantized cyclotron line effects begin to occur; we will refer to this as a quantum critical field value^[16] (See Appendix). According to our definition of the factor χ , we have

$$\chi \equiv \frac{B^2/8\pi}{43\rho_\gamma/8} = \frac{B^2}{43\pi a T^4} \simeq 0.76 \frac{\gamma^2}{T_{10}^4}, \quad (2.21)$$

where a is the Black-Body constant and $T_{10} = T/(10^{10} K)$. If we use the ‘‘critical’’ temperature $T_c \sim 1.28 \times 10^{10} K$ (note: $nkT_c \sim B_c^2/8\pi$, k is Boltzmann constant, $n \simeq 20T^3$ is number density of particles.). Eq.(2.21) can be re-expressed as

$$\chi \simeq 0.283 \left(\frac{T_c}{T}\right)^4 \gamma^2. \quad (2.22)$$

3. Numerical Results and a Limit on the Field Strength and Field Coherence Length

We will now take into account the two independent effects of magnetic fields on reaction rates and on expansion rates calculated in Ref. [12] and Section 2, respectively, in a reexamination of big bang nucleosynthesis. We use the new version of the Wagoner code developed by Kawano^[16]. Specifically, we have replaced the old formulae in the code, for both the reaction rates and the expansion rate, with the new derived equations (3.8), (3.9), (3.10), (3.11), in Ref.[12], and (2.18) above, and calculated the abundances. These are to be compared with observational data, to determine the implied constraints on the

strength and coherence scales of a primordial magnetic field. The observed abundances used are those summarized by Walker et al 1991^[2]. The main technique used is to adjust γ until the calculated abundances no longer match the observational data.

In order to obtain a limit on the strength of primordial magnetic fields and to focus on the effects on BBN explicitly, we have fixed all model parameters other than γ in our calculations: in particular, the neutron lifetime τ_n , the number of neutrino species N_ν , and the baryon to photon ratio $\eta \equiv \frac{n_b}{n_\gamma}$. For our purpose, we adopt the following values for these parameters^[2]:

$$\begin{aligned} \tau_n &= 889.6 \pm 2.9s, & N_\nu &= 3, \\ 2.8 \times 10^{-10} &\leq \eta \leq 4.0 \times 10^{-10}. \end{aligned} \quad (3.1)$$

Moreover, we have assumed non-degenerate neutrinos ($\phi_e = 0, \phi_\nu = 0$) (Thomas, Olive, and Schramm^[17], Steigman and Kang^[18]). For these choices, we then compute the primordial abundances numerically. Our computational results are displayed in both tables 1-3 and figures 2-4. Each figure contains seven sub-figures ((a), (b), (c), (d), (e), (f), and (g)), which represent the abundances of the elements for different strengths of the primordial magnetic fields on a coherence scale of L ($<$ the event horizon at that epoch $\sim 2 \times 10^{12}$ cm).

As expected, our calculations reveal that the abundances of the light elements can be dramatically affected by a strong magnetic field. For instance, if the magnetic fields on scales less than the horizon are as strong as $B \geq 10^{13}$ gauss, the abundances of most elements (except for protons) increase manifestly. In particular, the concentrations of 2H , 4He , 6Li , 7Li , 9Be , ^{14}N , all are enhanced. Some elements, for example, 4He , 7Li , 8Li , ^{11}B , ^{12}C , ^{13}C , ^{14}C , and ^{15}N , show sharp increases (i.e., $^4He \geq 0.5$, $^7Li/H \geq 8.43 \times 10^{-9} \sim 10^{-8}$, $^8Li/H \geq 7.5 \times 10^{-14} \sim 10^{-13}$, $^{11}B/H \sim 10^{-15}$, etc.), as illustrated in figures 1a-1d (I=2,3,4) and tables 1-3. On the contrary, if the magnetic fields are as weak as 10^{11} gauss, the emerging abundances of the light elements, according to our calculations, are only affected slightly and the variations become negligible. Figures 1e-1g (I=2,3,4) display these outcomes.

For comparison, we have also computed the effects on the abundances resulting from the energy density of the magnetic field only (without any variations on the reaction rates). We find that the effects on the abundances of the light elements from reaction rates dominate the contributions from the energy density, unless the magnetic field is very intense ($B > 10^{13}$ Gauss). These results are shown in Fig.5 and Fig.6, respectively. Also, as is true for the standard BBN model, a high ratio n_b/n_γ will globally enhance the high A element abundances.

We note that the observed mass fraction of helium is approximately (Skillman 1993)^[19]

$$Y_p^{\text{obs}} = 0.235 \pm 0.01. \quad (3.2)$$

For other elements, the adopted primordial abundances are displayed in Table 4 (Walker et al 1991)^[2]. By comparing the predicted abundances in the presence of B-fields with the observational determinations, (using particularly the abundance of ${}^4\text{He}$), we can constrain the B-field on scales less than the horizon. Figures I(a,b,c,d,e,f,g) (I = 2,3,4) and tables 1-3 show the abundances of the elements in the presence of magnetic fields, with field strengths ranging from zero to $B = 8.8 \times 10^{14}$ gauss. Through the comparison between our numerical calculations and the observational results, we ascertain that to keep the abundances of light elements compatible with the observations, the primordial magnetic fields at the BBN epoch ($\sim 1\text{min.}$) must satisfy the requirement

$$\gamma \leq 0.001, \quad B \leq 10^{11}G, \quad (3.3)$$

on scales less than the horizon. At this limit, the calculated abundances of light elements are shown in Table 5.

Incorporating our above upper limit into equations (2.21) and (2.5), we can further estimate the ratio of the energy density between magnetic fields and cosmic radiations at the BBN epoch $T_{10} \sim 0.1$ as

$$\frac{\rho_B}{\rho_\gamma} \sim 4\%. \quad (3.4)$$

4. Empirical Estimates of the Strength and the Coherence Scales of the Primeval Fields

We now come to the comparison of the diffusion time of magnetic fields with the expansion time of the universe. The characteristic diffusion time for magnetic fields on a scale $l(\sim L(\frac{R}{R_{\text{nuc}}}))$ is ^[20]

$$\tau_d \simeq \frac{4\pi l^2 \sigma}{c^2}. \quad (4.1)$$

Here R is the cosmological scale factor, and L and R_{nuc} are, respectively, the coherence scale of the B-field and the scale factor at the BBN epoch. Since the expansion time τ_{exp} of the universe (or Hubble time) goes as

$$\tau_{\text{exp}} \propto R^2,$$

we also have

$$\tau_d \sim \frac{4\pi l^2 \sigma}{c^2} \frac{\tau_{\text{exp}}}{t_{\text{nuc}}} \sigma. \quad (4.2)$$

where t_{nuc} is the time of the BBN epoch, $\sigma = \frac{c^2}{4\pi\eta}$ s⁻¹ is the electrical conductivity of the universe, $\eta \simeq \frac{4 \times 10^{12} Z \ln \Lambda}{T_c^{3/2}}$ esu describes the Spitzer resistivity (for electron-proton collisions)

due to electron collisions with neutral hydrogen^[21], T_e denotes the electron temperature in K, Z is the mean charge of the plasma, and Λ is a so called Coulomb integral, which has a typical value around 15 ± 5 .

The ratio of the diffusion time to the Hubble time can thus be estimated as

$$\frac{\tau_d}{\tau_{\text{exp}}} \sim 17.5 \left(\frac{L}{\text{cm}}\right)^2 \tau_{\text{exp}}^{-3/4}, \quad (4.3)$$

where we have used the approximations: $t_{\text{nuc}} \sim 1\text{s}$ and $T_{\text{nuc}} \sim 10^{10}\text{K}$. This ratio corresponding to a given physical scale is a monotonically decreasing function of time. Therefore scales that can not dissipate at recombination ($\sim 10^{12} - 10^{13}\text{s}$) could not have dissipated earlier. Taking the calculation at recombination, we find that if the coherence scale of the magnetic fields at the BBN epoch is larger than 10^4cm ($\sim 10^{-6}$ of the horizon scale) at that time, the field will not be dissipated prior to recombination. The dependence of the ratio with the expansion time is shown in Figure 8.

At the BBN epoch, based on our numerical results in Section 3, the primordial magnetic fields (or magnetic bubbles) at the BBN epoch are constrained as $B \leq 10^{11}\text{gauss}$ on scales of $L \geq 10^4\text{cm}$ (and $L \leq 10^{12}\text{cm}$). To the extent that the universe is a good conductor, this primeval field will evolve to the recombination era by relation $B \propto R^{-2}$ (and $L \propto R$). The implied magnetic field at the time of recombination, prior to the structure formation of the universe, would thus be

$$B_{\text{rec}} \leq 0.1 \text{ gauss}, \quad (4.4)$$

coherent on scales of $L_{\text{domain}} \geq 10^{10}\text{cm}$ (and $\leq 10^{18}\text{cm}$). On scales much larger than the size of the magnetic domains (or bubbles), the physical mechanisms driving field generation are uncorrelated. To put this in current perspective, Hogan^[22] has estimated with certain assumptions that such a field at recombination would correspond to a intergalactic field limit today of $\leq 7 \times 10^{-9}\text{G}$.

5. Conclusions and Future Work

The effects of the magnetic fields on big bang nucleosynthesis and the cosmological expansion rate have been investigated generally in this paper for coherent and chaotic fields on scales smaller than the event horizon. An upper limit has been provided on the strength of the primordial magnetic field on scales smaller than the event horizon. Our results show that, in the framework of standard big bang nucleosynthesis, the maximum strength of the primordial magnetic field on scales greater than 10^4cm but smaller than the horizon at the BBN epoch ($\sim 10^{12}\text{cm}$), can only be 10^{11} gauss , which implies that the magnetic

fields at recombination time would in principle be no stronger than 0.1 gauss. Moreover, in our calculations, we find that, of the two major effects of a primordial magnetic field, those arising from modification of the reaction rates will dominate those arising from modification of the expansion rate (or B-field energy density), unless the magnetic field is very intense ($B \gg 10^{13}$ gauss.)

Finally, we here would like to make two comments:

a. *Rates fluctuations (or Inhomogeneous model).*

If the magnetic field is not uniform or the size of the magnetic bubbles is much smaller than the horizon scale and the bubbles are disconnected from each other, the nuclear reaction rates will become inhomogeneous; i.e., the reaction rates inside a region will differ from those outside the region, even though the geometry of the universe is still not affected. This would require that we introduce reaction rate fluctuations into the big bang calculation (similar to the introduction of the density fluctuations associated with the first order QCD phase transition), and perform multizone calculations.

b. *Anisotropy (or effects on geometry).*

If the size of the magnetic bubbles were larger than the horizon scale, the effects on the geometry of the universe would need to be examined, and the Robertson-Walker metric would need to be replaced by other metrics. In addition, an anisotropy of the universe would result which might have important galactic consequences. (This has been explored by Thorne, 1967, but not with the more extensive network used here.)

A subsequent paper will examine these coherence scales.

Acknowledgements

We would like to thank A. V. Olinto for very helpful discussions. This research was supported in part by the NSF grants AST-93-96039 and AST-92-17969 at the University of Chicago, in part by NSF grant AST-90-22629, DOE grant DE FG0291ER40606 and NASA grant NAGW 1321 at the University of Chicago, and in part by the DOE and by NASA through grant NAGW 2381 at Fermilab.

APPENDIX

Quantum Mechanical Considerations

The applicability of classical electrodynamics to electrons requires that the wavelength of the synchrotron line in the presence of the magnetic field be much larger than the distances of order of $\hbar/m_e c$, for which classical electromagnetics will break down because of quantum effects^[23]. Let us now derive the limit at which classical electrodynamics leads to internal contradictions^[24].

Consider a system in which a charge e , with mass m , moves in a uniform magnetic field B . The synchrotron frequency ω of motion (angular frequency) can then be written as

$$\omega = e c B / E, \quad \text{or} \quad \lambda \sim \frac{m c^2}{e B}, \quad (\text{A.1})$$

where E is the total energy of the charged particle and λ is the wavelength of the synchrotron line.

If quantum effects become important, we would have

$$E = m c^2 = \hbar \omega, \quad \text{or} \quad \lambda \sim \frac{\hbar}{m c}. \quad (\text{A.2})$$

The magnetic field will then satisfy

$$B \sim B_c = \frac{\omega_c m c^2}{e c} = \frac{m^2 c^3}{e \hbar} = 4.4 \times 10^{13} \text{G}, \quad (\text{A.3})$$

which presents a limit where quantum electrodynamics plays a role. Therefore, Eq.(A.3) could be considered in some sense as a quantum critical limit for an organized primordial magnetic field on large scales. For this critical value, we can estimate the corresponding temperature by letting

$$B_c^2 / 8\pi = n k T_c, \quad n \simeq 20 T^3 K^{-3}, \quad (\text{A.4})$$

thus

$$T_c = \left(\frac{B_c^2}{160\pi k} \right)^{1/4} \simeq 1.28 \times 10^{10} \text{K}. \quad (\text{A.5})$$

Note that T_c is comparable to the temperature immediately prior to the BBN epoch. This means that, if the primordial magnetic field prior to the BBN epoch were to be as strong as $B > 10^{13}$ gauss, then it's origin would probably be some quantum process in the early universe. For such conditions, we would not have a field organized on a large scale as the universe expanded out of the quantum domain, because most of the energy released would be converted into small-wave-length radiation rather than into an ordered magnetic field. It is interesting that our numerical calculations ($B < 10^{11}$ gauss) in Section 3 appear to have ruled this possibility out. Instead, it could be suggested that any primordial magnetic field must have been initially in the classical regime $B < 10^{11}$ gauss.

REFERENCES

- [1] D. N. Schramm and R. V. Wagoner, *Ann. Rev. Nucl. & Part. Sci.*, (1977).
- [2] T. P. Walker, G. Steigman, D. N. Schramm, K. A. Olive, and H-S Kang, *Astrophys. J.*, 376 (1991) 51. P.24.
- [3] G. Steigman, D. N. Schramm, and J. Gunn, *Phys. Lett. B* 66 (1977) 202; J. Yang, M. S. Turner, G. Steigman, D. N. Schramm, and K. A. Olive, *Astrophys. J.*, 281 (1984) 493.
- [4] K. S. Thorne, *Astrophys. J.* 148 (1967) 51.
- [5] Ya. B. Zel'dovich and I. D. Novikov, *Relativistic Astrophysics, Vol II The Structure and Evolution of the Universe*, (Chicago: University of Chicago Press, 1983)
- [6] S. W. Hawking and R. J. Tayler, *Nature*, 209 (1966) 1278.
- [7] R. Juszkiewicz, S. Bajtlik, and K. Gorski, *Mon. Not. R. Astron. Soc.* 204 (1983) 63.
- [8] Ya. B. Zel'dovich, *Astron. J. (SSSR)*, 60 (1969) 656.
- [9] G. Greenstein, *Nature*, 223 (1969) 938.
- [10] J. D. Barrow, *Mon. Not. R. Astro. Soc.*, 175 (1976) 379; R. F. O'Connell and J. J. Matese, *Nature*, 222 (1969) 649.
- [11] T. Tajima, S. Cable, K. Shibata, and R. M. Kulsrud, *Astrophys. J.* 390 (1992) 309.
- [12] B. Cheng, D. N. Schramm, and J. W. Truran, Submitted to *Phys. Lett. B*.
- [13] H. Hurki-Suonio, R. A. Matzner, K. A. Olive, and D. N. Schramm, *Astrophys. J.*, 353 (1990) 406; J. H. Applegate, C. J. Hogan, and R. J. Scherrer, *Phys. Rev. D*, 35 (1987) 1151.
- [14] R. Wagoner, *Astrophys. J. Suppl. series* 162 Vol 18 (1969) 247.
- [15] P. J. E. Peebles, *Astrophys. J.*, 146 (1966) 542; R. V. Wagoner, W. A. Fowler, and F. Hoyle, *Astrophys. J.*, 148 (1966) 3.
- [16] L. Kawano, *nucl23 big bang code*.
- [17] Thomas, Olive, and Schramm, *Astrophys. J.*, 406 (1993) 569.
- [18] Steigman and Kang, *Nucl. Phys. B*, 372 (1992) 494.
- [19] Skillman, Preprint.
- [20] T. G. Cowling, *Proc. Roy. Soc. London A*, 183 (1945) 453.
- [21] L. Spitzer, Jr., *Physics of Fully Ionized Gases*, (New York: Interscience: 1962); Ya. B. Zel'dovich and I. D. Novikov, *Relativistic Astrophysics, Vol 2* (Chicago: University of Chicago Press, 1983), p.200.
- [22] C. J. Hogan, *Phys. Rev. Lett.*, 51 (1983) 1488.
- [23] H. Euler and B. Kockel, *Naturewissenschaften*, 23 (1935) 246.
- [24] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields* (Addison - Wesley Publishing Company, Inc., Reading Mass., 1965), revised 2nd ed., Chap. 9.

Figure Captions

Fig. 1a The dependence of T_9 with t under strong B-field but $P_B = 0$.

Fig. 1b The dependence of T_9 with t under strong B-field but $P_B \neq 0$.

Fig. 2 Abundance of elements (n - ${}^8\text{Li}$) in the presence of B-fields but non degenerate neutrinos.

Fig. 3 Abundance of the elements (${}^8\text{B}$ - ${}^{12}\text{C}$) in the presence of B-fields but nondegenerate neutrinos.

Fig. 4 Abundance of the elements (${}^{12}\text{N}$ - ${}^{15}\text{O}$) in the presence of B-fields but nondegenerate neutrinos.

Fig. 5 Abundance of the elements affected by energy density of B-field($\gamma = 10$) only.

Fig. 6 Abundance of the elements affected by energy density of B-field($\gamma = 100$) only.

Fig. 7 The ratio of diffusion time of magnetic fields to Hubble time.

Table Captions

Table 1 Abundance of elements ($n - {}^8\text{Li}$) in the presence of B-fields but non degenerate neutrinos.

Table 2 Abundance of the elements (${}^8\text{B} - {}^{12}\text{C}$) in the presence of B-fields but nondegenerate neutrinos.

Table 3 Abundance of the elements (${}^{12}\text{N} - {}^{15}\text{O}$) in the presence of B-fields but nondegenerate neutrinos.

Table 4 Observed abundances.

Table 5 The calculated abundances at limit $B \leq 10^{11}\text{G}$.

Table 1: Abundance of the Elements ($n-^8\text{Li}$) in the Presence of B-Fields But Nondegenerate Neutrinos.

Element	B-Field									
	$\gamma = 10.$	$\gamma = 1.$	$\gamma = 0.5$	$\gamma = 0.1$	$\gamma = 0.05$	$\gamma = 0.001$	$\gamma = 0$			
$\phi_e = 0$	4.58×10^{-14}	2.22×10^{-14}	1.12×10^{-14}	9.15×10^{-15}	8.25×10^{-15}	5.81×10^{-15}	5.77×10^{-15}			
n	0.441	0.495	0.624	0.716	0.751	0.778	0.777			
^1H	1.52×10^{-3}	1.19×10^{-3}	7.5×10^{-4}	5.59×10^{-4}	5.02×10^{-4}	4.65×10^{-4}	4.65×10^{-4}			
^2H	4.7×10^{-6}	3.64×10^{-6}	2.25×10^{-6}	1.67×10^{-6}	1.5×10^{-6}	1.39×10^{-6}	1.39×10^{-6}			
^3H	6.78×10^{-5}	5.68×10^{-5}	4.15×10^{-5}	3.48×10^{-5}	3.28×10^{-5}	3.15×10^{-5}	3.15×10^{-5}			
^4He	0.555	0.501	0.373	0.281	0.248	0.219	0.219			
^6Li	4.55×10^{-13}	2.83×10^{-13}	1.03×10^{-13}	4.96×10^{-14}	3.7×10^{-14}	2.95×10^{-14}	2.93×10^{-14}			
^7Li	1.41×10^{-8}	8.43×10^{-9}	2.66×10^{-9}	1.15×10^{-9}	8.23×10^{-10}	6.37×10^{-10}	6.37×10^{-10}			
^7Be	2.03×10^{-13}	2.34×10^{-13}	2.89×10^{-13}	3.3×10^{-13}	3.43×10^{-13}	3.54×10^{-13}	3.54×10^{-13}			
^8Li	1.45×10^{-13}	7.46×10^{-14}	1.54×10^{-14}	4.41×10^{-15}	2.58×10^{-15}	1.68×10^{-15}	1.68×10^{-15}			

Table 2: Abundance of the Elements (${}^8\text{B}$ - ${}^{12}\text{C}$) in the Presence of B-Fields But Nondegenerate Neutrinos.

Element	B-Field							
	$\gamma = 10.$	$\gamma = 1.$	$\gamma = 0.5$	$\gamma = 0.1$	$\gamma = 0.05$	$\gamma = 0.001$	$\gamma = 0$	
${}^8\text{B}$	2.27×10^{-25}	2.02×10^{-25}	1.60×10^{-25}	1.40×10^{-25}	1.33×10^{-25}	1.29×10^{-25}	1.29×10^{-25}	
${}^9\text{Be}$	1.20×10^{-24}	3.11×10^{-25}	1.60×10^{-25}	1.40×10^{-25}	1.33×10^{-25}	1.29×10^{-25}	1.29×10^{-25}	
${}^{10}\text{B}$	1.05×10^{-20}	1.58×10^{-20}	2.65×10^{-20}	2.93×10^{-20}	2.79×10^{-20}	2.59×10^{-20}	2.55×10^{-20}	
${}^{11}\text{B}$	4.32×10^{-16}	1.91×10^{-15}	3.03×10^{-16}	7.79×10^{-17}	4.48×10^{-17}	2.88×10^{-17}	2.88×10^{-17}	
${}^{11}\text{C}$	2.27×10^{-25}	2.02×10^{-25}	1.60×10^{-25}	1.40×10^{-25}	1.33×10^{-25}	1.29×10^{-25}	1.29×10^{-25}	
${}^{12}\text{B}$	2.27×10^{-25}	2.02×10^{-25}	1.60×10^{-25}	1.40×10^{-25}	1.33×10^{-25}	1.29×10^{-25}	1.29×10^{-25}	
${}^{12}\text{C}$	1.39×10^{-13}	7.19×10^{-14}	1.48×10^{-14}	4.27×10^{-15}	2.50×10^{-15}	1.63×10^{-15}	1.60×10^{-15}	

Table 3: Abundance of the Elements ($^{12}\text{N}-^{15}\text{O}$) in the Presence of B-Fields But Nongenerate Neutrinos.

Element	B-Field							
	$\gamma = 10.$	$\gamma = 1.$	$\gamma = 0.5$	$\gamma = 0.1$	$\gamma = 0.05$	$\gamma = 0.001$	$\gamma = 0$	
^{12}N	2.27×10^{-25}	2.02×10^{-25}	1.60×10^{-25}	1.40×10^{-25}	1.33×10^{-25}	1.29×10^{-25}	1.29×10^{-25}	
^{13}C	1.26×10^{-16}	7.30×10^{-16}	1.85×10^{-16}	5.96×10^{-17}	3.61×10^{-17}	2.38×10^{-17}	2.35×10^{-17}	
^{13}N	2.27×10^{-25}	2.02×10^{-25}	1.60×10^{-25}	1.40×10^{-25}	1.33×10^{-25}	1.29×10^{-25}	1.29×10^{-25}	
^{14}C	2.77×10^{-16}	1.31×10^{-16}	1.99×10^{-17}	4.15×10^{-18}	2.07×10^{-18}	1.17×10^{-18}	1.15×10^{-18}	
^{14}N	4.76×10^{-18}	4.17×10^{-18}	2.06×10^{-18}	9.24×10^{-19}	6.18×10^{-19}	4.36×10^{-19}	4.30×10^{-19}	
^{14}O	2.27×10^{-25}	2.02×10^{-25}	1.60×10^{-25}	1.40×10^{-25}	1.33×10^{-25}	1.29×10^{-25}	1.29×10^{-25}	
^{15}N	1.94×10^{-20}	9.31×10^{-21}	1.55×10^{-21}	3.95×10^{-22}	2.26×10^{-22}	1.44×10^{-22}	1.44×10^{-22}	
^{15}O	2.27×10^{-25}	2.02×10^{-25}	1.60×10^{-25}	1.40×10^{-25}	1.33×10^{-25}	1.29×10^{-25}	1.29×10^{-25}	

Table 4: Observed Abundances.

Element	Where Observed	Mass Fraction
D	presolar	$1.8 \times 10^{-5} \leq y_2 \leq 3.3 \times 10^{-5} (2\sigma)$
3He	presolar	$1.3 \times 10^{-6} \leq y_3 \leq 1.8 \times 10^{-5} (2\sigma)$
$D + {}^3He$	presolar	$3.3 \times 10^{-5} \leq y_{23} \leq 4.9 \times 10^{-5} (2\sigma)$
4He	H II region	$Y_p = 0.23 \pm 0.007 (\sigma_Y = 0.009)$
7Li	pre-Population II	$12 + \log({}^7Li/H) \leq 2.15 (95\%CL)$
6Li	pre-Population II	$\leq 0.1 {}^7Li$
9Be	Hyades	$\leq (1 - 2) \times 10^{-11}$
$\geq {}^{12}X$	—	$\leq 10^{-12}$

Table 5: The Calculated Abundances at Limit $B \leq 10^{11} G$.

LightElement	Abundance
D/H	$\leq 4.65 \times 10^{-4}$
T/H	$\leq 1.39 \times 10^{-6}$
${}^3He/H$	$\leq 3.17 \times 10^{-5}$
4He	$= 0.22$
${}^6Li/H$	$\leq 2.95 \times 10^{-14}$
${}^7Li/H$	$\leq 6.374 \times 10^{-10}$
${}^7Be/H$	$\leq 3.54 \times 10^{-13}$
${}^8Li/H$	$\leq 1.68 \times 10^{-15}$

Fig. 1a The dependence of T_θ with t under strong B-field but $P_B = 0$




Fig. 1b The dependence of T_9 with t under strong B-field but $P_B \neq 0$




Fig. 2 Abundance of elements ($n - {}^8\text{Li}$) in the presence of B-fields
but non degenerate neutrinos




Fig. 2






Fig. 3 Abundance of the elements (${}^8\text{B} - {}^{12}\text{C}$) in the presence of B-fields but nondegenerate neutrinos




Fig. 3






Fig. 4 Abundance of the elements ($^{12}\text{N} - ^{15}\text{O}$) in the presence of B-fields but nondegenerate neutrinos




Fig. 4






Fig. 5 Abundance of the elements affected by energy density of B-field($\gamma = 10$) only






Fig. 6 Abundance of the elements affected by energy density of B-field($\gamma = 100$) only






Fig. 7 The ratio of diffusion time of magnetic fields to Hubble time

