



**Fermi National Accelerator Laboratory**

FERMILAB-Pub-98/075-E

D0

**Measurement of the Shape of the Transverse Momentum  
Distribution of W Bosons Produced in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.8$  TeV**

B. Abbott et al.  
The D0 Collaboration

*Fermi National Accelerator Laboratory  
P.O. Box 500, Batavia, Illinois 60510*

February 1998

Submitted to *Physical Review Letters*

**Measurement of the Shape of the Transverse Momentum  
Distribution of  $W$  Bosons Produced in  $p\bar{p}$  Collisions at  
 $\sqrt{s} = 1.8$  TeV**

B. Abbott,<sup>31</sup> M. Abolins,<sup>27</sup> B.S. Acharya,<sup>46</sup> I. Adam,<sup>12</sup> D.L. Adams,<sup>40</sup> M. Adams,<sup>17</sup>  
S. Ahn,<sup>14</sup> H. Aihara,<sup>23</sup> G.A. Alves,<sup>10</sup> N. Amos,<sup>26</sup> E.W. Anderson,<sup>19</sup> R. Astur,<sup>45</sup>  
M.M. Baarmand,<sup>45</sup> L. Babukhadia,<sup>2</sup> A. Baden,<sup>25</sup> V. Balamurali,<sup>35</sup> J. Balderston,<sup>16</sup>  
B. Baldin,<sup>14</sup> S. Banerjee,<sup>46</sup> J. Bantly,<sup>5</sup> E. Barberis,<sup>23</sup> J.F. Bartlett,<sup>14</sup> A. Belyaev,<sup>29</sup>  
S.B. Beri,<sup>37</sup> I. Bertram,<sup>34</sup> V.A. Bezzubov,<sup>38</sup> P.C. Bhat,<sup>14</sup> V. Bhatnagar,<sup>37</sup>  
M. Bhattacharjee,<sup>45</sup> N. Biswas,<sup>35</sup> G. Blazey,<sup>33</sup> S. Blessing,<sup>15</sup> P. Bloom,<sup>7</sup> A. Boehnlein,<sup>14</sup>  
N.I. Bojko,<sup>38</sup> F. Borchering,<sup>14</sup> C. Boswell,<sup>9</sup> A. Brandt,<sup>14</sup> R. Brock,<sup>27</sup> A. Bross,<sup>14</sup>  
D. Buchholz,<sup>34</sup> V.S. Burtovoi,<sup>38</sup> J.M. Butler,<sup>3</sup> W. Carvalho,<sup>10</sup> D. Casey,<sup>27</sup> Z. Casilum,<sup>45</sup>  
H. Castilla-Valdez,<sup>11</sup> D. Chakraborty,<sup>45</sup> S.-M. Chang,<sup>32</sup> S.V. Chekulaev,<sup>38</sup> L.-P. Chen,<sup>23</sup>  
W. Chen,<sup>45</sup> S. Choi,<sup>44</sup> S. Chopra,<sup>26</sup> B.C. Choudhary,<sup>9</sup> J.H. Christenson,<sup>14</sup> M. Chung,<sup>17</sup>  
D. Claes,<sup>30</sup> A.R. Clark,<sup>23</sup> W.G. Cobau,<sup>25</sup> J. Cochran,<sup>9</sup> L. Coney,<sup>35</sup> W.E. Cooper,<sup>14</sup>  
C. Cretsinger,<sup>42</sup> D. Cullen-Vidal,<sup>5</sup> M.A.C. Cummings,<sup>33</sup> D. Cutts,<sup>5</sup> O.I. Dahl,<sup>23</sup> K. Davis,<sup>2</sup>  
K. De,<sup>47</sup> K. Del Signore,<sup>26</sup> M. Demarteau,<sup>14</sup> D. Denisov,<sup>14</sup> S.P. Denisov,<sup>38</sup> H.T. Diehl,<sup>14</sup>  
M. Diesburg,<sup>14</sup> G. Di Loreto,<sup>27</sup> P. Draper,<sup>47</sup> Y. Ducros,<sup>43</sup> L.V. Dudko,<sup>29</sup> S.R. Dugad,<sup>46</sup>  
D. Edmunds,<sup>27</sup> J. Ellison,<sup>9</sup> V.D. Elvira,<sup>45</sup> R. Engelmann,<sup>45</sup> S. Eno,<sup>25</sup> G. Eppley,<sup>40</sup>  
P. Ermolov,<sup>29</sup> O.V. Eroshin,<sup>38</sup> V.N. Evdokimov,<sup>38</sup> T. Fahland,<sup>8</sup> M.K. Fatyga,<sup>42</sup> S. Feher,<sup>14</sup>  
D. Fein,<sup>2</sup> T. Ferbel,<sup>42</sup> G. Finocchiaro,<sup>45</sup> H.E. Fisk,<sup>14</sup> Y. Fisyak,<sup>4</sup> E. Flattum,<sup>14</sup>  
G.E. Forden,<sup>2</sup> M. Fortner,<sup>33</sup> K.C. Frame,<sup>27</sup> S. Fuess,<sup>14</sup> E. Gallas,<sup>47</sup> A.N. Galyaev,<sup>38</sup>  
P. Gartung,<sup>9</sup> V. Gavrilov,<sup>28</sup> T.L. Geld,<sup>27</sup> R.J. Genik II,<sup>27</sup> K. Genser,<sup>14</sup> C.E. Gerber,<sup>14</sup>  
Y. Gershtein,<sup>28</sup> B. Gibbard,<sup>4</sup> S. Glenn,<sup>7</sup> B. Gobbi,<sup>34</sup> A. Goldschmidt,<sup>23</sup> B. Gómez,<sup>1</sup>  
G. Gómez,<sup>25</sup> P.I. Goncharov,<sup>38</sup> J.L. González Solís,<sup>11</sup> H. Gordon,<sup>4</sup> L.T. Goss,<sup>48</sup>  
K. Gounder,<sup>9</sup> A. Goussiou,<sup>45</sup> N. Graf,<sup>4</sup> P.D. Grannis,<sup>45</sup> D.R. Green,<sup>14</sup> H. Greenlee,<sup>14</sup>  
S. Grinstein,<sup>6</sup> P. Grudberg,<sup>23</sup> S. Grünendahl,<sup>14</sup> G. Guglielmo,<sup>36</sup> J.A. Guida,<sup>2</sup> J.M. Guida,<sup>5</sup>  
A. Gupta,<sup>46</sup> S.N. Gurzhiev,<sup>38</sup> G. Gutierrez,<sup>14</sup> P. Gutierrez,<sup>36</sup> N.J. Hadley,<sup>25</sup> H. Haggerty,<sup>14</sup>  
S. Hagopian,<sup>15</sup> V. Hagopian,<sup>15</sup> K.S. Hahn,<sup>42</sup> R.E. Hall,<sup>8</sup> P. Hanlet,<sup>32</sup> S. Hansen,<sup>14</sup>  
J.M. Hauptman,<sup>19</sup> D. Hedin,<sup>33</sup> A.P. Heinson,<sup>9</sup> U. Heintz,<sup>14</sup> R. Hernández-Montoya,<sup>11</sup>  
T. Heuring,<sup>15</sup> R. Hirosky,<sup>17</sup> J.D. Hobbs,<sup>45</sup> B. Hoeneisen,<sup>1,\*</sup> J.S. Hoftun,<sup>5</sup> F. Hsieh,<sup>26</sup>  
Ting Hu,<sup>45</sup> Tong Hu,<sup>18</sup> T. Huehn,<sup>9</sup> A.S. Ito,<sup>14</sup> E. James,<sup>2</sup> J. Jaques,<sup>35</sup> S.A. Jerger,<sup>27</sup>  
R. Jesik,<sup>18</sup> J.Z.-Y. Jiang,<sup>45</sup> T. Joffe-Minor,<sup>34</sup> K. Johns,<sup>2</sup> M. Johnson,<sup>14</sup> A. Jonckheere,<sup>14</sup>  
M. Jones,<sup>16</sup> H. Jöstlein,<sup>14</sup> S.Y. Jun,<sup>34</sup> C.K. Jung,<sup>45</sup> S. Kahn,<sup>4</sup> G. Kalbfleisch,<sup>36</sup> J.S. Kang,<sup>20</sup>  
D. Karmanov,<sup>29</sup> D. Karmgard,<sup>15</sup> R. Kehoe,<sup>35</sup> M.L. Kelly,<sup>35</sup> C.L. Kim,<sup>20</sup> S.K. Kim,<sup>44</sup>  
B. Klima,<sup>14</sup> C. Klopfenstein,<sup>7</sup> J.M. Kohli,<sup>37</sup> D. Koltick,<sup>39</sup> A.V. Kostritskiy,<sup>38</sup> J. Kotcher,<sup>4</sup>  
A.V. Kotwal,<sup>12</sup> J. Kourlas,<sup>31</sup> A.V. Kozelov,<sup>38</sup> E.A. Kozlovsky,<sup>38</sup> J. Krane,<sup>30</sup>  
M.R. Krishnaswamy,<sup>46</sup> S. Krzywdzinski,<sup>14</sup> S. Kuleshov,<sup>28</sup> S. Kunori,<sup>25</sup> F. Landry,<sup>27</sup>  
G. Landsberg,<sup>14</sup> B. Lauer,<sup>19</sup> A. Leflat,<sup>29</sup> H. Li,<sup>45</sup> J. Li,<sup>47</sup> Q.Z. Li-Demarteau,<sup>14</sup>  
J.G.R. Lima,<sup>41</sup> D. Lincoln,<sup>14</sup> S.L. Linn,<sup>15</sup> J. Linnemann,<sup>27</sup> R. Lipton,<sup>14</sup> Y.C. Liu,<sup>34</sup>  
F. Lobkowicz,<sup>42</sup> S.C. Loken,<sup>23</sup> S. Lökös,<sup>45</sup> L. Lueking,<sup>14</sup> A.L. Lyon,<sup>25</sup> A.K.A. Maciel,<sup>10</sup>  
R.J. Madaras,<sup>23</sup> R. Madden,<sup>15</sup> L. Magaña-Mendoza,<sup>11</sup> V. Manankov,<sup>29</sup> S. Mani,<sup>7</sup>  
H.S. Mao,<sup>14,†</sup> R. Markeloff,<sup>33</sup> T. Marshall,<sup>18</sup> M.I. Martin,<sup>14</sup> K.M. Mauritz,<sup>19</sup> B. May,<sup>34</sup>  
A.A. Mayorov,<sup>38</sup> R. McCarthy,<sup>45</sup> J. McDonald,<sup>15</sup> T. McKibben,<sup>17</sup> J. McKinley,<sup>27</sup>

T. McMahon,<sup>36</sup> H.L. Melanson,<sup>14</sup> M. Merkin,<sup>29</sup> K.W. Merritt,<sup>14</sup> H. Miettinen,<sup>40</sup>  
 A. Mincer,<sup>31</sup> C.S. Mishra,<sup>14</sup> N. Mokhov,<sup>14</sup> N.K. Mondal,<sup>46</sup> H.E. Montgomery,<sup>14</sup>  
 P. Mooney,<sup>1</sup> H. da Motta,<sup>10</sup> C. Murphy,<sup>17</sup> F. Nang,<sup>2</sup> M. Narain,<sup>14</sup> V.S. Narasimham,<sup>46</sup>  
 A. Narayanan,<sup>2</sup> H.A. Neal,<sup>26</sup> J.P. Negret,<sup>1</sup> P. Nemethy,<sup>31</sup> D. Norman,<sup>48</sup> L. Oesch,<sup>26</sup>  
 V. Oguri,<sup>41</sup> E. Oliveira,<sup>10</sup> E. Oltman,<sup>23</sup> N. Oshima,<sup>14</sup> D. Owen,<sup>27</sup> P. Padley,<sup>40</sup> A. Para,<sup>14</sup>  
 Y.M. Park,<sup>21</sup> R. Partridge,<sup>5</sup> N. Parua,<sup>46</sup> M. Paterno,<sup>42</sup> B. Pawlik,<sup>22</sup> J. Perkins,<sup>47</sup>  
 M. Peters,<sup>16</sup> R. Piegaiia,<sup>6</sup> H. Piekarczyk,<sup>15</sup> Y. Pischalnikov,<sup>39</sup> B.G. Pope,<sup>27</sup> H.B. Prosper,<sup>15</sup>  
 S. Protopopescu,<sup>4</sup> D. Pušeljčić,<sup>23</sup> J. Qian,<sup>26</sup> P.Z. Quintas,<sup>14</sup> R. Raja,<sup>14</sup> S. Rajagopalan,<sup>4</sup>  
 O. Ramirez,<sup>17</sup> L. Rasmussen,<sup>45</sup> S. Reucroft,<sup>32</sup> M. Rijssenbeek,<sup>45</sup> T. Rockwell,<sup>27</sup> M. Roco,<sup>14</sup>  
 P. Rubinov,<sup>34</sup> R. Ruchti,<sup>35</sup> J. Rutherford,<sup>2</sup> A. Sánchez-Hernández,<sup>11</sup> A. Santoro,<sup>10</sup>  
 L. Sawyer,<sup>24</sup> R.D. Schamberger,<sup>45</sup> H. Schellman,<sup>34</sup> J. Sculli,<sup>31</sup> E. Shabalina,<sup>29</sup> C. Shaffer,<sup>15</sup>  
 H.C. Shankar,<sup>46</sup> R.K. Shivpuri,<sup>13</sup> M. Shupe,<sup>2</sup> H. Singh,<sup>9</sup> J.B. Singh,<sup>37</sup> V. Sirotenko,<sup>33</sup>  
 W. Smart,<sup>14</sup> E. Smith,<sup>36</sup> R.P. Smith,<sup>14</sup> R. Snihur,<sup>34</sup> G.R. Snow,<sup>30</sup> J. Snow,<sup>36</sup> S. Snyder,<sup>4</sup>  
 J. Solomon,<sup>17</sup> M. Sosebee,<sup>47</sup> N. Sotnikova,<sup>29</sup> M. Souza,<sup>10</sup> A.L. Spadafora,<sup>23</sup> G. Steinbrück,<sup>36</sup>  
 R.W. Stephens,<sup>47</sup> M.L. Stevenson,<sup>23</sup> D. Stewart,<sup>26</sup> F. Stichelbaut,<sup>45</sup> D. Stoker,<sup>8</sup> V. Stolin,<sup>28</sup>  
 D.A. Stoyanova,<sup>38</sup> M. Strauss,<sup>36</sup> K. Streets,<sup>31</sup> M. Strovink,<sup>23</sup> A. Sznajder,<sup>10</sup>  
 P. Tamburello,<sup>25</sup> J. Tarazi,<sup>8</sup> M. Tartaglia,<sup>14</sup> T.L.T. Thomas,<sup>34</sup> J. Thompson,<sup>25</sup>  
 T.G. Trippe,<sup>23</sup> P.M. Tuts,<sup>12</sup> N. Varelas,<sup>17</sup> E.W. Varnes,<sup>23</sup> D. Vititoe,<sup>2</sup> A.A. Volkov,<sup>38</sup>  
 A.P. Vorobiev,<sup>38</sup> H.D. Wahl,<sup>15</sup> G. Wang,<sup>15</sup> J. Warchol,<sup>35</sup> G. Watts,<sup>5</sup> M. Wayne,<sup>35</sup>  
 H. Weerts,<sup>27</sup> A. White,<sup>47</sup> J.T. White,<sup>48</sup> J.A. Wightman,<sup>19</sup> S. Willis,<sup>33</sup> S.J. Wimpenny,<sup>9</sup>  
 J.V.D. Wirjawan,<sup>48</sup> J. Womersley,<sup>14</sup> E. Won,<sup>42</sup> D.R. Wood,<sup>32</sup> H. Xu,<sup>5</sup> R. Yamada,<sup>14</sup>  
 P. Yamin,<sup>4</sup> J. Yang,<sup>31</sup> T. Yasuda,<sup>32</sup> P. Yepes,<sup>40</sup> C. Yoshikawa,<sup>16</sup> S. Youssef,<sup>15</sup> J. Yu,<sup>14</sup>  
 Y. Yu,<sup>44</sup> Z. Zhou,<sup>19</sup> Z.H. Zhu,<sup>42</sup> D. Zieminska,<sup>18</sup> A. Zieminski,<sup>18</sup> E.G. Zverev,<sup>29</sup>  
 and A. Zylberstein<sup>43</sup>

(DØ Collaboration)

<sup>1</sup>*Universidad de los Andes, Bogotá, Colombia*

<sup>2</sup>*University of Arizona, Tucson, Arizona 85721*

<sup>3</sup>*Boston University, Boston, Massachusetts 02215*

<sup>4</sup>*Brookhaven National Laboratory, Upton, New York 11973*

<sup>5</sup>*Brown University, Providence, Rhode Island 02912*

<sup>6</sup>*Universidad de Buenos Aires, Buenos Aires, Argentina*

<sup>7</sup>*University of California, Davis, California 95616*

<sup>8</sup>*University of California, Irvine, California 92697*

<sup>9</sup>*University of California, Riverside, California 92521*

<sup>10</sup>*LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil*

<sup>11</sup>*CINVESTAV, Mexico City, Mexico*

<sup>12</sup>*Columbia University, New York, New York 10027*

<sup>13</sup>*Delhi University, Delhi, India 110007*

<sup>14</sup>*Fermi National Accelerator Laboratory, Batavia, Illinois 60510*

<sup>15</sup>*Florida State University, Tallahassee, Florida 32306*

<sup>16</sup>*University of Hawaii, Honolulu, Hawaii 96822*

<sup>17</sup>*University of Illinois at Chicago, Chicago, Illinois 60607*

<sup>18</sup>*Indiana University, Bloomington, Indiana 47405*

- <sup>19</sup>*Iowa State University, Ames, Iowa 50011*
- <sup>20</sup>*Korea University, Seoul, Korea*
- <sup>21</sup>*Kyungshung University, Pusan, Korea*
- <sup>22</sup>*Institute of Nuclear Physics, Kraków, Poland*
- <sup>23</sup>*Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720*
- <sup>24</sup>*Louisiana Tech University, Ruston, Louisiana 71272*
- <sup>25</sup>*University of Maryland, College Park, Maryland 20742*
- <sup>26</sup>*University of Michigan, Ann Arbor, Michigan 48109*
- <sup>27</sup>*Michigan State University, East Lansing, Michigan 48824*
- <sup>28</sup>*Institute for Theoretical and Experimental Physics, Moscow, Russia*
- <sup>29</sup>*Moscow State University, Moscow, Russia*
- <sup>30</sup>*University of Nebraska, Lincoln, Nebraska 68588*
- <sup>31</sup>*New York University, New York, New York 10003*
- <sup>32</sup>*Northeastern University, Boston, Massachusetts 02115*
- <sup>33</sup>*Northern Illinois University, DeKalb, Illinois 60115*
- <sup>34</sup>*Northwestern University, Evanston, Illinois 60208*
- <sup>35</sup>*University of Notre Dame, Notre Dame, Indiana 46556*
- <sup>36</sup>*University of Oklahoma, Norman, Oklahoma 73019*
- <sup>37</sup>*University of Panjab, Chandigarh 16-00-14, India*
- <sup>38</sup>*Institute for High Energy Physics, Protvino 142284, Russia*
- <sup>39</sup>*Purdue University, West Lafayette, Indiana 47907*
- <sup>40</sup>*Rice University, Houston, Texas 77005*
- <sup>41</sup>*Universidade do Estado do Rio de Janeiro, Brazil*
- <sup>42</sup>*University of Rochester, Rochester, New York 14627*
- <sup>43</sup>*CEA, DAPNIA/Service de Physique des Particules, CE-SACLAY, Gif-sur-Yvette, France*
- <sup>44</sup>*Seoul National University, Seoul, Korea*
- <sup>45</sup>*State University of New York, Stony Brook, New York 11794*
- <sup>46</sup>*Tata Institute of Fundamental Research, Colaba, Mumbai 400005, India*
- <sup>47</sup>*University of Texas, Arlington, Texas 76019*
- <sup>48</sup>*Texas A&M University, College Station, Texas 77843*

## Abstract

The shape of the transverse momentum distribution of  $W$  bosons ( $p_T^W$ ) produced in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV is measured with the DØ detector at Fermilab. The result is compared to QCD perturbative and resummation calculations over the  $p_T^W$  range from 0 – 200 GeV/ $c$ . The shape of the distribution is consistent with the theoretical prediction.

The transverse momentum ( $p_T^W$ ) of  $W$  intermediate vector bosons produced in proton-antiproton collisions is due to the production of one or more gluons or quarks along with the boson. At low transverse momentum ( $p_T^W < 10$  GeV/ $c$ ), multiple soft gluon emission is expected to dominate the cross section. A soft gluon resummation technique [1-5] is therefore used to make QCD predictions. At high transverse momentum ( $p_T^W > 20$  GeV/ $c$ ), the cross section is dominated by the radiation of a single parton with large transverse momentum. Perturbative QCD [6] is therefore expected to be reliable in this regime. A prescription [4] has been proposed for matching the low and high  $p_T^W$  regions to provide a continuous prediction for all  $p_T^W$ . Thus, a measurement of the transverse momentum distribution may be used to check the soft gluon resummation calculations in the low  $p_T^W$  range, and to test the perturbative QCD calculations at high  $p_T^W$ .

The transverse momentum spectrum of  $W$  bosons has been measured previously by the UA1 [7], UA2 [8], and CDF [9] collaborations, but with smaller data samples than the one used here. This paper presents a measurement of the shape of the  $p_T$  spectrum of  $W$  bosons produced in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV with the DØ detector [10] at Fermilab, and extends the  $p_T^W$  range of the previous measurements. The data come from a sample of  $12.4 \pm 0.7$  pb $^{-1}$  collected during the 1992–1993 run. A measurement of the inclusive cross section for  $W$  and  $Z$  boson production based on the same data set has been reported [11] and agrees with QCD predictions.

This measurement uses the decay mode  $W \rightarrow e\nu$ . Electrons were detected in a hermetic uranium/liquid-argon calorimeter with an energy resolution of about 15%/ $\sqrt{E(\text{GeV})}$ . The calorimeter has a transverse granularity of  $\Delta\eta \times \Delta\phi = 0.1 \times 0.1$ , where  $\eta$  is the pseudorapidity and  $\phi$  is the azimuthal angle. Electrons were accepted in the central pseudorapidity region only,  $|\eta| < 1.1$ , to keep the background contamination from multijet events at a reasonably low level for high values of  $p_T^W$ . The transverse momentum of the neutrino was calculated using the calorimetric measurement of the missing transverse energy ( $\cancel{E}_T$ ) in the event. We take the  $p_T^W$  to be the sum of the electron and neutrino transverse momenta, measuring it only from the recoiling hadrons. The analysis used a single electron trigger, which required one electron with transverse energy ( $E_T$ ) greater than 20 GeV.

The offline electron identification requirements consisted of four criteria: (i) the electron had to deposit at least 95% of its energy in the electromagnetic calorimeter (21 radiation lengths deep); (ii) the transverse and longitudinal shower shapes had to be consistent with those expected for an electron [12]; (iii) a good match had to exist between a reconstructed track in the drift chamber system and the shower position in the calorimeter; and (iv) the electron had to be isolated from other energy deposits in the calorimeter, with  $I < 0.1$ . This isolation variable is defined as  $I = [E_{\text{TOT}}(0.4) - E_{\text{EM}}(0.2)]/E_{\text{EM}}(0.2)$ , where  $E_{\text{TOT}}(0.4)$  is the total calorimeter energy inside a cone of radius  $\sqrt{\Delta\eta^2 + \Delta\phi^2} = 0.4$  and  $E_{\text{EM}}(0.2)$  is the electromagnetic energy inside a cone of radius 0.2. To select the  $W$  boson candidate sample, we required one electron with  $E_T > 25$  GeV, and  $\cancel{E}_T > 25$  GeV. Events having a second electron with  $E_T > 20$  GeV that satisfies criteria (i), (ii), and (iv) were excluded from the candidate sample as possible  $Z \rightarrow e^+e^-$  events. Criterion (iii) was not applied to this second electron in order to allow for possible tracking inefficiencies. These selection criteria yielded 7132  $W \rightarrow e\nu$  candidates.

The trigger and selection efficiencies were determined using  $Z \rightarrow e^+e^-$  events in which one of the electrons satisfied the trigger and selection criteria. The second electron then

provided an unbiased sample with which to measure the efficiencies. No dependence of the trigger or selection efficiency on  $p_T^W$  was found, to an accuracy of 5%.

A Monte Carlo program [13] was used to simulate the DØ detector response and calculate the kinematic and geometric acceptance as a function of  $p_T^W$ . The detector resolutions used in the Monte Carlo were determined from data, and were parametrized as a function of energy and angle. The relative response of the hadronic and EM calorimeters was studied using the transverse momentum of the  $Z$  boson as measured by the  $p_T$  of the two electrons compared to the hadronic recoil system in the  $Z$  event. This parametrized representation of the DØ detector was used to smear the theoretical prediction by detector effects and compare it to our measured  $p_T^W$ . We prefer this method of comparison to a standard unfolding procedure [14] that proved to be sensitive to the choice of the prior distribution function. This sensitivity is caused by the Jacobian zero in  $dN/dp_T^W$  at the origin, which induces a peak that appears near  $p_T^W \approx 4$  GeV/ $c$  after it is broadened by our  $p_T^W$  resolution. Only below 4 GeV/ $c$  do the true  $p_T^W$  distributions predicted by the two available models [4,5] exhibit a difference, which is masked in the data by these same resolution effects.

The dominant source of background in the  $W$  boson sample was multijet events where one or more of the jets fluctuated to fake an electron. Some multijet events also have significant  $\cancel{E}_T$  due to fluctuations and mismeasurements of the jet energies. This could fake a neutrino from  $W$  boson decay. The amount of multijet background in the  $W \rightarrow e\nu$  candidate sample was determined by first defining a “loose” event sample which had the same selection criteria as the candidate sample except that electron identification criteria (i) and (ii) were not applied. This loose sample consisted of  $N_s$  signal events and  $N_b$  multijet background events. The  $W \rightarrow e\nu$  candidate sample (described above) consisted of  $\varepsilon_s N_s$  signal events and  $\varepsilon_b N_b$  multijet background events, where  $\varepsilon_s$  and  $\varepsilon_b$  are the electron selection efficiencies for the candidate relative to the loose samples, for signal and background respectively. We obtained  $\varepsilon_s$  from the  $Z \rightarrow e^+e^-$  sample, and  $\varepsilon_b$  from events with  $\cancel{E}_T < 15$  GeV. The total multijet background was determined to be  $(4.2 \pm 2.3)\%$ . The shape of the multijet background as a function of  $p_T^W$  was determined by repeating the procedure in different  $p_T^W$  bins.

To cross-check the multijet background estimate, the transverse mass and the  $\cancel{E}_T$  distributions of the final data sample were compared to a model of the expected distributions obtained from a combination of  $W \rightarrow e\nu$  Monte Carlo events plus the estimated multijet background. The comparison was performed in three  $p_T^W$  bins: 0–30 GeV/ $c$ , 30–60 GeV/ $c$ , and  $> 60$  GeV/ $c$ ; the number of  $W \rightarrow e\nu$  candidates in each bin was 6726, 282, and 124. The amount of multijet background in each  $p_T^W$  bin was estimated as  $(2.9 \pm 1.6)\%$ ,  $(20.9 \pm 11.7)\%$ , and  $(38.3 \pm 21.5)\%$ , respectively. Figure 1 shows the results of the comparisons. The distinctive shape of the transverse mass distribution for multijet background arises from applying the kinematic cuts and the minimum  $p_T^W$  requirement to a sample predominantly composed of dijet events that lie back-to-back in the transverse plane. The goodness of the fit between the data and the model is evaluated by performing a  $\chi^2_\lambda$  test [15]. The numerical results for  $\chi^2_\lambda/\text{d.o.f.}$  for each fit in Fig. 1 are (a) 20.2/25, (b) 26.4/19, (c) 4.7/7, (d) 5.4/9, (e) 3.0/13, and (f) 2.1/7. They correspond to fit probabilities of (a) 74%, (b) 12%, (c) 69%, (d) 80%, (e) 99%, and (f) 95% respectively. We therefore conclude that the tests show good agreement between the data and the expectation.

The normalized multijet background was subtracted bin by bin from the  $W$  boson candidate sample transverse momentum spectrum. Additional corrections (all less than 5%) were

made to account for top quark background events and for  $Z \rightarrow e^+e^-$  events where one of the electrons was lost or not identified. Since  $p_T^W$  was measured from the recoiling hadrons, the events originating from  $W \rightarrow \tau\nu$  (where  $\tau \rightarrow e\nu\nu$ ) contributed properly to the differential distribution; this source of background therefore was included in the Monte Carlo simulation of the  $p_T^W$  distribution.

The normalized distribution of the  $W$  boson transverse momentum ( $\frac{1}{N} \frac{dN}{dp_T^W}$ ) is shown in Fig. 2 and given in Table I. The largest contributions to the systematic error in the  $p_T^W$  measurement are the uncertainty in the magnitude of the multijet background, the uncertainty in the hadronic recoil energy scale factor and resolution used in the detector simulation, and the uncertainty in the selection efficiency. These are all added in quadrature since they are independent. We compare our experimental result to the theoretical prediction [4] computed using the MRSA' [16] parton distribution function and smeared for detector resolutions. The measurement and the prediction are independently area-normalized to unity. These points are used to perform a more detailed comparison between data and theory by plotting the ratio (Data-Theory)/Theory, which is shown in Fig. 3. Data and theory differ by less than a half of a standard deviation above  $p_T^W$  of 60 GeV/ $c$ . We therefore conclude that the shape of the distribution is consistent with the theoretical prediction.

In summary, we have measured the shape of the transverse momentum distribution of  $W$  bosons produced in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV, and have found that it is consistent with the combined QCD perturbative and resummation calculations.

We thank the staffs at Fermilab and collaborating institutions for their contributions to this work, and acknowledge support from the Department of Energy and National Science Foundation (U.S.A.), Commissariat à l'Énergie Atomique (France), State Committee for Science and Technology and Ministry for Atomic Energy (Russia), CAPES and CNPq (Brazil), Departments of Atomic Energy and Science and Education (India), Colciencias (Colombia), CONACyT (Mexico), Ministry of Education and KOSEF (Korea), and CONICET and UBACyT (Argentina).

## REFERENCES

\* Visitor from Universidad San Francisco de Quito, Quito, Ecuador.

† Visitor from IHEP, Beijing, China.

- [1] J. Collins, D. Soper, and G. Sterman, Nucl. Phys. **B250**, 199 (1985); J. Collins and D. Soper, Nucl. Phys. **B193**, 381 (1981), Nucl. Phys. **B197**, 446 (1982), and erratum Nucl. Phys. **B213**, 545 (1983).
- [2] C. Davies, B. Webber, and W. J. Stirling, Nucl. Phys. **B256**, 413 (1985); C. Davies and W. J. Stirling, Nucl. Phys. **B244**, 337 (1984).
- [3] G. Altarelli, R. K. Ellis, M. Greco, and G. Martinelli, Nucl. Phys. **B246**, 12 (1984).
- [4] P. B. Arnold and R. Kauffman, Nucl. Phys. **B349**, 381 (1991).
- [5] G. A. Ladinsky and C. P. Yuan, Phys. Rev. D **50**, 4239 (1994).
- [6] P. B. Arnold and M. H. Reno, Nucl. Phys. **B319**, 37 (1989); R. J. Gonsalves, J. Pawlowski, and C-F. Wai, Phys. Rev. D **40**, 2245 (1989).
- [7] UA1 Collaboration, C. Albajar *et al.*, Z. Phys. C **44**, 15 (1989).
- [8] UA2 Collaboration, J. Alitti *et al.*, Z. Phys. C **47**, 523 (1990).
- [9] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **66**, 2951 (1991), and Phys. Rev. Lett. **67**, 2937 (1991).
- [10] DØ Collaboration, S. Abachi *et al.*, Nucl. Instrum. Methods A **338**, 185 (1994).
- [11] DØ Collaboration, S. Abachi *et al.*, Phys. Rev. Lett. **75**, 1456 (1995).
- [12] DØ Collaboration, S. Abachi *et al.*, Phys. Rev. D **552**, 4877 (1995).
- [13] DØ Collaboration, B. Abbott *et al.*, Fermilab-Pub-97/328-E, hep-ex/9710007 (October 1997), submitted to Phys. Rev. D and Fermilab-Pub-97/422-E, hep-ex/9712029 (December 1997), submitted to Phys. Rev. D.
- [14] G. D'Agostini, Nucl. Instrum. Methods A **362**, 487 (1995).
- [15] S. Baker and R. D. Cousins, Nucl. Instrum. Methods A **221**, 437 (1984).
- [16] A. D. Martin, R. G. Roberts, and W. J. Stirling, Phys. Rev. D **50**, 6734 (1994); **51**, 4756 (1995).



TABLE I. The  $W$  boson transverse momentum distribution corresponding to Fig. 2. The column labeled “Stat Error” shows the statistical uncertainty; “Syst Error” shows the systematic uncertainty in background and efficiency; “Detector Error” shows the systematic uncertainty in the detector modelling; “Total Error” is the sum in quadrature of the previous three columns.

Bin Width	$\langle p_T^W \rangle$	$N_{\text{signal}}$	$(1/N)(dN/dp_T^W)$	Stat Error	Syst Error	Detector Error	Total Error
(GeV/ $c$ )	(GeV/ $c$ )		( $c$ /TeV)	( $c$ /TeV)	( $c$ /TeV)	( $c$ /TeV)	( $c$ /TeV)
2	1.2	506.8	37.4	1.6	1.9	6.1	6.6
2	3.0	1232.0	90.8	2.6	4.6	7.9	9.5
2	5.0	1253.0	92.4	2.6	4.8	4.4	7.0
2	7.0	1006.6	74.2	2.3	3.9	3.7	5.8
2	9.0	718.4	53.0	1.9	2.8	2.6	4.2
2	11.0	431.2	31.8	1.5	1.7	2.0	3.0
2	13.0	368.4	27.2	1.4	1.4	1.6	2.6
2	15.1	228.0	16.8	1.1	1.1	1.2	1.9
2	17.1	184.4	13.6	1.0	0.8	0.8	1.5
2	19.0	167.9	12.4	0.9	0.7	0.9	1.5
5	22.6	252.0	7.43	0.46	0.42	0.65	0.89
5	27.3	145.4	4.29	0.34	0.30	0.43	0.61
5	32.3	83.0	2.45	0.25	0.22	0.26	0.42
5	37.2	56.7	1.67	0.20	0.19	0.18	0.33
10	44.2	59.4	0.875	0.099	0.147	0.150	0.232
20	57.7	45.2	0.333	0.037	0.144	0.045	0.155
20	78.0	25.2	0.186	0.028	0.066	0.020	0.075
30	100.7	10.7	0.052	0.011	0.035	0.010	0.038
30	133.2	5.1	0.025	0.008	0.009	0.002	0.013
50	172.7	3.6	0.011	0.005	0.002	0.001	0.005




FIG. 1. The transverse mass (left) and  $\cancel{E}_T$  (right) distributions for three  $p_T^W$  bins. The points are the  $D\bar{O}$  data. The solid histogram is the sum of the Monte Carlo signal and the estimated background. The dotted histogram is the estimated background alone.




FIG. 2. The  $W$  boson transverse momentum spectrum, showing the  $D\bar{O}$  result (solid points) with statistical uncertainty. The theoretical calculation by Arnold and Kauffman [4], smeared for detector resolutions, is shown as two lines corresponding to the  $\pm 1\sigma$  variations of the uncertainties in the detector modelling. Within each bin, the values are plotted at the mean  $p_T^W$ . The fractional systematic uncertainty on the data is shown as a band in the lower portion of the plot. The values of the uncertainties for different  $p_T^W$  bins are 100% correlated with each other. Upward fluctuations in the magnitude of the multijet background cause the widening observed in the band at about 60 and 100  $\text{GeV}/c$  in  $p_T^W$ .




FIG. 3. The ratio  $(\text{Data} - \text{Theory}) / \text{Theory}$  shown as a function of  $p_T^W$  with its statistical uncertainty as error bars. Within each bin, the values are plotted at the mean  $p_T^W$ . The theory corresponds to reference [4], smeared for detector resolutions. The systematic uncertainties from data (background and efficiency) and from the detector modelling are added in quadrature and shown as a band.