

Fermi National Accelerator Laboratory

FERMILAB-Pub-98/084-E

CDF

**Search for Chargino-Neutralino Production at the Fermilab
Tevatron Collider**

F. Abe et al.

The CDF Collaboration

*Fermi National Accelerator Laboratory
P.O. Box 500, Batavia, Illinois 60510*

March 1998

Submitted to *Physical Review Letters*

Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Distribution

Approved for public release; further dissemination unlimited.

Search for Chargino-Neutralino Associated Production at the Fermilab Tevatron Collider

F. Abe,¹⁷ H. Akimoto,³⁹ A. Akopian,³¹ M. G. Albrow,⁷ A. Amadon,⁵ S. R. Amendolia,²⁷
 D. Amidei,²⁰ J. Antos,³³ S. Aota,³⁷ G. Apollinari,³¹ T. Arisawa,³⁹ T. Asakawa,³⁷
 W. Ashmanskas,¹⁸ M. Atac,⁷ P. Azzi-Bacchetta,²⁵ N. Bacchetta,²⁵ S. Bagdasarov,³¹
 M. W. Bailey,²² P. de Barbaro,³⁰ A. Barbaro-Galtieri,¹⁸ V. E. Barnes,²⁹ B. A. Barnett,¹⁵
 M. Barone,⁹ G. Bauer,¹⁹ T. Baumann,¹¹ F. Bedeschi,²⁷ S. Behrends,³ S. Belforte,²⁷
 G. Bellettini,²⁷ J. Bellinger,⁴⁰ D. Benjamin,³⁵ J. Bensinger,³ A. Beretvas,⁷ J. P. Berge,⁷
 J. Berryhill,⁵ S. Bertolucci,⁹ S. Bettelli,²⁷ B. Bevensee,²⁶ A. Bhatti,³¹ K. Biery,⁷
 C. Bigongiari,²⁷ M. Binkley,⁷ D. Bisello,²⁵ R. E. Blair,¹ C. Blocker,³ S. Blusk,³⁰
 A. Bodek,³⁰ W. Bokhari,²⁶ G. Bolla,²⁹ Y. Bonushkin,⁴ D. Bortoletto,²⁹ J. Boudreau,²⁸
 L. Breccia,² C. Bromberg,²¹ N. Bruner,²² R. Brunetti,² E. Buckley-Geer,⁷ H. S. Budd,³⁰
 K. Burkett,²⁰ G. Busetto,²⁵ A. Byon-Wagner,⁷ K. L. Byrum,¹ M. Campbell,²⁰ A. Caner,²⁷
 W. Carithers,¹⁸ D. Carlsmith,⁴⁰ J. Cassada,³⁰ A. Castro,²⁵ D. Cauz,³⁶ A. Cerri,²⁷
 P. S. Chang,³³ P. T. Chang,³³ H. Y. Chao,³³ J. Chapman,²⁰ M. -T. Cheng,³³ M. Chertok,³⁴
 G. Chiarelli,²⁷ C. N. Chiou,³³ F. Chlebana,⁷ L. Christofek,¹³ M. L. Chu,³³ S. Cihangir,⁷
 A. G. Clark,¹⁰ M. Cobal,²⁷ E. Cocca,²⁷ M. Contreras,⁵ J. Conway,³² J. Cooper,⁷
 M. Cordelli,⁹ D. Costanzo,²⁷ C. Couyoumtzelis,¹⁰ D. Cronin-Hennessy,⁶ R. Culbertson,⁵
 D. Dagenhart,³⁸ T. Daniels,¹⁹ F. DeJongh,⁷ S. Dell'Agnello,⁹ M. Dell'Orso,²⁷ R. Demina,⁷
 L. Demortier,³¹ M. Deninno,² P. F. Derwent,⁷ T. Devlin,³² J. R. Dittmann,⁶ S. Donati,²⁷
 J. Done,³⁴ T. Dorigo,²⁵ N. Eddy,²⁰ K. Einsweiler,¹⁸ J. E. Elias,⁷ R. Ely,¹⁸ E. Engels, Jr.,²⁸
 W. Erdmann,⁷ D. Errede,¹³ S. Errede,¹³ Q. Fan,³⁰ R. G. Feild,⁴¹ Z. Feng,¹⁵ C. Ferretti,²⁷
 I. Fiori,² B. Flaughner,⁷ G. W. Foster,⁷ M. Franklin,¹¹ J. Freeman,⁷ J. Friedman,¹⁹
 H. Frisch,⁵ Y. Fukui,¹⁷ S. Gadomski,¹⁴ S. Galeotti,²⁷ M. Gallinaro,²⁶ O. Ganel,³⁵
 M. Garcia-Sciveres,¹⁸ A. F. Garfinkel,²⁹ C. Gay,⁴¹ S. Geer,⁷ D. W. Gerdes,¹⁵ P. Giannetti,²⁷
 N. Giokaris,³¹ P. Giromini,⁹ G. Giusti,²⁷ M. Gold,²² A. Gordon,¹¹ A. T. Goshaw,⁶
 Y. Gotra,²⁵ K. Goulianos,³¹ H. Grassmann,³⁶ L. Groer,³² C. Grosso-Pilcher,⁵ G. Guillian,²⁰
 J. Guimaraes da Costa,¹⁵ R. S. Guo,³³ C. Haber,¹⁸ E. Hafen,¹⁹ S. R. Hahn,⁷ R. Hamilton,¹¹
 T. Handa,¹² R. Handler,⁴⁰ F. Happacher,⁹ K. Hara,³⁷ A. D. Hardman,²⁹ R. M. Harris,⁷
 F. Hartmann,¹⁶ J. Hauser,⁴ E. Hayashi,³⁷ J. Heinrich,²⁶ W. Hao,³⁵ B. Hinrichsen,¹⁴
 K. D. Hoffman,²⁹ M. Hohlmann,⁵ C. Holck,²⁶ R. Hollebeek,²⁶ L. Holloway,¹³ Z. Huang,²⁰
 B. T. Huffman,²⁸ R. Hughes,²³ J. Huston,²¹ J. Huth,¹¹ H. Ikeda,³⁷ M. Incagli,²⁷
 J. Incandela,⁷ G. Introzzi,²⁷ J. Iwai,³⁹ Y. Iwata,¹² E. James,²⁰ H. Jensen,⁷ U. Joshi,⁷
 E. Kajfasz,²⁵ H. Kambara,¹⁰ T. Kamon,³⁴ T. Kaneko,³⁷ K. Karr,³⁸ H. Kasha,⁴¹ Y. Kato,²⁴
 T. A. Keaffaber,²⁹ K. Kelley,¹⁹ R. D. Kennedy,⁷ R. Kephart,⁷ D. Kestenbaum,¹¹
 D. Khazins,⁶ T. Kikuchi,³⁷ B. J. Kim,²⁷ H. S. Kim,¹⁴ S. H. Kim,³⁷ Y. K. Kim,¹⁸ L. Kirsch,³
 S. Klimentenko,⁸ D. Knoblauch,¹⁶ P. Koehn,²³ A. Königeter,¹⁶ K. Kondo,³⁷ J. Konigsberg,⁸
 K. Kordas,¹⁴ A. Korytov,⁸ E. Kovacs,¹ W. Kowald,⁶ J. Kroll,²⁶ M. Kruse,³⁰
 S. E. Kuhlmann,¹ E. Kuns,³² K. Kurino,¹² T. Kuwabara,³⁷ A. T. Laasanen,²⁹ I. Nakano,¹²

S. Lami,²⁷ S. Lammel,⁷ J. I. Lamoureux,³ M. Lancaster,¹⁸ M. Lanzoni,²⁷ G. Latino,²⁷
 T. LeCompte,¹ S. Leone,²⁷ J. D. Lewis,⁷ P. Limon,⁷ M. Lindgren,⁴ T. M. Liss,¹³ J. B. Liu,³⁰
 Y. C. Liu,³³ N. Lockyer,²⁶ O. Long,²⁶ C. Loomis,³² M. Loreti,²⁵ D. Lucchesi,²⁷ P. Lukens,⁷
 S. Lusin,⁴⁰ J. Lys,¹⁸ K. Maeshima,⁷ P. Maksimovic,¹⁹ M. Mangano,²⁷ M. Mariotti,²⁵
 J. P. Marriner,⁷ A. Martin,⁴¹ J. A. J. Matthews,²² P. Mazzanti,² P. McIntyre,³⁴ P. Melese,³¹
 M. Menguzzato,²⁵ A. Menzione,²⁷ E. Meschi,²⁷ S. Metzler,²⁶ C. Miao,²⁰ T. Miao,⁷
 G. Michail,¹¹ R. Miller,²¹ H. Minato,³⁷ S. Miscetti,⁹ M. Mishina,¹⁷ S. Miyashita,³⁷
 N. Moggi,²⁷ E. Moore,²² Y. Morita,¹⁷ A. Mukherjee,⁷ T. Muller,¹⁶ P. Murat,²⁷ S. Murgia,²¹
 H. Nakada,³⁷ I. Nakano,¹² C. Nelson,⁷ D. Neuberger,¹⁶ C. Newman-Holmes,⁷
 C.-Y. P. Ngan,¹⁹ L. Nodulman,¹ A. Nomerotski,⁸ S. H. Oh,⁶ T. Ohmoto,¹² T. Ohsugi,¹²
 R. Oishi,³⁷ M. Okabe,³⁷ T. Okusawa,²⁴ J. Olsen,⁴⁰ C. Pagliarone,²⁷ R. Paoletti,²⁷
 V. Papadimitriou,³⁵ S. P. Pappas,⁴¹ N. Parashar,²⁷ A. Parri,⁹ J. Patrick,⁷ G. Pauletta,³⁶
 M. Paulini,¹⁸ A. Perazzo,²⁷ L. Pescara,²⁵ M. D. Peters,¹⁸ T. J. Phillips,⁶ G. Piacentino,²⁷
 M. Pillai,³⁰ K. T. Pitts,⁷ R. Plunkett,⁷ A. Pompos,²⁹ L. Pondrom,⁴⁰ J. Proudfoot,¹
 F. Ptohos,¹¹ G. Punzi,²⁷ K. Ragan,¹⁴ D. Reher,¹⁸ M. Reischl,¹⁶ A. Ribon,²⁵ F. Rimondi,²
 L. Ristori,²⁷ W. J. Robertson,⁶ T. Rodrigo,²⁷ S. Rolli,³⁸ L. Rosenson,¹⁹ R. Roser,¹³
 T. Saab,¹⁴ W. K. Sakumoto,³⁰ D. Saltzberg,⁴ A. Sansoni,⁹ L. Santi,³⁶ H. Sato,³⁷
 P. Schlabach,⁷ E. E. Schmidt,⁷ M. P. Schmidt,⁴¹ A. Scott,⁴ A. Scribano,²⁷ S. Segler,⁷
 S. Seidel,²² Y. Seiya,³⁷ F. Semeria,² T. Shah,¹⁹ M. D. Shapiro,¹⁸ N. M. Shaw,²⁹
 P. F. Shepard,²⁸ T. Shibayama,³⁷ M. Shimojima,³⁷ M. Shochet,⁵ J. Siegrist,¹⁸ A. Sill,³⁵
 P. Sinervo,¹⁴ P. Singh,¹³ K. Sliwa,³⁸ C. Smith,¹⁵ F. D. Snider,¹⁵ J. Spalding,⁷ T. Speer,¹⁰
 P. Sphicas,¹⁹ F. Spinella,²⁷ M. Spiropulu,¹¹ L. Spiegel,⁷ L. Stanco,²⁵ J. Steele,⁴⁰
 A. Stefanini,²⁷ R. Ströhmer,^{7a} J. Strologas,¹³ F. Strumia,¹⁰ D. Stuart,⁷ K. Sumorok,¹⁹
 J. Suzuki,³⁷ T. Suzuki,³⁷ T. Takahashi,²⁴ T. Takano,²⁴ R. Takashima,¹² K. Takikawa,³⁷
 M. Tanaka,³⁷ B. Tannenbaum,²² F. Tartarelli,²⁷ W. Taylor,¹⁴ M. Tecchio,²⁰ P. K. Teng,³³
 Y. Teramoto,²⁴ K. Terashi,³⁷ S. Tether,¹⁹ D. Theriot,⁷ T. L. Thomas,²²
 R. Thurman-Keup,¹ M. Timko,³⁸ P. Tipton,³⁰ A. Titov,³¹ S. Tkaczyk,⁷ D. Toback,⁵
 K. Tollefson,¹⁹ A. Tollestrup,⁷ H. Toyoda,²⁴ W. Trischuk,¹⁴ J. F. de Troconiz,¹¹ S. Truitt,²⁰
 J. Tseng,¹⁹ N. Turini,²⁷ T. Uchida,³⁷ F. Ukegawa,²⁶ J. Valls,³² S. C. van den Brink,²⁸
 S. Vejcik, III,²⁰ G. Velez,²⁷ R. Vidal,⁷ R. Vilar,^{7a} D. Vucinic,¹⁹ R. G. Wagner,¹
 R. L. Wagner,⁷ J. Wahl,⁵ N. B. Wallace,²⁷ A. M. Walsh,³² C. Wang,⁶ C. H. Wang,³³
 M. J. Wang,³³ A. Warburton,¹⁴ T. Watanabe,³⁷ T. Watts,³² R. Webb,³⁴ C. Wei,⁶
 H. Wenzel,¹⁶ W. C. Wester, III,⁷ A. B. Wicklund,¹ E. Wicklund,⁷ R. Wilkinson,²⁶
 H. H. Williams,²⁶ P. Wilson,⁵ B. L. Winer,²³ D. Winn,²⁰ D. Wolinski,²⁰ J. Wolinski,²¹
 S. Worm,²² X. Wu,¹⁰ J. Wyss,²⁷ A. Yagil,⁷ W. Yao,¹⁸ K. Yasuoka,³⁷ G. P. Yeh,⁷ P. Yeh,³³
 J. Yoh,⁷ C. Yosef,²¹ T. Yoshida,²⁴ I. Yu,⁷ A. Zanetti,³⁶ F. Zetti,²⁷ and S. Zucchelli²

(CDF Collaboration)

¹ Argonne National Laboratory, Argonne, Illinois 60439

² Istituto Nazionale di Fisica Nucleare, University of Bologna, I-40127 Bologna, Italy

³ Brandeis University, Waltham, Massachusetts 02254

⁴ University of California at Los Angeles, Los Angeles, California 90024

⁵ University of Chicago, Chicago, Illinois 60637

- ⁶ *Duke University, Durham, North Carolina 27708*
- ⁷ *Fermi National Accelerator Laboratory, Batavia, Illinois 60510*
- ⁸ *University of Florida, Gainesville, FL 32611*
- ⁹ *Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*
- ¹⁰ *University of Geneva, CH-1211 Geneva 4, Switzerland*
- ¹¹ *Harvard University, Cambridge, Massachusetts 02138*
- ¹² *Hiroshima University, Higashi-Hiroshima 724, Japan*
- ¹³ *University of Illinois, Urbana, Illinois 61801*
- ¹⁴ *Institute of Particle Physics, McGill University, Montreal H3A 2T8, and University of Toronto, Toronto M5S 1A7, Canada*
- ¹⁵ *The Johns Hopkins University, Baltimore, Maryland 21218*
- ¹⁶ *Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany*
- ¹⁷ *National Laboratory for High Energy Physics (KEK), Tsukuba, Ibaraki 305, Japan*
- ¹⁸ *Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720*
- ¹⁹ *Massachusetts Institute of Technology, Cambridge, Massachusetts 02139*
- ²⁰ *University of Michigan, Ann Arbor, Michigan 48109*
- ²¹ *Michigan State University, East Lansing, Michigan 48824*
- ²² *University of New Mexico, Albuquerque, New Mexico 87131*
- ²³ *The Ohio State University, Columbus, OH 43210*
- ²⁴ *Osaka City University, Osaka 588, Japan*
- ²⁵ *Università di Padova, Istituto Nazionale di Fisica Nucleare, Sezione di Padova, I-35131 Padova, Italy*
- ²⁶ *University of Pennsylvania, Philadelphia, Pennsylvania 19104*
- ²⁷ *Istituto Nazionale di Fisica Nucleare, University and Scuola Normale Superiore of Pisa, I-56100 Pisa, Italy*
- ²⁸ *University of Pittsburgh, Pittsburgh, Pennsylvania 15260*
- ²⁹ *Purdue University, West Lafayette, Indiana 47907*
- ³⁰ *University of Rochester, Rochester, New York 14627*
- ³¹ *Rockefeller University, New York, New York 10021*
- ³² *Rutgers University, Piscataway, New Jersey 08855*
- ³³ *Academia Sinica, Taipei, Taiwan 11530, Republic of China*
- ³⁴ *Texas A&M University, College Station, Texas 77843*
- ³⁵ *Texas Tech University, Lubbock, Texas 79409*
- ³⁶ *Istituto Nazionale di Fisica Nucleare, University of Trieste/ Udine, Italy*
- ³⁷ *University of Tsukuba, Tsukuba, Ibaraki 315, Japan*
- ³⁸ *Tufts University, Medford, Massachusetts 02155*
- ³⁹ *Waseda University, Tokyo 169, Japan*
- ⁴⁰ *University of Wisconsin, Madison, Wisconsin 53706*
- ⁴¹ *Yale University, New Haven, Connecticut 06520*

(April 3, 1998)

Abstract

We have searched in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV for events with three charged leptons and missing transverse energy. In the Minimal Supersymmetric Standard Model, we expect trilepton events from chargino-neutralino ($\tilde{\chi}_1^\pm \tilde{\chi}_2^0$) pair production, with subsequent decay into leptons. We observe no candidate $e^+e^-e^\pm$, $e^+e^-\mu^\pm$, $e^\pm\mu^+\mu^-$ or $\mu^+\mu^-\mu^\pm$ events in 106 pb^{-1} integrated luminosity. We present limits on the sum of the branching ratios times cross section for the four channels: $\sigma_{\tilde{\chi}_1^\pm \tilde{\chi}_2^0} \cdot \text{BR}(\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow 3\ell + X) < 0.34 \text{ pb}$, $M_{\tilde{\chi}_1^\pm} > 81.5 \text{ GeV}/c^2$ and $M_{\tilde{\chi}_2^0} > 82.2 \text{ GeV}/c^2$ for $\tan\beta = 2$, $\mu = -600 \text{ GeV}/c^2$ and $M_{\tilde{q}} = M_{\tilde{g}}$.

PACS numbers: 14.80.Ly, 12.60.Jv, 13.85.Rm

The Minimal Supersymmetric Standard Model (MSSM) [1] contains two Higgs doublets and supersymmetric partners to all the Standard Model (SM) particles. The superpartners of the electroweak gauge bosons and Higgs bosons are two charged and four neutral fermions ($\tilde{\chi}$'s). Further assumptions, namely the Grand Unified Theory hypothesis provided by Supergravity [2], Supergravity-inspired slepton/sneutrino mass constraints [3], degeneracy of five of the squarks, and R-parity conservation, lead to models with only six parameters. R-parity conservation implies the creation of superpartners in pairs and the stability of the lightest supersymmetric partner (LSP). Within this framework we expect, for certain regions of parameter space, a measurable rate for the reaction $q\bar{q}' \rightarrow \tilde{\chi}_1^\pm \tilde{\chi}_2^0$, where $\tilde{\chi}_1^\pm \rightarrow \tilde{\chi}_1^0 \ell^\pm \nu$ and $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 \ell^+ \ell^-$, and $\tilde{\chi}_1^0$ is the LSP. The ν and two LSPs do not interact with the detector and manifest themselves as missing energy. The resulting final state is three isolated charged leptons plus missing energy [4]. In this Letter, we report on a search for direct production of $\tilde{\chi}_1^\pm \tilde{\chi}_2^0$, via virtual W^\pm s -channel and virtual squark t -channel diagrams, in the trilepton channels e^+e^-e , $e^+e^-\mu$, $e\mu^+\mu^-$ and $\mu^+\mu^-\mu$. Additional trilepton production arising from squark and gluino cascade decays was not included. We add 87 pb $^{-1}$ of data recorded in 1994-95 to a previously analyzed sample of 19 pb $^{-1}$ collected in 1992-93 [5].

The Collider Detector at Fermilab (CDF) is described in detail elsewhere [6]. The components of the detector relevant to this analysis are the vertex chamber, which provides r - z tracking information; the central tracking chamber, which is situated inside a 1.4 T solenoidal magnetic field and provides a combination of r - ϕ , z and transverse momentum (p_T) information for charged particles; the central ($|\eta| < 1.1$) and endplug ($1.1 < |\eta| < 2.4$) electromagnetic calorimeters, which are located outside the solenoid and are segmented in a projective tower geometry; and the central muon chambers. We define pseudorapidity $\eta \equiv -\ln \tan(\theta/2)$ and θ and ϕ to be the polar and azimuthal angles with respect to the beam axis.

We begin with a sample of 87 pb $^{-1}$ recorded in 1994-95, which contains 3.3×10^6 events that have an electron or muon with $p_T > 8$ GeV/c and $|\eta^e| < 1.1$ or $|\eta^\mu| < 0.6$, and an additional charged lepton (excluding taus) with $p_T > 3$ GeV/c and $|\eta^e| < 2.4$ or $|\eta^\mu| < 1.0$. We select events from this sample by requiring an electron with $E_T^e > 11$ GeV and $|\eta^e| < 1.1$ or a muon with $p_T^\mu > 11$ GeV/c and $|\eta^\mu| < 0.6$. We require two additional charged leptons with $E_T^e > 5$ GeV and $|\eta^e| < 2.4$ or $p_T^\mu > 4$ GeV/c and $|\eta^\mu| < 1.0$. At least one lepton passing the high threshold cut must pass stringent lepton identification cuts [5-9]. To improve the integrity of these events, we require that all three leptons originate from a common vertex within 60 cm of the center of the detector. The 60 cm requirement is to preserve the projective tower geometry of the calorimetry. We find 59 events meeting these requirements.

The principal backgrounds to the $\tilde{\chi}_1^\pm \tilde{\chi}_2^0$ signature are real trilepton events from $W^\pm Z^0$, $Z^0 Z^0$, $t\bar{t}$ and $b\bar{b}$ and dilepton plus fake lepton [10] events from W^+W^- , Z^0 and the Drell-Yan process. To remove background from $b\bar{b}$, $c\bar{c}$ and $t\bar{t}$ production and fake leptons, each lepton must be isolated, where isolation is defined by requiring less than 2 GeV E_T in the calorimeter inside an η - ϕ cone of radius $\Delta R \equiv \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2} = 0.4$ around the lepton, excluding the energy deposited by the lepton. There must be at least one e^+e^- or $\mu^+\mu^-$ pair, the η - ϕ distance between any two leptons ($\Delta R_{\ell\ell}$) must be greater than 0.4 (to remove background from $b\bar{b}$ production, as well as some anomalously reconstructed cosmic

ray events) and the difference in azimuthal angle between the two highest p_T leptons ($\Delta\phi_{\ell_1\ell_2}$) in the event must be less than 170° (to remove background from the Drell-Yan process and cosmic rays) [9]. Events containing a same flavor $\ell^+\ell^-$ pair with invariant mass in the regions of the resonances J/ψ (2.9-3.3 GeV/ c^2), Υ (9-11 GeV/ c^2) and Z^0 (75-105 GeV/ c^2) are removed. These requirements select 6 events (see Table I). In the previous data sample [5] these criteria selected zero events.

The presence of two LSPs and a neutrino in the final state of the signal can lead to substantial missing transverse energy (\cancel{E}_T). The dominant remaining backgrounds, $b\bar{b}$ production and the Drell-Yan process, do not have significant \cancel{E}_T . As seen in Table I, requiring $\cancel{E}_T > 15$ GeV reduces the background by 85% while retaining 82% of the expected signal for $M_{\tilde{\chi}_1^\pm} \approx 70$ GeV/ c^2 . No events pass the \cancel{E}_T cut.

For the remainder of the analysis we combine this data with the previous 19 ± 1 pb $^{-1}$ sample [5] for a total Run I integrated luminosity ($\int \mathcal{L} dt$) of 106 ± 7 pb $^{-1}$ and zero candidate trilepton events.

To determine the SM background and the signal acceptance, we use the ISAJET Monte Carlo program [11] with a CDF detector simulation. For background due to vector boson pair production we use theoretical calculations of cross sections and branching ratios [12]. For background due to $t\bar{t}$ production and the Drell-Yan process we use cross sections measured by CDF [7]. The rate of lepton misidentification was determined from a $W^\pm \rightarrow \ell\nu$ data sample to be $(0.29 \pm 0.04)\%$ per event. After all cuts are applied the total expected background is 1.2 ± 0.2 events in 106 pb $^{-1}$.

The total detection efficiency (ϵ^{tot}) is a product of the trigger efficiency, the isolation efficiency, the lepton identification efficiency and a geometric and kinematic acceptance factor. The triggers used were single lepton and dilepton triggers, with single lepton trigger efficiencies of $\epsilon_e^{trig} = (87_{-5}^{+4})\%$ above 11 GeV and $\epsilon_\mu^{trig} = (87 \pm 3)\%$ above 11 GeV/ c . We determine the lepton isolation and identification efficiencies by studying the second lepton in $Z^0 \rightarrow \ell^+\ell^-$ events. The isolation efficiency is $(90 \pm 4)\%$ per lepton. The per-event trilepton identification efficiency ranges from $(59 \pm 1)\%$ to $(82 \pm 1)\%$, depending on the combination of leptons in the event. The geometric and kinematic acceptance is determined using ISAJET and the CDF detector simulation. The total efficiency (ϵ^{tot}) is mainly a function of the $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ masses, which are nearly equal for the region of the search. The efficiency increases linearly from 3% at 50 GeV/ c^2 to 12% at 100 GeV/ c^2 , because massive $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ lead to more central and more energetic leptons which are detected with higher efficiency.

We see no signal candidates and thus set limits on the available parameter space. A particular point in parameter space is excluded if the predicted number of events exceeds the number of events (s) expected at the 95% confidence level limit given that zero events were observed. The predicted number of events is a function of the cross section times branching ratio ($\sigma(pp \rightarrow \tilde{\chi}_1^\pm \tilde{\chi}_2^0 + X) \cdot \text{BR}(\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow 3\ell + X)$) and $\epsilon^{tot} \cdot \int \mathcal{L} dt$. We calculate cross section times branching ratio ($\sigma \cdot \text{BR}$) using ISAJET 7.20 with CTEQ-3L [13] parton distribution functions and calculate s by convolving the total systematic uncertainty as a Gaussian smearing with a Poisson distribution. The total systematic uncertainty is 15%, which includes uncertainty in the total integrated luminosity ($\pm 7\%$), the parton distribution ($\pm 7\%$), the trigger efficiency ($\pm 6\%$), and the trilepton-finding efficiency ($\pm 2\%$), leading to $s = 3.1$. To calculate the uncertainty due to the parton distribution function we compare CTEQ-3L with a variety of other parton distribution functions. We use the largest uncertainty in the efficiency of any

single lepton trigger for all events.

Using the model assumptions listed in the first paragraph, four parameters determine the $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ masses, production cross sections and decay branching ratios: the ratio of the Higgs vacuum expectation values ($\tan\beta$), the Higgsino mass parameter (μ), the gluino mass ($M_{\tilde{g}}$) and the squark-to-gluino mass ratio ($M_{\tilde{q}}/M_{\tilde{g}}$). To make the analysis more independent of details of the Higgs sector, we consider a region in the MSSM where there is no significant chargino or neutralino branching fraction into Higgs particles. Technically, we do this by choosing the mass of the pseudoscalar Higgs (M_A) to be above the chargino/neutralino mass and use $M_A = 500 \text{ GeV}/c^2$. The production and decay of $\tilde{\chi}_1^\pm\tilde{\chi}_2^0$ are independent of the remaining MSSM parameter, the trilinear top squark coupling (A_t). We fix $A_t = \mu/\tan\beta$ for consistency with other CDF analyses [14]. The search is more sensitive at low values of $\tan\beta$; $\tan\beta \gtrsim 10$ leads to higher branching ratios to τ 's, for which we do not search. We consider $1.1 \leq \tan\beta \leq 8$. We use $-1000 \text{ GeV}/c^2 < \mu < -200 \text{ GeV}/c^2$ because the search is more sensitive to negative values of μ and $|\mu|$ is expected to be on the order of the energy scale at which supersymmetric phenomena should be observable. Also, small $|\mu|$ increases the Higgsino content of the $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$, which decreases the branching ratio to leptons. ISAJET requires $M_{\tilde{g}}$ and $M_{\tilde{q}}$ as input parameters to calculate $M_{\tilde{\chi}_1^\pm}$. The slepton/sneutrino mass constraints [3] use $M_{\tilde{g}}$ and $M_{\tilde{q}}$ to determine $M_{\tilde{\ell}}$ and $M_{\tilde{\nu}}$; large differences in $M_{\tilde{g}}$ and $M_{\tilde{q}}$ lead to heavy sleptons and sneutrinos and decreases the branching ratio to leptons. Thus, this analysis considers $M_{\tilde{q}}/M_{\tilde{g}} > 1$ to avoid invisible decays through light sneutrinos and $M_{\tilde{q}}/M_{\tilde{g}} < 2$ to enhance leptonic final states. For the regions of parameter space we examine $M_{\tilde{g}} \approx 3M_{\tilde{\chi}_1^\pm}$, so we use $150 \text{ GeV}/c^2 \leq M_{\tilde{g}} \leq 340 \text{ GeV}/c^2$.

Figure 1 shows the limit for a few representative points in the $M_{\tilde{\chi}_1^\pm} - (\sigma \cdot \text{BR})$ plane. All points above the solid line are excluded. For comparison, we include the result of the DØ collaboration [15]. DØ reports the *average* $\sigma \cdot \text{BR}$; we use the sum. Figure 2 shows the limit on $M_{\tilde{\chi}_1^\pm}$ as a function of μ and $\tan\beta$. These limits are compared to the limits from ALEPH [16] in Figure 2. The ALEPH result is from a search for all possible final states. The OPAL, L3 and DELPHI collaborations report similar limits [17].

We also examined a string-inspired $SU(5) \times U(1)$ one-parameter supergravity model [18]. This model requires $M_{\tilde{\chi}_1^\pm} \lesssim 87 \text{ GeV}/c^2$ and $M_{\tilde{\chi}_2^0} \lesssim 91 \text{ GeV}/c^2$ and has a nearly maximized trilepton branching ratio via $\tilde{\chi}_2^0 \rightarrow \tilde{\ell}_R \ell$ and $\tilde{\ell}_R \rightarrow \ell \tilde{\chi}_1^0$. As shown in Figure 3, we exclude most of this model and set $M_{\tilde{\chi}_1^\pm} > 80.5 \text{ GeV}/c^2$, $M_{\tilde{\chi}_2^0} > 86.7 \text{ GeV}/c^2$ and $\sigma \cdot \text{BR}(\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow 3\ell + X) < 0.48 \text{ pb}$.

In conclusion, we find no evidence for $\tilde{\chi}_1^\pm \tilde{\chi}_2^0$ production in 1.8 TeV $p\bar{p}$ collisions and set limits on $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ masses and $\sigma \cdot \text{BR}$ within the framework of MSSM models which have $M_{\tilde{\chi}_1^\pm} \approx M_{\tilde{\chi}_2^0} \approx 2M_{\tilde{\chi}_1^0}$ and three-body decays of $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$. The strongest limit is $\sigma_{\tilde{\chi}_1^\pm \tilde{\chi}_2^0} \cdot \text{BR}(\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow 3\ell + X) < 0.34 \text{ pb}$, $M_{\tilde{\chi}_1^\pm} > 81.5 \text{ GeV}/c^2$ and $M_{\tilde{\chi}_2^0} > 82.2 \text{ GeV}/c^2$ for $\tan\beta = 2$, $\mu = -600 \text{ GeV}/c^2$ and $M_{\tilde{q}} = M_{\tilde{g}}$.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Science and Culture of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the A. P. Sloan Foundation.

REFERENCES

- [1] H.P. Nilles, Phys. Rep. **110**, 1 (1984); H.E. Haber and G.L. Kane, Phys. Rep. **117**, 75 (1985).
- [2] A.H. Chamseddine *et al.*, Phys. Rev. Lett. **49**, 970 (1982); R. Barbieri *et al.*, Phys. Lett. **B119**, 343 (1982); L. Hall *et al.*, Phys. Rev. **D27**, 2359 (1983). For a review, see R. Arnowitt and P. Nath, “Supersymmetry and Supergravity,” VIIth Swieca Summer School, Campos de Jordao, Brazil, (World Scientific, Singapore, 1994).
- [3] L.E. Ibañez *et al.*, Nucl. Phys. **B256**, 218 (1985); G.G. Ross and R.G. Roberts, Nucl. Phys. **B377**, 571 (1992); R. Arnowitt and P. Nath, Phys. Rev. Lett. **69**, 725 (1992); S. Kelley *et al.*, Nucl. Phys. **B398**, 31 (1993); G.L. Kane *et al.*, Phys. Rev. **D49**, 6173 (1994). In this analysis we use formulae in H. Baer *et al.*, Phys. Rev. **D47**, 2739 (1993).
- [4] P. Nath and R. Arnowitt, Mod. Phys. Lett. **A2**, 331 (1987); R. Barbieri *et al.*, Nucl. Phys. **B367**, 28 (1991); J.L. Lopez *et al.*, Phys. Rev. **D48**, 2062 (1993); H. Baer and X. Tata, Phys. Rev. **D48**, 5175 (1993).
- [5] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **76**, 4307 (1996).
- [6] F. Abe *et al.* (CDF Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **271**, 387 (1988); F. Abe *et al.*, Phys. Rev. **D50**, 2966 (1994).
- [7] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **74**, 2626 (1994); F. Abe *et al.* (CDF Collaboration), Phys. Rev. **D49**, 1 (1994).
- [8] F. Abe *et al.* (CDF Collaboration), Phys. Rev. **D52**, 4784 (1995).
- [9] B. Tannenbaum, Ph. D. dissertation, University of New Mexico, NMCPP 97/12 (1997).
- [10] “Fake lepton” includes both non-prompt leptons such as decay-in-flight muons and non-leptonic objects passing the lepton identification cuts.
- [11] H. Baer *et al.*, “Simulating Supersymmetry with ISAJET 7.0/ISASUSY 1.0,” Proc. of Workshop on Physics at Current Accelerators and the Supercollider, (Argonne Nat. Lab., 1993). We use ISAJET V7.20.
- [12] J. Ohnemus, Phys. Rev. **D 44**, 1403 (1991) and Phys. Rev. **D44**, 3477 (1991); J. Ohnemus and J.F. Owens, Phys. Rev. **D 43**, 3626 (1991).
- [13] H.L. Lai *et al.* (CTEQ Collaboration), Phys. Rev. **D51**, 4763 (1995).
- [14] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **76**, 2006 (1996); F. Abe *et al.* (CDF Collaboration), Phys. Rev. **D56**, 1357 (1997).
- [15] B. Abbott *et al.* (DØ Collaboration), Phys. Rev. Lett. **80**, 1591 (1998).
- [16] R. Barate *et al.* (ALEPH Collaboration), CERN-PPE-97-128 (1997).
- [17] K. Ackerstaff *et al.* (OPAL Collaboration), CERN-PPE-97-083 (1997); M. Acciarri *et al.* (L3 Collaboration), CERN-PPE/97-130 (1997); P. Abreu *et al.* (DELPHI Collaboration), Eur. Phys. J. C1, 1 (1998).
- [18] J. Lopez *et al.*, Phys. Rev. **D52**, 4178 (1995) and Phys. Rev. **D53**, 5253 (1996).

TABLES

TABLE I. Events remaining after each cut in the 1994-95 data (87 pb^{-1}). One Z^0 event and one J/ψ event are removed with the resonance cuts. For comparison we indicate the expected background (BG) and an expected signal from a representative MSSM Monte Carlo (MC) sample ($M_{\tilde{q}} = M_{\tilde{g}} = 200 \text{ GeV}/c^2$, $\tan \beta = 2$, $\mu = -400 \text{ GeV}/c^2$, $M_{\tilde{\chi}_1^\pm} \simeq M_{\tilde{\chi}_2^0} \simeq 70 \text{ GeV}/c^2$, $\sigma_{\tilde{\chi}_1^\pm \tilde{\chi}_2^0} = 4.8 \text{ pb}$, $\epsilon^{\text{tot}} = 6.7\%$).

Cut	Events	Expected BG	MSSM MC
Dilepton data set	3,270,488		
Trilepton data set	59		
◦ Isolation $< 2 \text{ GeV}$	23		
◦ Require e^+e^- or $\mu^+\mu^-$	23		
◦ $\Delta R_{\ell\ell} > 0.4$	9		
◦ $\Delta\phi_{\ell_1\ell_2} < 170^\circ$	8	9.6 ± 1.5	6.2 ± 0.6
◦ J/ψ , Υ , Z^0 removal	6	6.6 ± 1.1	5.5 ± 0.5
◦ $\cancel{E}_T > 15 \text{ GeV}$	0	1.0 ± 0.2	4.5 ± 0.4
Total Run I data (106 pb^{-1})	0	1.2 ± 0.2	5.5 ± 0.4

FIGURES




FIG. 1. $\sigma \cdot \text{BR}(\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow 3\ell + X)$ versus $\tilde{\chi}_1^\pm$ mass for representative points in the MSSM parameter space, namely $\mu = -400 \text{ GeV}/c^2$, $\tan \beta = 2$ and (a) $M_{\tilde{q}}/M_{\tilde{g}} = 2.0$, (b) $M_{\tilde{q}}/M_{\tilde{g}} = 1.5$, (c) $M_{\tilde{q}}/M_{\tilde{g}} = 1.2$ and (d) $M_{\tilde{q}}/M_{\tilde{g}} = 1.0$. BR is the summed branching ratio into the four trilepton modes (e^+e^-e , $e^+e^-\mu$, $e\mu^+\mu^-$ and $\mu^+\mu^-\mu$). The solid line is the 95% confidence level upper limit based on an observation of zero events. We set a mass limit of $77.0 \text{ GeV}/c^2$ when $M_{\tilde{q}}=M_{\tilde{g}}$. All MSSM points in this plot yield three body decays of the $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$. The $D\emptyset$ limit [15] is for single trilepton mode which we scale up by 4 to match our notation.




FIG. 2. The experimental limit on $M_{\tilde{\chi}_1^\pm}$ as a function of μ for $\tan\beta = 2$ and $M_{\tilde{q}} = M_{\tilde{g}}$ (upper) and as a function of $\tan\beta$ for $\mu = -400 \text{ GeV}/c^2$ and $M_{\tilde{q}} = M_{\tilde{g}}$ (lower). The ALEPH limits shown [16] are from a search for all possible final states. In this analysis $M_{\tilde{\nu}} \approx 100 \text{ GeV}/c^2$. Minimal SUGRA models favor the region of small $|\mu|$ values.




FIG. 3. $\sigma \cdot \text{BR}(\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow 3\ell + X)$ versus $\tilde{\chi}_1^\pm$ mass for the $SU(5) \times U(1)$ model [18]. This sets a mass limit of $80.5 \text{ GeV}/c^2$. In this model, the $\tilde{\ell}_R$ is lighter than the $\tilde{\chi}_2^0$, resulting in two body decays of the $\tilde{\chi}_2^0$. Note that the acceptance for events from this model decreases at large $\tilde{\chi}_1^\pm$ mass. In this region, the LSP mass approaches that of the $\tilde{\ell}$, resulting in soft final state leptons which are difficult to detect.