

Fermi National Accelerator Laboratory

FERMILAB-Conf-99/167-E

D0

**Inclusive Jet Cross Sections in
 $\bar{p}p$ Collisions at $\sqrt{s} = 630$ and 1800 GeV**

V. Daniel Elvira

For the D0 Collaboration

*State University of New York at Stony Brook
Stony Brook, New York 11794-3800*

*Fermi National Accelerator Laboratory
P.O. Box 500, Batavia, Illinois 60510*

September 1999

Published Proceedings of the *International Workshop on Deep Inelastic Scattering and QCD*,
Zeuthen, Germany, April 19-23, 1999

Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Distribution

Approved for public release; further dissemination unlimited.

Copyright Notification

This manuscript has been authored by Universities Research Association, Inc. under contract No. DE-AC02-76CH03000 with the U.S. Department of Energy. The United States Government and the publisher, by accepting the article for publication, acknowledges that the United States Government retains a nonexclusive, paid-up, irrevocable, worldwide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government Purposes.

Inclusive Jet Cross Sections in $\bar{p}p$ Collisions at $\sqrt{s}= 630$ and 1800 GeV

V. Daniel Elvira^{a *}

^aDepartment of Physics and Astronomy, State University of New York at Stony Brook, Stony Brook, NY 11794-3800

We have made a precise measurement of the inclusive jet cross section at $\sqrt{s}=1800$ GeV. The result is based on an integrated luminosity of 92 pb^{-1} collected at the Fermilab Tevatron $\bar{p}p$ Collider with the DØ detector. The measurement is reported as a function of jet transverse energy ($60 \text{ GeV} \leq E_T \leq 500 \text{ GeV}$), and in the pseudorapidity intervals $|\eta| \leq 0.5$ and $0.1 \leq |\eta| \leq 0.7$. A preliminary measurement of the pseudorapidity dependence of inclusive jet production ($|\eta| \leq 1.5$) is also discussed. The results are in good agreement with predictions from next-to-leading order (NLO) quantum chromodynamics (QCD). DØ has also determined the ratio of jet cross sections at $\sqrt{s}=630$ GeV and $\sqrt{s}=1800$ GeV ($|\eta| \leq 0.5$). This preliminary measurement differs from NLO QCD predictions.

1. Introduction

Within the framework of quantum chromodynamics (QCD), inelastic scattering between a proton and antiproton is described as a hard collision between their constituents (partons). After the collision, the outgoing partons manifest themselves as localized streams of particles or “jets”. Predictions for the inclusive jet cross section have improved in the early nineties with next-to-leading order (NLO) perturbative QCD calculations [1] and new, accurately measured parton density functions (pdf)[2].

DØ has recently measured and published [3] the cross section for the production of jets as a function of the jet energy transverse to the incident beams, E_T . The measurement is based on an integrated luminosity of about 92 pb^{-1} of $\bar{p}p$ hard collisions collected with the DØ Detector [4] at the Fermilab Tevatron Collider. This result allows a stringent test of QCD, with a total uncertainty substantially reduced relative to previous results [5,6]. We also measure the ratio of jet cross sections at two center-of-mass energies: 630 (based on an integrated luminosity of about 0.537 pb^{-1}) and 1800 GeV [7]. Experimental and theoretical uncertainties are significantly reduced in the ratio. This is due to the large correlation in the errors of the two cross section measure-

ments, and the suppression of the sensitivity to parton distribution functions (pdf) in the prediction. The ratio of cross sections thus provides a stronger test of the matrix element portion of the calculation than a single cross section measurement alone. Previous measurements of cross section ratios have been performed with smaller data sets by the UA2 and CDF [8] experiments.

2. Jet Reconstruction and Data Selection

Jets are reconstructed using an iterative jet cone algorithm with a cone radius of $\mathcal{R}=0.7$ in η - ϕ space, (pseudorapidity is defined as $\eta = -\ln[\tan(\frac{\theta}{2})]$) [9]. The offline data selection procedure, which eliminates background caused by electrons, photons, noise, or cosmic rays, follows the methods described in Refs. [10,11].

3. Energy Corrections

The jet energy scale correction, described in [12], removes instrumentation effects associated with calorimeter response, showering, and noise, as well as the contribution from spectator partons (underlying event).

The energy scale corrects jets from their reconstructed E_T to their “true” E_T on average. An unsmearing correction is applied later to remove the effect of a finite E_T resolution [3].

*Representing the DØ Collaboration

4. The Inclusive Jet Cross Section

The resulting inclusive double differential jet cross sections, $\langle d^2\sigma/(dE_T d\eta) \rangle$, for $|\eta| \leq 0.5$ and $0.1 \leq |\eta| \leq 0.7$ (the second region for comparison to Ref. [6]), are compared with a NLO QCD theoretical prediction [1]. Discussions on the different choices in the theoretical calculation: pdfs, renormalization and factorization scales (μ), and clustering algorithm parameter (R_{sep}) can be found in Refs. [9].

Figure 1 shows the ratios $(D - T)/T$ for the data (D) and JETRAD NLO theoretical (T) predictions based on the CTEQ3M, CTEQ4M and MRST pdf's [4,5] for $|\eta| \leq 0.5$. (The tabulated data for both $|\eta| \leq 0.5$ and $0.1 \leq |\eta| \leq 0.7$ measurements can be found in Ref. [13].)

The predictions are in good quantitative agreement with the data, as verified with a $\chi^2 = \sum_{i,j} (D_i - T_i)(C^{-1})_{ij}(D_j - T_j)$ test, which incorporates the uncertainty covariance matrix C . Here D_i and T_i represent the i -th data and theory points, respectively. The overall systematic uncertainty is largely correlated.

Table 1 lists χ^2 values for several JETRAD predictions using various parton distribution functions [2]. The predictions describe both the $|\eta| \leq 0.5$ and $0.1 \leq |\eta| \leq 0.7$ cross section very well. The measurement by DØ and CDF are also in good quantitative agreement within their systematic uncertainties [3].

Table 1

χ^2 comparisons between JETRAD and $|\eta| \leq 0.5$ and $0.1 \leq |\eta| \leq 0.7$ data for $\mu = 0.5E_T^{\max}$, $R_{sep}=1.3R$, and various pdfs. There are 24 degrees of freedom.

pdf	$ \eta \leq 0.5$	$0.1 \leq \eta \leq 0.7$
CTEQ3M	23.9	28.4
CTEQ4M	17.6	23.3
CTEQ4HJ	15.7	20.5
MRSA'	20.0	27.8
MRST	17.0	19.5

5. η Dependence of $\langle d^2\sigma/(dE_T d\eta) \rangle$

DØ has made a preliminary measurement of the pseudorapidity dependence of the inclusive jet cross section. Figure 2 shows the ratios $(D - T)/T$ for the data (D) and JETRAD NLO theoretical




Figure 1. The difference between data and JETRAD QCD predictions normalized to predictions. The bands are the total experimental uncertainty.

(T) predictions using the CTEQ3M pdf set for $0.5 \leq |\eta| \leq 1$ and $1 \leq |\eta| \leq 1.5$. The measurements and the predictions are in good qualitative agreement. A detailed error analysis is currently being completed.

6. Ratio of Scale Invariant Jet Cross Sections

A simple parton model would predict a jet cross section that scales with center-of-mass energy. In this scenario, $E_T^4 \cdot E \frac{d^3\sigma}{dp^3}$, plotted as a function of jet $x_T \equiv \frac{2E_T}{\sqrt{s}}$, would remain constant with respect to the center-of-mass energy. Figure 3 shows the DØ measurement of $E_T^4 \cdot E \frac{d^3\sigma}{dp^3}$ (stars) compared to JETRAD predictions (lines). There is poor agreement between data and NLO QCD calculations using the same μ in the numerator and the denominator (probability of agreement not greater than 10%). The agreement improves for predictions with different μ at the two center-of-mass energies [7].

In conclusion, we have made precise measurements of jet production cross sections. At $\sqrt{s}=1800$ GeV, there is good agreement between the measurements and the data. The ratio of cross sections at $\sqrt{s}=1800$ and 630 GeV, however, differs from NLO QCD predictions, unless different renormalization scales are introduced for the two center-of-mass energies.




Figure 2. Pseudorapidity dependence of the inclusive jet cross section ($0.5 < |\eta| < 1$ and $1 < |\eta| < 1.5$). Comparison between data and NLO QCD predictions. The bands are the total systematic uncertainty in the experiment.

REFERENCES

- W. T. Giele, E. W. N. Glover, and D. A. Kosower, Phys. Rev. Letters **73**, 2019 (1994); S. D. Ellis, Z. Kunszt, and D. E. Soper, Phys. Rev. Letters **64**, 2121 (1990); F. Aversa *et al.*, Phys. Rev. Letters **65**, (1990)
- H. L. Lai *et al.*, Phys. Rev. D **51**, 4763 (1995); A. D. Martin *et al.*, Eur. Phys. J.C **4**, 463-496 (1998).
- B. Abbott *et al.*, (DØ Collaboration), Phys. Rev. Letters **82**, 2451 (1999).
- S. Abachi *et al.*, (DØ Collaboration), Nucl. Instrum. Methods **A338**, 185 (1994).
- J. Alitti *et al.*, (UA2 Collaboration), Phys. Letters B **257**, 232 (1991); Z.Phys.C **49**, 17 (1991); F. Abe *et al.*, (CDF Collaboration), Phys. Rev. Letters **70**, 1376 (1993).
- F. Abe *et al.*, (CDF Collaboration), Phys. Rev. Letters **77**, 438 (1996).
- J. Krane (DØ Collaboration). To be published in the proceedings of DPF (1999).
- J. Appel *et al.*, (UA2 Collaboration), Phys.




Figure 3. The ratio of scale invariant jet cross sections. The stars are the DØ data, the band is the systematic uncertainty, and the lines are the NLO QCD predictions.

- Letters B **160**, 349 (1985); F. Abe *et al.*, (CDF Collaboration), Phys. Rev. Lett. **70**, 1376 (1993)
- S. Abachi *et al.*, (DØ Collaboration), Phys. Letters B **357**, 500 (1995); B. Abbott *et al.*, Eur. Phys. J. C. **5**, 687 (1998).
 - D. Elvira, Ph.D. Thesis, Universidad de Buenos Aires, Buenos Aires, Argentina, 1995 (unpublished), [HTTP://WWW-D0.FNAL.GOV/RESULTS/PUBLICATIONS_TALKS/THESIS/THESIS.HTML](http://www-d0.fnal.gov/results/publications_talks/thesis/thesis.html).
 - J. Krane, Ph.D. Thesis, University of Nebraska-Lincoln, 1998 (unpublished), [HTTP://WWW-D0.FNAL.GOV/RESULTS/PUBLICATIONS_TALKS/THESIS/THESIS.HTML](http://www-d0.fnal.gov/results/publications_talks/thesis/thesis.html).
 - B. Abbott *et al.*, (DØ Collaboration), Nuc. Inst. and Meth. A **424**, nos. 2-3, 352-394 (1999)
 - Available from the AIP E-PAPS service and at [HTTP://WWW-D0.FNAL.GOV/ D0QCD/INCLUSIVE](http://www-d0.fnal.gov/d0qcd/inclusive)