

International Journal of Modern Physics A  
 © World Scientific Publishing Company

**Measurements of the  $t\bar{t}$  production cross section at the Tevatron  
 Run II CDF experiment using b-tagging**

Henri Bachacou

*Lawrence Berkeley National Laboratory  
 1 cyclotron road, Bldg 50B-5239  
 Berkeley CA 94720*

Received (Day Month Year)  
 Revised (Day Month Year)

We present measurements of the  $t\bar{t}$  production cross section in b-tagged lepton + jets events from  $p\bar{p}$  collisions at 1.96 TeV using the CDF detector at Fermilab. B-jets are tagged with either a secondary vertex algorithm, or a soft lepton tagger that identifies muons from B hadron semileptonic decays. With Tevatron Run II data, we estimate the  $t\bar{t}$  signal fraction in two different ways: by estimating the various background contributions, and by fitting directly the leading jet transverse energy spectrum for the signal and background contributions. A subset of the sample, with two secondary vertex tagged jets, yields a production cross section consistent with the inclusive measurements. Results are consistent with a Standard Model  $t\bar{t}$  signal and current measurements of the top quark mass.

*Keywords:* top; CDF; Tevatron

## 1. Introduction

The Tevatron (Run II) collides protons and anti-protons head-on at a center-of-mass energy of 1.96 TeV. In such collisions, the Standard Model (SM) predicts a  $t\bar{t}$  production cross section of  $\sigma_{t\bar{t}} = 6.7^{+0.7}_{-0.9} \text{ pb}$  at  $m_t = 175 \text{ GeV}/c^2$ <sup>1</sup>. Top quarks are expected to decay almost exclusively to a W boson and a b quark. When one W decays leptonically, the  $t\bar{t}$  event contains a high transverse momentum lepton, missing energy from the unrecorded neutrino, and 4 high transverse momentum jets, 2 of which originate from b quarks. We use this decay channel to measure the total  $t\bar{t}$  production cross section. A deviation from the predicted value would be an indication of new physics either in the production mechanism or in the top decay. We select events with an isolated electron  $E_T$  (muon  $P_T$ ) greater than 20 GeV, missing  $E_T > 20 \text{ GeV}$  and at least 3 jets with  $E_T > 15 \text{ GeV}$  and  $|\eta| < 2.0$ . Finally, we require at least one jet in the event to be identified as a heavy flavor jet, either using a secondary vertex algorithm (SECVTX), or a soft lepton tagger (SLT) that identifies muons from B hadron semileptonic decays. The analyses using SECVTX (resp. SLT) are based on  $162 \text{ pb}^{-1}$  (resp.  $194 \text{ pb}^{-1}$ ) of data. The CDF detector is described in detail elsewhere<sup>2</sup>.




Fig. 1. Jet multiplicity of  $W+jets$  events tagged with the SecVtx algorithm in  $162 \text{ pb}^{-1}$  of data. The  $H_T > 200 \text{ GeV}$  cut is only applied to events with three or more jets.




Fig. 2.  $H_T$  distribution of the candidate events compared to the expected backgrounds and  $t\bar{t}$  signal.

## 2. Measurement with secondary vertex b-tagging.

We optimize the event selection by requiring that the total transverse energy in the event ( $H_T$ , the scalar sum of all jets  $E_T$ , lepton  $p_T$ , and missing  $E_T$ ) be larger than 200 GeV. The SECVTX algorithm selects tracks within the jet with large impact parameter to reconstruct secondary vertices. Jets containing a secondary vertex more than  $3\sigma$  away from the primary vertex (in the plane transverse to the beam) are identified as b-jets. After tuning the simulation on a control sample, the efficiency for tagging at least one jet in a  $t\bar{t}$  event that passes all other selection requirements is  $(53 \pm 4)\%$ . The main sources of background are  $W + \text{Heavy Flavor}$  events,  $W + \text{light jets}$  events where one jet is wrongly tagged, and QCD events that fake a  $W$  signal; they are estimated with techniques that use both Monte Carlo and data control samples. We expect  $13.8 \pm 2.0$  background events and observe 48 events in the data; we measure a cross section of  $5.6^{+1.2}_{-1.1}(\text{stat.})^{+1.0}_{-0.7}(\text{syst.}) \text{ pb}$ . Fig. 1 shows the number of candidate events vs jet multiplicity together with the expected background contributions. Fig. 2 shows the  $H_T$  variable distribution of the candidates compared to the expected background and  $t\bar{t}$  signal (normalized to 6.7 pb).

The sub-sample of events with at least two tagged jets contains 8 events, compared to an expected background of  $0.9 \pm 0.2$  events, from which we measure a cross section of  $5.4 \pm 2.2(\text{stat.}) \pm 1.1(\text{syst.}) \text{ pb}$ .

## 3. Measurement with SECVTX using a kinematic fit.

Instead of explicitly evaluating the contribution to the sample from backgrounds, one can extract the  $t\bar{t}$  fraction by fitting some kinematic variable in the data. The leading jet  $E_T$  variable was chosen for this purpose. Template shapes for the background are evaluated from the data; the template shape for  $t\bar{t}$  is from Monte Carlo.




Fig. 3. Leading jet transverse energy of candidates in  $162 \text{ pb}^{-1}$  of data, together with fitted contribution from  $t\bar{t}$  signal and background.




Fig. 4. Jet multiplicity of  $W+\text{jets}$  events tagged with the SLT algorithm in  $162 \text{ pb}^{-1}$  of data.

The fit (Fig. 3) measures a  $t\bar{t}$  fraction of  $(67^{+13}_{-16})\%$ , leading to a cross section of  $6.0^{+1.5}_{-1.8}(\text{stat.}) \pm 0.8(\text{syst.}) \text{ pb}$ .

#### 4. Measurement with soft muon b-tagging.

The muon SLT algorithm matches tracks in the central drift chamber with segments in the muon chambers. It uses a global  $\chi^2$  built from the matching distributions, to define a pseudo-likelihood variable,  $L$ , that separates muon candidates from background. A jet is considered "tagged" if it contains an SLT muon with  $P_T > 3 \text{ GeV}/c$ , with  $L < 3.5$  and within  $\Delta R < 0.6$  of the jet axis. Efficiency and fake rate are measured on control samples. Backgrounds are estimated with techniques similar to Sec. 2. We expect  $11.6 \pm 1.5$  background events, and observe 20, and we measure a cross section of  $4.2^{+2.9}_{-1.9}(\text{stat.}) \pm 1.4(\text{syst.}) \text{ pb}$ . Fig. 4 shows the jet multiplicity of the candidates compared to the expected background.

#### References

1. M. Cacciari, et al., *JHEP* **404**, 68 (2004).
2. F. Abe, et al., *Nucl. Instrum. Methods Phys. Res. A* **271**, 387 (1988); D. Amidei, et al., *Nucl. Instrum. Methods Phys. Res. A* **350**, 73 (1994); F. Abe, et al., *Phys. Rev. D* **52**, 4784 (1995); P. Azzi, et al., *Nucl. Instrum. Methods Phys. Res. A* **360**, 137 (1995); *The CDFII Detector Technical Design Report*, Fermilab-Pub-96/390-E