

## W boson polarization measurement in the $t\bar{t}$ dilepton channel using the CDF II Detector

T. Aaltonen,<sup>21</sup> B. Álvarez González<sup>z,9</sup> S. Amerio,<sup>40</sup> D. Amidei,<sup>32</sup> A. Anastassov<sup>x,15</sup> A. Annovi,<sup>17</sup> J. Antos,<sup>12</sup> G. Apollinari,<sup>15</sup> J.A. Appel,<sup>15</sup> T. Arisawa,<sup>54</sup> A. Artikov,<sup>13</sup> J. Asaadi,<sup>49</sup> W. Ashmanskas,<sup>15</sup> B. Auerbach,<sup>57</sup> A. Aurisano,<sup>49</sup> F. Azfar,<sup>39</sup> W. Badgett,<sup>15</sup> T. Bae,<sup>25</sup> A. Barbaro-Galtieri,<sup>26</sup> V.E. Barnes,<sup>44</sup> B.A. Barnett,<sup>23</sup> P. Barria<sup>hh,42</sup> P. Bartos,<sup>12</sup> M. Bauce<sup>ff,40</sup> F. Bedeschi,<sup>42</sup> S. Behari,<sup>23</sup> G. Bellettini<sup>gg,42</sup> J. Bellinger,<sup>56</sup> D. Benjamin,<sup>14</sup> A. Beretvas,<sup>15</sup> A. Bhatti,<sup>46</sup> D. Bisello<sup>ff,40</sup> I. Bizjak,<sup>28</sup> K.R. Bland,<sup>5</sup> B. Blumenfeld,<sup>23</sup> A. Bocci,<sup>14</sup> A. Bodek,<sup>45</sup> D. Bortoletto,<sup>44</sup> J. Boudreau,<sup>43</sup> A. Boveia,<sup>11</sup> L. Brigliadori<sup>ee,6</sup> C. Bromberg,<sup>33</sup> E. Brucken,<sup>21</sup> J. Budagov,<sup>13</sup> H.S. Budd,<sup>45</sup> K. Burkett,<sup>15</sup> G. Busetto<sup>ff,40</sup> P. Bussey,<sup>19</sup> A. Buzatu,<sup>31</sup> A. Calamba,<sup>10</sup> C. Calancha,<sup>29</sup> S. Camarda,<sup>4</sup> M. Campanelli,<sup>28</sup> M. Campbell,<sup>32</sup> F. Canelli,<sup>11,15</sup> B. Carls,<sup>22</sup> D. Carlsmith,<sup>56</sup> R. Carosi,<sup>42</sup> S. Carrillo<sup>m,16</sup> S. Carron,<sup>15</sup> B. Casal<sup>k,9</sup> M. Casarsa,<sup>50</sup> A. Castro<sup>ee,6</sup> P. Catastini,<sup>20</sup> D. Cauz,<sup>50</sup> V. Cavaliere,<sup>22</sup> M. Cavalli-Sforza,<sup>4</sup> A. Cerri<sup>f,26</sup> L. Cerrito<sup>s,28</sup> Y.C. Chen,<sup>1</sup> M. Chertok,<sup>7</sup> G. Chiarelli,<sup>42</sup> G. Chlachidze,<sup>15</sup> F. Chlebana,<sup>15</sup> K. Cho,<sup>25</sup> D. Chokheli,<sup>13</sup> W.H. Chung,<sup>56</sup> Y.S. Chung,<sup>45</sup> M.A. Ciocci<sup>hh,42</sup> A. Clark,<sup>18</sup> C. Clarke,<sup>55</sup> G. Compostella<sup>ff,40</sup> M.E. Convery,<sup>15</sup> J. Conway,<sup>7</sup> M. Corbo,<sup>15</sup> M. Cordelli,<sup>17</sup> C.A. Cox,<sup>7</sup> D.J. Cox,<sup>7</sup> F. Crescioli<sup>gg,42</sup> J. Cuevas<sup>z,9</sup> R. Culbertson,<sup>15</sup> D. Dagenhart,<sup>15</sup> N. d'Ascenzo<sup>w,15</sup> M. Datta,<sup>15</sup> P. de Barbaro,<sup>45</sup> M. Dell'Orso<sup>gg,42</sup> L. Demortier,<sup>46</sup> M. Deninno,<sup>6</sup> F. Devoto,<sup>21</sup> M. d'Errico<sup>ff,40</sup> A. Di Canto<sup>gg,42</sup> B. Di Ruzza,<sup>15</sup> J.R. Dittmann,<sup>5</sup> M. D'Onofrio,<sup>27</sup> S. Donati<sup>gg,42</sup> P. Dong,<sup>15</sup> M. Dorigo,<sup>50</sup> T. Dorigo,<sup>40</sup> K. Ebina,<sup>54</sup> A. Elagin,<sup>49</sup> A. Eppig,<sup>32</sup> R. Erbacher,<sup>7</sup> S. Errede,<sup>22</sup> N. Ershaidat<sup>dd,15</sup> R. Eusebi,<sup>49</sup> S. Farrington,<sup>39</sup> M. Feindt,<sup>24</sup> J.P. Fernandez,<sup>29</sup> R. Field,<sup>16</sup> G. Flanagan<sup>u,15</sup> R. Forrest,<sup>7</sup> M.J. Frank,<sup>5</sup> M. Franklin,<sup>20</sup> J.C. Freeman,<sup>15</sup> Y. Funakoshi,<sup>54</sup> I. Furic,<sup>16</sup> M. Gallinaro,<sup>46</sup> J.E. Garcia,<sup>18</sup> A.F. Garfinkel,<sup>44</sup> P. Garosi<sup>hh,42</sup> H. Gerberich,<sup>22</sup> E. Gerchtein,<sup>15</sup> S. Giagu,<sup>47</sup> V. Giakoumopoulou,<sup>3</sup> P. Giannetti,<sup>42</sup> K. Gibson,<sup>43</sup> C.M. Ginsburg,<sup>15</sup> N. Giokaris,<sup>3</sup> P. Giromini,<sup>17</sup> G. Giurgiu,<sup>23</sup> V. Glagolev,<sup>13</sup> D. Glenzinski,<sup>15</sup> M. Gold,<sup>35</sup> D. Goldin,<sup>49</sup> N. Goldschmidt,<sup>16</sup> A. Golossanov,<sup>15</sup> G. Gomez,<sup>9</sup> G. Gomez-Ceballos,<sup>30</sup> M. Goncharov,<sup>30</sup> O. González,<sup>29</sup> I. Gorelov,<sup>35</sup> A.T. Goshaw,<sup>14</sup> K. Goulianos,<sup>46</sup> S. Grinstein,<sup>4</sup> C. Grosso-Pilcher,<sup>11</sup> R.C. Group<sup>53,15</sup> J. Guimaraes da Costa,<sup>20</sup> S.R. Hahn,<sup>15</sup> E. Halkiadakis,<sup>48</sup> A. Hamaguchi,<sup>38</sup> J.Y. Han,<sup>45</sup> F. Happacher,<sup>17</sup> K. Hara,<sup>51</sup> D. Hare,<sup>48</sup> M. Hare,<sup>52</sup> R.F. Harr,<sup>55</sup> K. Hatakeyama,<sup>5</sup> C. Hays,<sup>39</sup> M. Heck,<sup>24</sup> J. Heinrich,<sup>41</sup> M. Herndon,<sup>56</sup> S. Hewamanage,<sup>5</sup> A. Hocker,<sup>15</sup> W. Hopkins<sup>g,15</sup> D. Horn,<sup>24</sup> S. Hou,<sup>1</sup> R.E. Hughes,<sup>36</sup> M. Hurwitz,<sup>11</sup> U. Husemann,<sup>57</sup> N. Hussain,<sup>31</sup> M. Hussein,<sup>33</sup> J. Huston,<sup>33</sup> G. Introzzi,<sup>42</sup> M. Iori<sup>jj,47</sup> A. Ivanov<sup>p,7</sup> E. James,<sup>15</sup> D. Jang,<sup>10</sup> B. Jayatilaka,<sup>14</sup> E.J. Jeon,<sup>25</sup> S. Jindariani,<sup>15</sup> M. Jones,<sup>44</sup> K.K. Joo,<sup>25</sup> S.Y. Jun,<sup>10</sup> T.R. Junk,<sup>15</sup> T. Kamon<sup>25,49</sup> P.E. Karchin,<sup>55</sup> A. Kasmi,<sup>5</sup> Y. Kato<sup>o,38</sup> W. Ketchum,<sup>11</sup> J. Keung,<sup>41</sup> V. Khotilovich,<sup>49</sup> B. Kilminster,<sup>15</sup> D.H. Kim,<sup>25</sup> H.S. Kim,<sup>25</sup> J.E. Kim,<sup>25</sup> M.J. Kim,<sup>17</sup> S.B. Kim,<sup>25</sup> S.H. Kim,<sup>51</sup> Y.K. Kim,<sup>11</sup> Y.J. Kim,<sup>25</sup> N. Kimura,<sup>54</sup> M. Kirby,<sup>15</sup> S. Klimenko,<sup>16</sup> K. Knoepfel,<sup>15</sup> K. Kondo<sup>\*,54</sup> D.J. Kong,<sup>25</sup> J. Konigsberg,<sup>16</sup> A.V. Kotwal,<sup>14</sup> M. Kreps,<sup>24</sup> J. Kroll,<sup>41</sup> D. Krop,<sup>11</sup> M. Kruse,<sup>14</sup> V. Krutelyov<sup>c,49</sup> T. Kuhr,<sup>24</sup> M. Kurata,<sup>51</sup> S. Kwang,<sup>11</sup> A.T. Laasanen,<sup>44</sup> S. Lami,<sup>42</sup> S. Lammel,<sup>15</sup> M. Lancaster,<sup>28</sup> R.L. Lander,<sup>7</sup> K. Lannon<sup>y,36</sup> A. Lath,<sup>48</sup> G. Latino<sup>hh,42</sup> T. LeCompte,<sup>2</sup> E. Lee,<sup>49</sup> H.S. Lee<sup>q,11</sup> J.S. Lee,<sup>25</sup> S.W. Lee<sup>bb,49</sup> S. Leo<sup>gg,42</sup> S. Leone,<sup>42</sup> J.D. Lewis,<sup>15</sup> A. Limosani<sup>t,14</sup> C.-J. Lin,<sup>26</sup> M. Lindgren,<sup>15</sup> E. Lipeles,<sup>41</sup> A. Lister,<sup>18</sup> D.O. Litvintsev,<sup>15</sup> C. Liu,<sup>43</sup> H. Liu,<sup>53</sup> Q. Liu,<sup>44</sup> T. Liu,<sup>15</sup> S. Lockwitz,<sup>57</sup> A. Loginov,<sup>57</sup> D. Lucchesi<sup>ff,40</sup> J. Lueck,<sup>24</sup> P. Lujan,<sup>26</sup> P. Lukens,<sup>15</sup> G. Lungu,<sup>46</sup> J. Lys,<sup>26</sup> R. Lysak<sup>e,12</sup> R. Madrak,<sup>15</sup> K. Maeshima,<sup>15</sup> P. Maestro<sup>hh,42</sup> S. Malik,<sup>46</sup> G. Manca<sup>a,27</sup> A. Manousakis-Katsikakis,<sup>3</sup> F. Margaroli,<sup>47</sup> C. Marino,<sup>24</sup> M. Martínez,<sup>4</sup> P. Mastrandrea,<sup>47</sup> K. Matera,<sup>22</sup> M.E. Mattson,<sup>55</sup> A. Mazzacane,<sup>15</sup> P. Mazzanti,<sup>6</sup> K.S. McFarland,<sup>45</sup> P. McIntyre,<sup>49</sup> R. McNulty<sup>j,27</sup> A. Mehta,<sup>27</sup> P. Mehtala,<sup>21</sup> C. Mesropian,<sup>46</sup> T. Miao,<sup>15</sup> D. Mietlicki,<sup>32</sup> A. Mitra,<sup>1</sup> H. Miyake,<sup>51</sup> S. Moed,<sup>15</sup> N. Moggi,<sup>6</sup> M.N. Mondragon<sup>m,15</sup> C.S. Moon,<sup>25</sup> R. Moore,<sup>15</sup> M.J. Morello<sup>ii,42</sup> J. Morlock,<sup>24</sup> P. Movilla Fernandez,<sup>15</sup> A. Mukherjee,<sup>15</sup> Th. Muller,<sup>24</sup> P. Murat,<sup>15</sup> M. Mussini<sup>ee,6</sup> J. Nachtman<sup>n,15</sup> Y. Nagai,<sup>51</sup> J. Naganoma,<sup>54</sup> I. Nakano,<sup>37</sup> A. Napier,<sup>52</sup> J. Nett,<sup>49</sup> C. Neu,<sup>53</sup> M.S. Neubauer,<sup>22</sup> J. Nielsen<sup>d,26</sup> L. Nodulman,<sup>2</sup> S.Y. Noh,<sup>25</sup> O. Norriella,<sup>22</sup> L. Oakes,<sup>39</sup> S.H. Oh,<sup>14</sup> Y.D. Oh,<sup>25</sup> I. Oksuzian,<sup>53</sup> T. Okusawa,<sup>38</sup> R. Orava,<sup>21</sup> L. Ortolan,<sup>4</sup> S. Pagan Griso<sup>ff,40</sup> C. Pagliarone,<sup>50</sup> E. Palencia<sup>f,9</sup> V. Papadimitriou,<sup>15</sup> A.A. Paramonov,<sup>2</sup> J. Patrick,<sup>15</sup> G. Pauletta<sup>kk,50</sup> M. Paulini,<sup>10</sup> C. Paus,<sup>30</sup> D.E. Pellett,<sup>7</sup> A. Penzo,<sup>50</sup> T.J. Phillips,<sup>14</sup> G. Piacentino,<sup>42</sup> E. Pianori,<sup>41</sup> J. Pilot,<sup>36</sup> K. Pitts,<sup>22</sup> C. Plager,<sup>8</sup> L. Pondrom,<sup>56</sup> S. Poprocki<sup>g,15</sup> K. Potamianos,<sup>44</sup> F. Prokoshin<sup>cc,13</sup> A. Pranko,<sup>26</sup> F. Ptohos<sup>h,17</sup> G. Punzi<sup>gg,42</sup> A. Rahaman,<sup>43</sup> V. Ramakrishnan,<sup>56</sup> N. Ranjan,<sup>44</sup> I. Redondo,<sup>29</sup> P. Renton,<sup>39</sup> M. Rescigno,<sup>47</sup> T. Riddick,<sup>28</sup> F. Rimondi<sup>ee,6</sup> L. Ristori<sup>42,15</sup> A. Robson,<sup>19</sup> T. Rodrigo,<sup>9</sup> T. Rodriguez,<sup>41</sup> E. Rogers,<sup>22</sup> S. Rolli<sup>i,52</sup> R. Roser,<sup>15</sup> F. Ruffini<sup>hh,42</sup> A. Ruiz,<sup>9</sup> J. Russ,<sup>10</sup> V. Rusu,<sup>15</sup> A. Safonov,<sup>49</sup> W.K. Sakumoto,<sup>45</sup> Y. Sakurai,<sup>54</sup> L. Santi<sup>kk,50</sup> K. Sato,<sup>51</sup>

V. Saveliev<sup>w,15</sup> A. Savoy-Navarro<sup>aa,15</sup> P. Schlabach,<sup>15</sup> A. Schmidt,<sup>24</sup> E.E. Schmidt,<sup>15</sup> T. Schwarz,<sup>15</sup> L. Scodellaro,<sup>9</sup> A. Scribano<sup>hh,42</sup> F. Scuri,<sup>42</sup> S. Seidel,<sup>35</sup> Y. Seiya,<sup>38</sup> A. Semenov,<sup>13</sup> F. Sforza<sup>hh,42</sup> S.Z. Shalhout,<sup>7</sup> T. Shears,<sup>27</sup> P.F. Shepard,<sup>43</sup> M. Shimojima<sup>v,51</sup> M. Shochet,<sup>11</sup> I. Shreyber-Tecker,<sup>34</sup> A. Simonenko,<sup>13</sup> P. Sinervo,<sup>31</sup> K. Sliwa,<sup>52</sup> J.R. Smith,<sup>7</sup> F.D. Snider,<sup>15</sup> A. Soha,<sup>15</sup> V. Sorin,<sup>4</sup> H. Song,<sup>43</sup> P. Squillacioti<sup>hh,42</sup> M. Stancari,<sup>15</sup> R. St. Denis,<sup>19</sup> B. Stelzer,<sup>31</sup> O. Stelzer-Chilton,<sup>31</sup> D. Stentz<sup>x,15</sup> J. Strologas,<sup>35</sup> G.L. Strycker,<sup>32</sup> Y. Sudo,<sup>51</sup> A. Sukhanov,<sup>15</sup> I. Suslov,<sup>13</sup> K. Takemasa,<sup>51</sup> Y. Takeuchi,<sup>51</sup> J. Tang,<sup>11</sup> M. Tecchio,<sup>32</sup> P.K. Teng,<sup>1</sup> J. Thom<sup>g,15</sup> J. Thome,<sup>10</sup> G.A. Thompson,<sup>22</sup> E. Thomson,<sup>41</sup> D. Toback,<sup>49</sup> S. Tokar,<sup>12</sup> K. Tollefson,<sup>33</sup> T. Tomura,<sup>51</sup> D. Tonelli,<sup>15</sup> S. Torre,<sup>17</sup> D. Torretta,<sup>15</sup> P. Totaro,<sup>40</sup> M. Trovato<sup>ii,42</sup> F. Ukegawa,<sup>51</sup> S. Uozumi,<sup>25</sup> A. Varganov,<sup>32</sup> F. Vázquez<sup>m,16</sup> G. Velev,<sup>15</sup> C. Vellidis,<sup>15</sup> M. Vidal,<sup>44</sup> I. Vila,<sup>9</sup> R. Vilar,<sup>9</sup> J. Vizán,<sup>9</sup> M. Vogel,<sup>35</sup> G. Volpi,<sup>17</sup> P. Wagner,<sup>41</sup> R.L. Wagner,<sup>15</sup> T. Wakisaka,<sup>38</sup> R. Wallny,<sup>8</sup> S.M. Wang,<sup>1</sup> A. Warburton,<sup>31</sup> D. Waters,<sup>28</sup> W.C. Wester III,<sup>15</sup> D. Whiteson<sup>b,41</sup> A.B. Wicklund,<sup>2</sup> E. Wicklund,<sup>15</sup> S. Wilbur,<sup>11</sup> F. Wick,<sup>24</sup> H.H. Williams,<sup>41</sup> J.S. Wilson,<sup>36</sup> P. Wilson,<sup>15</sup> B.L. Winer,<sup>36</sup> P. Wittich<sup>g,15</sup> S. Wolbers,<sup>15</sup> H. Wolfe,<sup>36</sup> T. Wright,<sup>32</sup> X. Wu,<sup>18</sup> Z. Wu,<sup>5</sup> K. Yamamoto,<sup>38</sup> D. Yamato,<sup>38</sup> T. Yang,<sup>15</sup> U.K. Yang<sup>r,11</sup> Y.C. Yang,<sup>25</sup> W.-M. Yao,<sup>26</sup> G.P. Yeh,<sup>15</sup> K. Yi<sup>n,15</sup> J. Yoh,<sup>15</sup> K. Yorita,<sup>54</sup> T. Yoshida<sup>l,38</sup> G.B. Yu,<sup>14</sup> I. Yu,<sup>25</sup> S.S. Yu,<sup>15</sup> J.C. Yun,<sup>15</sup> A. Zanetti,<sup>50</sup> Y. Zeng,<sup>14</sup> C. Zhou,<sup>14</sup> and S. Zucchelli<sup>ee6</sup>  
(CDF Collaboration<sup>†</sup>)

### The CDF collaboration

<sup>1</sup>*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*

<sup>2</sup>*Argonne National Laboratory, Argonne, Illinois 60439, USA*

<sup>3</sup>*University of Athens, 157 71 Athens, Greece*

<sup>4</sup>*Institut de Física d'Altes Energies, ICREA, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*

<sup>5</sup>*Baylor University, Waco, Texas 76798, USA*

<sup>6</sup>*Istituto Nazionale di Fisica Nucleare Bologna, <sup>ee</sup>University of Bologna, I-40127 Bologna, Italy*

<sup>7</sup>*University of California, Davis, Davis, California 95616, USA*

<sup>8</sup>*University of California, Los Angeles, Los Angeles, California 90024, USA*

<sup>9</sup>*Instituto de Física de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*

<sup>10</sup>*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*

<sup>11</sup>*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*

<sup>12</sup>*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*

<sup>13</sup>*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*

<sup>14</sup>*Duke University, Durham, North Carolina 27708, USA*

<sup>15</sup>*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*

<sup>16</sup>*University of Florida, Gainesville, Florida 32611, USA*

<sup>17</sup>*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*

<sup>18</sup>*University of Geneva, CH-1211 Geneva 4, Switzerland*

<sup>19</sup>*Glasgow University, Glasgow G12 8QQ, United Kingdom*

<sup>20</sup>*Harvard University, Cambridge, Massachusetts 02138, USA*

<sup>21</sup>*Division of High Energy Physics, Department of Physics,*

*University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland*

<sup>22</sup>*University of Illinois, Urbana, Illinois 61801, USA*

<sup>23</sup>*The Johns Hopkins University, Baltimore, Maryland 21218, USA*

<sup>24</sup>*Institut für Experimentelle Kernphysik, Karlsruhe Institute of Technology, D-76131 Karlsruhe, Germany*

<sup>25</sup>*Center for High Energy Physics: Kyungpook National University,*

*Daegu 702-701, Korea; Seoul National University, Seoul 151-742,*

*Korea; Sungkyunkwan University, Suwon 440-746,*

*Korea; Korea Institute of Science and Technology Information,*

*Daejeon 305-806, Korea; Chonnam National University, Gwangju 500-757,*

*Korea; Chonbuk National University, Jeonju 561-756, Korea*

<sup>26</sup>*Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*

<sup>27</sup>*University of Liverpool, Liverpool L69 7ZE, United Kingdom*

<sup>28</sup>*University College London, London WC1E 6BT, United Kingdom*

<sup>29</sup>*Centro de Investigaciones Energeticas Medioambientales y Tecnológicas, E-28040 Madrid, Spain*

<sup>30</sup>*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

<sup>31</sup>*Institute of Particle Physics: McGill University, Montréal, Québec,*

*Canada H3A 2T8; Simon Fraser University, Burnaby, British Columbia,*

*Canada V5A 1S6; University of Toronto, Toronto, Ontario,*

*Canada M5S 1A7; and TRIUMF, Vancouver, British Columbia, Canada V6T 2A3*

<sup>32</sup>*University of Michigan, Ann Arbor, Michigan 48109, USA*

<sup>33</sup>*Michigan State University, East Lansing, Michigan 48824, USA*

<sup>34</sup>*Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia*

<sup>35</sup>*University of New Mexico, Albuquerque, New Mexico 87131, USA*

<sup>36</sup>*The Ohio State University, Columbus, Ohio 43210, USA*

<sup>37</sup>*Okayama University, Okayama 700-8530, Japan*

<sup>38</sup>*Osaka City University, Osaka 588, Japan*

<sup>39</sup>*University of Oxford, Oxford OX1 3RH, United Kingdom*

<sup>40</sup>*Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, <sup>fj</sup>University of Padova, I-35131 Padova, Italy*

<sup>41</sup>*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*

<sup>42</sup>*Istituto Nazionale di Fisica Nucleare Pisa, <sup>gg</sup>University of Pisa,*

<sup>hh</sup>*University of Siena and <sup>ii</sup>Scuola Normale Superiore, I-56127 Pisa, Italy*

<sup>43</sup>*University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA*

<sup>44</sup>*Purdue University, West Lafayette, Indiana 47907, USA*

<sup>45</sup>*University of Rochester, Rochester, New York 14627, USA*

<sup>46</sup>*The Rockefeller University, New York, New York 10065, USA*

<sup>47</sup>*Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1,*

<sup>jj</sup>*Sapienza Università di Roma, I-00185 Roma, Italy*

<sup>48</sup>*Rutgers University, Piscataway, New Jersey 08855, USA*

<sup>49</sup>*Texas A&M University, College Station, Texas 77843, USA*

<sup>50</sup>*Istituto Nazionale di Fisica Nucleare Trieste/Udine,*

*I-34100 Trieste, <sup>kk</sup>University of Udine, I-33100 Udine, Italy*

<sup>51</sup>*University of Tsukuba, Tsukuba, Ibaraki 305, Japan*

<sup>52</sup>*Tufts University, Medford, Massachusetts 02155, USA*

<sup>53</sup>*University of Virginia, Charlottesville, Virginia 22906, USA*

<sup>54</sup>*Waseda University, Tokyo 169, Japan*

<sup>55</sup>*Wayne State University, Detroit, Michigan 48201, USA*

<sup>56</sup>*University of Wisconsin, Madison, Wisconsin 53706, USA*

<sup>57</sup>*Yale University, New Haven, Connecticut 06520, USA*

(Dated: May 1, 2012)

We present a measurement of  $W$  boson polarization in top-quark decays in  $t\bar{t}$  events with decays to dilepton final states using  $5.1 \text{ fb}^{-1}$  integrated luminosity in  $p\bar{p}$  collisions collected by the CDF II detector at the Tevatron. A simultaneous measurement of the fractions of longitudinal ( $f_0$ ) and right-handed ( $f_+$ )  $W$  bosons yields the results  $f_0 = 0.71_{-0.17}^{+0.18}(\text{stat}) \pm 0.06(\text{syst})$  and  $f_+ = -0.07 \pm 0.09(\text{stat}) \pm 0.03(\text{syst})$ . Combining this measurement with our previous result based on single lepton final states, we obtain  $f_0 = 0.84 \pm 0.09(\text{stat}) \pm 0.05(\text{syst})$  and  $f_+ = -0.16 \pm 0.05(\text{stat}) \pm 0.04(\text{syst})$ . The results are consistent with standard model expectation.

PACS numbers: 14.65.Ha,12.15.Ji,13.38.Be,13.88.+e

---

\*Deceased

<sup>†</sup>With visitors from <sup>a</sup>Istituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy, <sup>b</sup>University of CA Irvine, Irvine, CA 92697, USA, <sup>c</sup>University of CA Santa Barbara, Santa Barbara, CA 93106, USA, <sup>d</sup>University of CA Santa Cruz, Santa Cruz, CA 95064, USA, <sup>e</sup>Institute of Physics, Academy of Sciences of the Czech Republic, Czech Republic, <sup>f</sup>CERN, CH-1211 Geneva, Switzerland, <sup>g</sup>Cornell University, Ithaca, NY 14853, USA, <sup>h</sup>University of Cyprus, Nicosia CY-1678, Cyprus, <sup>i</sup>Office of Science, U.S. Department of Energy, Washington, DC 20585, USA, <sup>j</sup>University College Dublin, Dublin 4, Ireland, <sup>k</sup>ETH, 8092 Zurich, Switzerland, <sup>l</sup>University of Fukui, Fukui City, Fukui Prefecture, Japan 910-0017, <sup>m</sup>Universidad Iberoamericana, Mexico D.F., Mexico, <sup>n</sup>University of Iowa, Iowa City, IA 52242, USA, <sup>o</sup>Kinki University, Higashi-Osaka City, Japan 577-8502, <sup>p</sup>Kansas State University, Manhattan, KS 66506, USA, <sup>q</sup>Korea University, Seoul, 136-713, Korea, <sup>r</sup>University of Manchester, Manchester M13 9PL, United Kingdom, <sup>s</sup>Queen Mary, University of London, London, E1 4NS, United Kingdom, <sup>t</sup>University of Melbourne, Victoria 3010, Australia, <sup>u</sup>Muons, Inc., Batavia, IL 60510, USA, <sup>v</sup>Nagasaki Institute of Applied Science, Nagasaki, Japan, <sup>w</sup>National Research Nuclear University, Moscow, Russia, <sup>x</sup>Northwestern University, Evanston, IL 60208, USA, <sup>y</sup>University of Notre Dame, Notre Dame,

Since the top quark discovery by the CDF and D0 experiments in 1995, many of its properties have been measured. Due to its very short lifetime, the top quark does not hadronize and therefore its properties are transferred directly to its decay products. The standard model (SM) makes specific predictions for the  $W$ -boson polarization in top-quark decays. Precise measurement of the  $W$ -boson polarization fractions provides a test of the SM and could reveal new physics beyond the SM [1].

In the SM, the top quark decays to a  $W$  boson and  $b$  quark with almost 100% probability [2]. The  $W$  boson is a massive vector particle with three polarization states: right-handed (+1), longitudinal (0), and left-handed (–

---

IN 46556, USA, <sup>z</sup>Universidad de Oviedo, E-33007 Oviedo, Spain, <sup>aa</sup>CNRS-IN2P3, Paris, F-75205 France, <sup>bb</sup>Texas Tech University, Lubbock, TX 79609, USA, <sup>cc</sup>Universidad Tecnica Federico Santa Maria, 110v Valparaiso, Chile, <sup>dd</sup>Yarmouk University, Irbid 211-63, Jordan,

1). The differential decay rate is given by:

$$\frac{d\Gamma}{d\cos\theta^*} \propto (1-c)^2 f_- + 2(1-c^2)f_0 + (1+c)^2 f_+ \quad (1)$$

where  $c \equiv \cos\theta^*$ , and the polarization angle  $\theta^*$  is the angle between the direction of the charged lepton (or down-type quark) and the opposite of the direction of the top quark in the  $W$  boson rest frame. The polarization fractions satisfy the normalization condition  $f_- + f_0 + f_+ = 1$ . Right-handed  $W$  boson production is strongly suppressed due to the  $V-A$  structure of the charged-current weak interaction. In the SM at tree level [1], the fraction of right-handed  $W$  bosons is very close to zero ( $f_+ = 3.7 \times 10^{-4}$ ) while  $f_0 = 0.698$ , and  $f_- = 0.301$  for a top-quark mass of  $173.3 \text{ GeV}/c^2$  [3],  $W$ -boson mass of  $80.4 \text{ GeV}/c^2$  [2], and a  $b$ -quark mass of  $4.78 \text{ GeV}/c^2$  [2]. In the  $m_b \rightarrow 0$  limit,  $f_+ = 0$  and  $f_0 = m_{\text{top}}^2 / (2m_W^2 + m_{\text{top}}^2)$ . Polarization fractions that deviate from the SM are predicted in theories with anomalous  $tWb$  couplings [1].

Earlier measurements of the polarization fractions of the  $W$  boson in top-quark decay by the CDF and D0 collaborations focused on the single-lepton channel [4, 5]. Currently, the most precise result has been reported by the D0 collaboration, where the combination of single lepton and the dilepton channels has been performed [5]. In this Letter we report the CDF measurement in the dilepton channel ( $t\bar{t} \rightarrow W^+bW^- \bar{b} \rightarrow \ell^+ \nu b \ell^- \bar{\nu} \bar{b}$ ). We perform two types of measurements: a model-independent approach where  $f_0$  and  $f_+$  are determined simultaneously; and a model-dependent approach where  $f_0$  ( $f_+$ ) is fixed to its SM value, and  $f_+$  ( $f_0$ ) is measured. The model-independent and model-dependent approaches are referred as “2D” and “1D”, respectively, throughout this article. We also combine this result with our previous measurement [4] in the single-lepton channel.

This analysis is based on data corresponding to an integrated luminosity of  $5.1 \text{ fb}^{-1}$  collected with the CDF II detector [6] between March 2002 and June 2009 at the Fermilab Tevatron with a center of mass energy  $\sqrt{s} = 1.96 \text{ TeV}$ . The CDF II detector is described in detail elsewhere [6]. The components essential to this analysis are the tracking system consisting of a silicon microstrip tracker and a central drift chamber immersed in a 1.4 T solenoidal magnetic field; electromagnetic and hadronic calorimeters arranged in a projective geometry outside the magnet coil; and drift chambers and scintillation counters for muon detection outside the calorimeters.

The analysis uses the same event selection criteria that are used for the measurement of the  $t\bar{t}$  cross-section in the dilepton channel [7]. A brief description of the event selection is as follows: the data are collected with an on-line inclusive event-selection system (trigger) that requires a high transverse energy  $E_T$  [8] lepton (electron or muon with  $E_T > 18 \text{ GeV}$  [9]). From the inclusive lepton data, we selected events with opposite-charged lep-

tons of  $E_T > 20 \text{ GeV}$ . We require the first lepton to be well identified and isolated, while the second lepton is more loosely identified and has no isolation requirement. The pseudorapidity  $\eta$  [8] coverage is  $|\eta| < 2.0$  for electrons and  $|\eta| < 1.0$  for muons. We require missing transverse energy  $\cancel{E}_T > 25 \text{ GeV}$  [10] unless the  $\cancel{E}_T$  direction is along (within  $20^\circ$  in  $\phi$ ) either a lepton or a jet, in which case we require  $\cancel{E}_T > 50 \text{ GeV}$ . Additionally we require at least two jets [11] reconstructed with  $E_T > 15 \text{ GeV}$  and  $|\eta| < 2.5$ . Jet energies are corrected for the effects of calorimeter response, multiple interactions, and the hadronic calorimeter energy scale [12]. Backgrounds to the dilepton signal are further reduced through kinematic cuts on the dilepton invariant mass, total energy in the transverse plane and  $\cancel{E}_T$  significance. For the purpose of this measurement, we split the inclusive data sample into two non-overlapping subsamples (“ $b$ -tag” and “non-tag”) where for the “ $b$ -tag” subsample we require at least one jet in the event to be consistent with having originated from a  $b$  quark by using an algorithm that identifies a long-lived  $b$  hadron through the presence of a displaced vertex [13]. The two subsamples have different signal to background ratios and background compositions; therefore, we can improve the overall measurement uncertainties by analyzing each subsample separately.

The dominant background to  $t\bar{t}$  dilepton events is from “fake” events where a jet is misidentified as a lepton. The main source of “fake” events are  $W(\rightarrow \ell\nu) + \text{jets}$  events. The additional background is Drell-Yan production of electrons and muons ( $q\bar{q} \rightarrow Z/\gamma^* \rightarrow \ell^+\ell^-$ , where  $\ell = e, \mu$ ) with the fake  $\cancel{E}_T$ . Both of the above backgrounds are estimated using data-based methods. The remaining backgrounds are from Drell-Yan production of  $\tau$  leptons and SM diboson ( $WW, WZ, ZZ$ ) production which are estimated using Monte-Carlo (MC) simulation. The detailed description of the background estimation can be found in [7]. The number of events expected and observed passing dilepton selection is presented in Table I. The uncertainties on the number of events include the statistical and systematic part. The correlations between signal and various backgrounds systematic uncertainties are taken into account. There is good agreement between data and the expectation from  $t\bar{t}$  and backgrounds.

We use the  $\cos\theta^*$  of the leptons defined above to determine the  $W$  boson polarization fractions. In order to reconstruct  $\cos\theta^*$ , the full  $t\bar{t}$  kinematic chain must be reconstructed. The dilepton channel presents an under-constrained system due to the two undetected neutrinos. We use a simple modification of the kinematic method previously developed for the measurement of the top-quark mass, denoted as the KIN method in [14]. We solve the kinematic equations by Newton’s method for nonlinear systems of equations using the top-quark mass constraint with  $m_{\text{top}} = 175 \text{ GeV}/c^2$ . In principle each event provides two measurements of  $\cos\theta^*$ ; in practice, the re-

TABLE I: Expected and observed number of signal and background events after the dilepton selection assuming  $\sigma_{t\bar{t}} = 6.7$  pb ( $m_{top} = 175$  GeV/ $c^2$ ).

Source	Events
Diboson	17.45 $\pm$ 4.64
$Z/\gamma^* \rightarrow \tau\tau$	12.26 $\pm$ 2.18
$Z/\gamma^* \rightarrow ee + \mu\mu$	22.40 $\pm$ 3.24
Fakes	53.69 $\pm$ 14.71
Total background	105.80 $\pm$ 17.24
$t\bar{t}$ ( $\sigma = 6.7$ pb)	222.44 $\pm$ 10.61
Total SM expectation	328.24 $\pm$ 27.61
Observed	343

construction under our assumptions can fail in which case we don't consider such event.

In order to perform the fit to the  $\cos\theta^*$  distribution we create templates using  $t\bar{t}$  MC simulated samples for exclusive left-handed, longitudinal and right-handed  $W$  bosons using a customized HERWIG [15, 16] MC generator. We create the templates separately for  $b$ -tag and no-tag subsamples which turn out to be similar. Figure 1 shows the templates for the inclusive sample for both the signal and background.

Due to various selection and reconstruction effects (e.g., we consider the two highest  $E_T$  jets as jets coming from  $b$ -quark hadronization while there is a possibility that one of these two jets comes from initial (ISR) or final (FSR) state gluon radiation) the templates vary significantly from the theoretical distributions in Eq. (1). The effect of the lepton isolation cut is seen as a softening of the theoretical peaks near  $\cos\theta^* = -1$ . Furthermore, the KIN reconstruction method requires the lepton-jet- $\cancel{E}_T$  mass to be close to the mass of the top quark so that the reconstruction is inefficient for high lepton-jet pair masses ( $\cos\theta^* \simeq +1$ ). This gives a polarization-dependent reconstruction efficiency of about 95%, 92%, and 87%, respectively, for left-handed, longitudinal, and right-handed  $W$  bosons. For the background events, the reconstruction efficiency is only 71%. There are also differences in the acceptance of dilepton events. The largest difference is between events with two left-handed  $W$  bosons where the acceptance is about 30% smaller relative to the acceptance of events having two longitudinal  $W$  bosons. This is mainly due to the dependence of the acceptance on the lepton  $p_T$  and isolation (leptons from left-handed  $W$  bosons tend to be less isolated and have smaller  $p_T$ ).

We combine the signal and background templates taking into account the above  $W$  polarization dependent efficiencies. We use an unbinned likelihood method which determines the  $f_0$  and  $f_+$  polarization fractions that best correspond to the observed  $\cos\theta^*$  distribution. The Gaussian constraint on the number of background events




FIG. 1: The signal templates for left-handed, longitudinal and right-handed  $W$  bosons together with the background template for the inclusive dilepton selection.

and the Poisson constraint on the total number of events are included in the likelihood formula. We multiply the likelihoods for “ $b$ -tag” and “non-tag” subsamples to arrive at the final likelihood. The method has been extensively tested in simulated samples across a full range of physically possible values of  $f_0$  and  $f_+$  parameters. From these tests, we obtain small corrections to the measured values of  $f_0$  and  $f_+$ .

The determination of  $W$  boson polarization fractions by our method is sensitive to different sources of theoretical and experimental uncertainties, such as the MC simulated templates, the jet reconstruction algorithms, and jet corrections. We have performed MC studies of simulated experiments in order to estimate these systematic uncertainties. One of the largest sources of systematic uncertainty comes from the jet energy scale (JES). We have studied this uncertainty by changing the corrections by  $\pm 1\sigma$  of the JES uncertainty [12]. Another large systematic uncertainty is modeling of the signal which we estimate as variations in the ISR and FSR, using different parton distribution functions (PDF) and different MC generators (see [7] for details). We estimate the systematic uncertainty due to the background template shape by changing each individual background within its rate uncertainty thus changing the overall shape. We then combine all these shifts (in quadrature) to obtain an overall background shape uncertainty. The uncertainty in the total number of expected background events is taken into account in the fitting procedure where the amount of background is allowed to float. The method-specific systematic uncertainties are due to limited statistics of the signal and background templates and are evaluated by fluctuating the templates bin-by-bin. An additional (small) uncertainty is due to the instantaneous luminosity which determines the mean number of interactions per bunch crossing. The systematic uncertainties are summarized in Table II, with the total systematic uncertainty of

the measurement being the sum in quadrature of all the partial systematic uncertainties from the various sources.

TABLE II: Summary of systematic uncertainties.

Source	$\Delta f_0^{1D}$	$\Delta f_+^{1D}$	$\Delta f_0^{2D}$	$\Delta f_+^{2D}$
Jet energy scale	0.033	0.019	0.002	0.020
Generators	0.035	0.019	0.016	0.011
ISR/FSR	0.024	0.010	0.040	0.017
PDF	0.010	0.003	0.025	0.009
Background shape	0.012	0.005	0.023	0.010
Template statistics				
Signal	0.010	0.005	0.024	0.012
Background	0.007	0.004	0.015	0.007
Instant. luminosity	0.016	0.008	0.013	0.002
Total	0.059	0.031	0.063	0.034

We assume a fixed top-quark mass of  $m_{top} = 175$  GeV/ $c^2$  in our dilepton measurement. However, as already noted, within the SM the fraction of  $W$  bosons with a given polarization directly depends on the top-quark and  $W$ -boson masses. Therefore, we do not include this effect in the systematic uncertainties. Rather, we provide the  $m_{top}$  dependence of the reconstructed fractions. We estimate that there is a linear shift in reconstructed  $f_0$  ( $f_+$ ) of  $\pm 0.004$  ( $\pm 0.005$ ) and  $\pm 0.012$  ( $\pm 0.006$ ) per  $\pm 1$  GeV/ $c^2$  change in the top-quark mass for 2D and 1D measurements, respectively.

There are 304 events (118 in “ $b$ -tag” and 186 in “non-tag” subsamples) passing dilepton selection and kinematic reconstruction, consistent with the SM expectation of  $284.3 \pm 22.7$  events. The comparison of  $\cos \theta^*$  between data and the expectations for SM  $t\bar{t}$  signal and background can be seen in Fig. 2. There is a good agreement between data and the SM expectation ( $\chi^2 = 6.5$  for 9 degrees of freedom, corresponding to a  $p$ -value of 69%).

We perform a model-independent simultaneous determinations of both  $f_0$  and  $f_+$  fractions:  $f_0 = 0.71^{+0.18}_{-0.17}(\text{stat})$  and  $f_+ = -0.07 \pm 0.09(\text{stat})$ . There is a strong negative correlation of  $-0.88$  between the statistical uncertainties of  $f_0$  and  $f_+$ . We also measure each polarization fraction when the other is fixed to its SM value. We measure  $f_0 = 0.59 \pm 0.09(\text{stat})$  when  $f_+$  is so fixed and measure  $f_+ = -0.07 \pm 0.04(\text{stat})$  when  $f_0$  is so fixed. We also find  $f_+ < 0.07$  at 95% C.L. when  $f_0$  is so fixed following Bayesian procedure assuming constant a priori probability for the  $f_+$  within physically possible range.

The CDF measurement performed in the single lepton channel obtained the following result [4], assuming a top-quark mass 175 GeV/ $c^2$ :  $f_0 = 0.88 \pm 0.11(\text{stat}) \pm 0.06(\text{syst})$  and  $f_+ = -0.15 \pm 0.07(\text{stat}) \pm 0.06(\text{syst})$  with the correlation of  $-0.6$  between  $f_0$  and  $f_+$ . This is consistent with the result presented in this Letter. We combine both results using the analytic best linear unbi-




FIG. 2: The comparison of  $\cos \theta^*$  distribution between data and the expected SM  $t\bar{t}$  signal and background. Each event has two entries in the histogram.

ased estimator method [17, 18]. When combining these results, in order to be close to the world average top-quark mass [3], we use the top-quark mass dependence (presented above) to correct to a top-quark mass 172.5 GeV/ $c^2$ . We also include the top-quark mass-related systematic uncertainty corresponding to  $\pm 1.1$  GeV/ $c^2$  uncertainty on  $m_{top}$  [3]. The results of the dilepton and single lepton channels are statistically independent. There is a strong negative correlation of the statistical uncertainty between the  $f_0$  and  $f_+$  observables for both channels as mentioned above. The systematic uncertainties are theoretically dominated and are assumed to be 100% correlated between the measurements, with the exception of the method-specific systematic uncertainties (signal and background template statistics) which are treated as uncorrelated. The luminosity-related systematic uncertainty applies only to the dilepton measurement. For a given measurement, we assume that the  $f_0$  and  $f_+$  uncertainties are 100% anti-correlated for each systematic uncertainty category. Table III presents the full correlation matrix between the measurements and their weights in the combination. The combined result for the simultaneous measurement is  $f_0 = 0.843 \pm 0.093(\text{stat}) \pm 0.054(\text{syst})$  and  $f_+ = -0.155 \pm 0.052(\text{stat}) \pm 0.039(\text{syst})$ . The combination has a  $\chi^2$  value of 0.99 for two degrees of freedom, corresponding to a  $p$ -value of 61% for consistency between the input measurements. The combined values of  $f_0$  and  $f_+$  have a correlation coefficient  $-0.81$ . We also combine the measurements of one polarization fraction when the other one is fixed to SM expected value. In this case, we arrive at  $f_0 = 0.637 \pm 0.055(\text{stat}) \pm 0.047(\text{syst})$  ( $f_+$  is fixed) and  $f_+ = -0.068 \pm 0.024(\text{stat}) \pm 0.038(\text{syst})$  ( $f_0$  is fixed). The combination for  $f_0$  ( $f_+$ ) has a  $\chi^2$  of 1.04 (0.61) for one degree of freedom, corresponding to a  $p$ -value of 31% (44%) for consistency between the input

TABLE III: The correlations between the measurements and their weights in the  $f_0$  and  $f_+$  combined result. The results from single lepton channel are labeled as ‘LJ’, the dilepton results as ‘DIL’.

Measurement	Correlation matrix			Weight	Weight	
				for $f_0$ (%)	for $f_+$ (%)	
LJ $f_0$	1			80.6	-21.8	
DIL $f_0$	0.13	1		19.4	21.8	
LJ $f_+$	-0.72	-0.15	1	18.9	24.0	
DIL $f_+$	-0.12	-0.88	0.16	1	-18.9	76.0

measurements.

To summarize, we have performed the measurement of  $W$  boson polarization fractions in top-quark dilepton decays. We have also combined our dilepton measurement with our previous measurement in the single lepton channel. Our results are consistent with the SM expectations and do not require the introduction of new physics. They agree with the results obtained by the D0 collaboration [5] which are of comparable precision. Our method is also the first model-independent measurement of the  $W$  polarization in the dilepton channel from CDF.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korean World Class University Program, the National Research Foundation of Korea; the Science and Technology Facilities Council and the Royal Society, UK; the Institut National de Physique Nucleaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010, Spain; the Slovak R&D Agency; the Academy of Finland; and the Australian Research

Council (ARC).

- 
- [1] J. A. Aguilar-Saavedra *et al.*, Eur. Phys. J. C **50**, 519 (2007).
  - [2] K. Nakamura *et al.* (Particle Data Group), J. Phys. G **37**, 075021 (2010) and 2011 partial update for the 2012 edition.
  - [3] T. Aaltonen *et al.* (CDF and D0 Collaboration), arXiv:hep-ex/1007.3178.
  - [4] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **105**, 042002 (2010).
  - [5] V. M. Abazov *et al.* (D0 Collaboration), Phys. Rev. D **83**, 032009 (2011).
  - [6] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 032001 (2005).
  - [7] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. D **82**, 052002 (2010).
  - [8] CDF uses a cylindrical coordinate system with the  $z$  axis along the proton beam axis. Pseudorapidity is  $\eta \equiv -\ln \tan(\theta/2)$ , where  $\theta$  is the polar angle relative to the proton beam direction and  $\phi$  is the azimuthal angle while  $p_T = |p| \sin \theta$ ,  $E_T = E \sin \theta$ .
  - [9] For the muons, transverse momentum  $p_T$  rather than transverse energy  $E_T$  is considered in the text.
  - [10] The missing transverse energy ( $\vec{\cancel{E}}_T$ ) is defined by  $\vec{\cancel{E}}_T = -\sum_i E_T^i \hat{n}_i$ , where  $i$  denotes calorimeter tower number with  $|\eta| < 2.5$ , and  $\hat{n}_i$  is a unit vector perpendicular to the beam axis and pointing at the  $i^{\text{th}}$  calorimeter tower. We also define  $\cancel{E}_T = |\vec{\cancel{E}}_T|$ .
  - [11] A jet is defined as a fixed-cone cluster in the calorimeter with a cone size of 0.4 in azimuth-pseudorapidity ( $\eta$ ) space.
  - [12] A. Bhatti *et al.* (CDF Collaboration), Nucl. Instrum. Methods A **566**, 375 (2006).
  - [13] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 052003 (2005).
  - [14] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. D **73**, 112006 (2006).
  - [15] E. H. Guillian, Ph.D. thesis, University of Michigan, 1999, Fermilab-Thesis-1999-51.
  - [16] G. Corcella *et al.*, J. High Energy Phys. 01 (2001) 10.
  - [17] L. Lyons, D. Gibaut, and P. Clifford, Nucl. Instrum. Methods A **270** (1988) 110.
  - [18] A. Valassi, Nucl. Instrum. Methods A **500** (2003) 391.