

Non-Unitarity, sterile neutrinos, and Non-Standard neutrino Interactions

Mattias Blennow,^{1,*} Pilar Coloma,^{2,†} Enrique Fernandez-Martinez,^{3,4,‡}
 Josu Hernandez-Garcia,^{3,4,§} and Jacobo Lopez-Pavon^{5,¶}

¹*Department of Theoretical Physics,
 School of Engineering Sciences, KTH Royal Institute of Technology,
 Albanova University Center, 106 91 Stockholm, Sweden*

²*Theoretical Physics Department, Fermi National Accelerator Laboratory,
 P.O. Box 500, Batavia, IL 60510, USA*

³*Departamento de Física Teórica, Universidad Autónoma de Madrid,
 Cantoblanco E-28049 Madrid, Spain*

⁴*Instituto de Física Teórica UAM/CSIC,
 Calle Nicolás Cabrera 13-15, Cantoblanco E-28049 Madrid, Spain*

⁵*INFN, Sezione di Genova, via Dodecaneso 33, 16146 Genova, Italy*

Abstract

The simplest Standard Model extension to explain neutrino masses involves the addition of right-handed neutrinos. At some level, this extension will impact neutrino oscillation searches. In this work we explore the differences and similarities between the case in which these neutrinos are kinematically accessible (sterile neutrinos) or not (mixing matrix non-unitarity). We clarify apparent inconsistencies in the present literature when using different parametrizations to describe these effects and recast both limits in the popular neutrino non-standard interaction (NSI) formalism. We find that, in the limit in which sterile oscillations are averaged out at the near detector, their effects at the far detector coincide with non-unitarity at leading order, even in presence of a matter potential. We also summarize the present bounds existing in both limits and compare them with the expected sensitivities of near-future facilities taking the DUNE proposal as a benchmark. We conclude that non-unitarity effects are too constrained to impact present or near future neutrino oscillation facilities but that sterile neutrinos can play an important role at long baseline experiments. The role of the near detector is also discussed in detail.

*Electronic address: emb@kth.se

†Electronic address: pcoloma@fnal.gov

‡Electronic address: enrique.fernandez-martinez@uam.es

§Electronic address: josu.hernandez@uam.es

¶Electronic address: jpgavon@ge.infn.it

I. INTRODUCTION

The simplest extension of the Standard Model (SM) of particle physics able to account for the evidence for neutrino masses and mixings is the addition of right-handed neutrinos to its particle content. A Majorana mass term for these new singlet fermions is then allowed by all symmetries of the Lagrangian. This new mass scale at which lepton number is violated could provide a mechanism to also explain the origin of the observed matter-antimatter asymmetry of the Universe [1] and is a necessary missing piece to solve the mysterious flavour puzzle. However, given that this new mass scale is not related to the electroweak symmetry breaking mechanism, there is no theoretical guidance for its value. A large Majorana scale leads to the celebrated seesaw mechanism [2–5], providing a very natural explanation of the lightness of neutrino masses. On the other hand, it also leads to unnaturally large contributions to the Higgs mass, worsening the hierarchy problem [6]. Conversely, a light neutrino mass could also naturally stem from a symmetry argument [7–13]. Indeed, neutrino masses are protected by the lepton number symmetry and, if this is an approximate symmetry of the theory, a large hierarchy of scales is not required to naturally accommodate the lightness of neutrinos. Thus, the value of this scale of new physics can only be probed experimentally and, depending on its value, very different and interesting phenomenology would be induced in different observables.

In this work we analyze the phenomenological impact of these new physics in neutrino oscillation facilities. If the new mass scale is kinematically accessible in meson decays, the sterile states will be produced in the neutrino beam. On the other hand, if the extra neutrinos are too heavy to be produced, the effective three by three PMNS matrix will show unitarity deviations. We will refer to these situations as sterile and non-unitary neutrino oscillations, respectively. The aim of this work is to discuss in which limits these two regimes lead to the same impact on the oscillation probabilities, and reconcile apparently inconsistent results in previous literature.

This work is organized as follows. In section II we will compare the non-unitarity and sterile neutrino phenomenology and discuss in which cases both limits are equivalent. In Section III we will present and solve the apparent inconsistency present in the literature studying non-unitarity effects in different parametrizations, and provide a mapping between the two. In Section IV we will recast both scenarios using the popular NSI parametrization. The existing bounds, applicable in both regimes, from present observables are summarized in Section V. Finally, in Section VI we present the sensitivity of the prospective Deep Underground Neutrino Experiment (DUNE) to these effects, and our conclusions are summarized in Section VII.

II. NON-UNITARITY AND STERILE NEUTRINO PHENOMENOLOGY COMPARISON

In this section we will show how, under certain conditions, the phenomenology of non-unitarity and sterile neutrino oscillations are equivalent to leading order in the active-heavy mixing parameters, not only in vacuum but also in matter. If n extra right-handed neutrinos are added to the SM Lagrangian, the full mixing matrix (including both light and heavy

states) can be written as

$$\mathcal{U} = \begin{pmatrix} N & \Theta \\ R & S \end{pmatrix}, \quad (1)$$

where N represents the 3×3 active-light sub-block (i.e., the PMNS matrix), which will no longer be unitary¹. Here, Θ is the $3 \times n$ sub-block that includes the mixing between active and heavy states, while the R and S sub-blocks define the mixing of the sterile states with the light and heavy states, respectively. Note that both R and S are only defined up to an unphysical rotation of the sterile states and that neither of them will be involved when considering oscillations among active flavours.

A. Non-unitarity case

In the case of non-unitarity, only the light states are kinematically accessible and the amplitude for producing one of these states in conjunction with a charged lepton of flavour α in a particular decay is proportional to the mixing matrix element $N_{\alpha i}^*$. In the mass eigenstate basis, the evolution of the produced neutrino state is given by the Hamiltonian [14]

$$H = \frac{1}{2E} \begin{pmatrix} 0 & 0 & 0 \\ 0 & \Delta m_{21}^2 & 0 \\ 0 & 0 & \Delta m_{31}^2 \end{pmatrix} + N^\dagger \begin{pmatrix} V_{\text{CC}} + V_{\text{NC}} & 0 & 0 \\ 0 & V_{\text{NC}} & 0 \\ 0 & 0 & V_{\text{NC}} \end{pmatrix} N, \quad (2)$$

where $V_{\text{CC}} = \sqrt{2}G_F n_e$ and $V_{\text{NC}} = -G_F n_n / \sqrt{2}$ are the charged-current (CC) and neutral-current (NC) matter potentials, respectively. The oscillation evolution matrix S^0 in this basis is now defined as the solution to the differential equation

$$i\dot{S}^0 = HS^0 \quad (3)$$

with the initial condition $S^0(0) = I$, I being the identity matrix. For a constant matter potential, this equation has the formal solution

$$S^0 = \exp(-iHL). \quad (4)$$

The amplitude for a neutrino in the mass eigenstate j to interact as a neutrino of flavour β is given by the mixing matrix element $N_{\beta j}$, which means that the oscillation probability will be given by

$$P_{\alpha\beta} = |(NS^0N^\dagger)_{\beta\alpha}|^2. \quad (5)$$

Here $P_{\alpha\beta}$ denotes the ‘‘theoretical’’ oscillation probabilities (although it should be noted that they do not add up to one), defined as the ratio between the observed number of events divided by the product of the SM-predicted flux times cross section. In other words, $P_{\alpha\beta}$ is the factor that would be needed to obtain the number of events after convolution with the standard model flux and cross sections.

However, in practice neutrino oscillation experiments do not measure $P_{\alpha\beta}$. Most present and future experiments rather determine the flux and cross sections via near detector data

¹ Note that this is true regardless whether the extra states are kinematically accessible or not.

and extrapolate to the far detector by correcting for the different geometries, angular apertures, and detection cross sections. In this scenario, the oscillation probability would then be inferred from the ratio

$$\mathcal{P}_{\alpha\beta} = \frac{R_\beta}{R_\alpha}, \quad (6)$$

where R_β and R_α are the observed event rates at the far detector and the corresponding extrapolation of the near detector result, respectively. For the near detector, we assume that the phases corresponding to the propagation of the light neutrinos have not yet developed significantly and therefore $S^0 = I$, resulting in the experimentally inferred probability

$$\mathcal{P}_{\alpha\beta} = \frac{|(NS^0N^\dagger)_{\beta\alpha}|^2}{((NN^\dagger)_{\alpha\alpha})^2}. \quad (7)$$

In the SM limit the matrix N becomes unitary and, thus, $NN^\dagger = I$ and $\mathcal{P}_{\alpha\beta} = P_{\alpha\beta}$ as expected.

B. Sterile neutrino case

In the sterile neutrino scenario, all of the states are kinematically accessible and the full oscillation evolution matrix \mathcal{S} , involving both light and heavy states, takes the form

$$\mathcal{S} = \mathcal{U}\mathcal{S}^0\mathcal{U}^\dagger, \quad (8)$$

where \mathcal{S}^0 is the full $n \times n$ evolution matrix expressed in the mass eigenbasis. For vacuum oscillations, we find that $\mathcal{S}^0 = \text{diag}(\exp(-i\Delta m_{j1}^2 L/2E))$. Therefore, the active neutrino 3×3 sub-block S can be simplified to

$$S_{\alpha\beta} = \sum_{i \in \text{light}} N_{\alpha i} S_{ij}^0 N_{\beta j}^* + \sum_{J \in \text{heavy}} \Theta_{\alpha J} \Theta_{\beta J}^* \Phi_J, \quad (9)$$

where α, β stand for active neutrino flavors, Φ_J is the phase factor acquired by the heavy state J as it propagates, and S^0 is defined in the same way as in the non-unitarity case.

In the limit of large mass squared splitting between the light and heavy states (*i.e.*, $\Delta m_{iJ}^2 L/E \gg 1$) the oscillations are too fast to be resolved at the detector and only an averaged-out effect is observable. In this averaged-out limit, the cross terms between the first and second term in the evolution matrix average to zero and we find

$$P_{\alpha\beta} = |S_{\alpha\beta}|^2 = \left| \sum_i N_{\alpha i} S_{ij}^0 N_{\beta j}^* \right|^2 + \mathcal{O}(\Theta^4), \quad (10)$$

which recovers the same expression as Eq. (5) up to the $\mathcal{O}(\Theta^4)$ corrections.² Thus, we can conclude that averaged-out sterile neutrino oscillations in vacuum are equivalent to

² Note that this expression is also applicable whenever the light and heavy states decohere due to wave packet separation.

non-unitarity to leading order (this equivalence is indeed lost at higher orders). We will therefore concentrate on this averaged-out limit for the rest of this paper.

For oscillations in the presence of matter, the sterile neutrino oscillations will be subjected to a matter potential that in the flavour basis takes the form

$$\mathcal{H}_{\text{mat}}^f = \begin{pmatrix} V_{3\times 3} & 0 \\ 0 & 0 \end{pmatrix}, \quad (11)$$

where

$$V_{3\times 3} = \begin{pmatrix} V_{\text{CC}} + V_{\text{NC}} & 0 & 0 \\ 0 & V_{\text{NC}} & 0 \\ 0 & 0 & V_{\text{NC}} \end{pmatrix}. \quad (12)$$

If the matter potential is small in comparison to the light-heavy energy splitting $\Delta m_{iJ}^2/2E$, the light-heavy mixing in matter will be given by

$$\tilde{\Theta}_{\alpha J} = \Theta_{\alpha J} + \frac{2E}{\Delta m_{iJ}^2} (\delta_{\alpha e} V_{\text{CC}} \Theta_{eJ} + \Theta_{\alpha J} V_{\text{NC}}) \quad (13)$$

to first order in perturbation theory. In the limit $\Delta m_{iJ}^2/2E \gg V_{\text{CC}}, V_{\text{NC}}$, we can therefore neglect the difference between the heavy mass eigenstates in vacuum and in matter, and apply Eq. (10). Thus, we conclude from this that the matter Hamiltonian in the light sector can be computed in exactly the same way as for the non-unitarity scenario and we therefore find the very same expressions for the ‘‘theoretical’’ probability in Eq. (5) as for the non-unitarity case, at leading order in Θ .

In the case of sterile neutrinos one also needs to consider the impact of the near detector measurements on the extraction of the experimentally measurable probability. In this work we will always assume that the oscillations due to the additional heavy states are averaged out at the far detector. However, this might not be the case at the near detector. Ideally, both sets of observables should be simulated and analyzed together consistently. Nevertheless, the following simplified limiting cases can be identified:

1. The light-heavy oscillations are averaged out already at the near detector. For practical purposes, this case is identical to the non-unitarity case and Eq. (7) also applies in the sterile case.
2. The light-heavy oscillations have not yet developed at the near detector. In this case, the near detector would measure the SM fluxes and cross sections, and therefore the denominator in Eq. (7) is equal to 1. In this case, the experimental probability would coincide with the ‘‘theoretical probability’’ in Eq. (5).
3. The oscillation frequency dictated by the light-heavy frequency matches the near detector distance. In this case, oscillations could be seen at the near detector as a function of neutrino energy, leading to more striking signals. This case will not be considered any further in this work.

III. PARAMETRIZATIONS

The two most widely used parametrizations to encode these non-unitarity effects stemming from the heavy-active mixing are

$$N = (I - \eta)U' \quad \text{or} \quad N = TU = (I - \alpha)U, \quad (14)$$

where η is a Hermitian matrix [15, 16] and T is a lower triangular matrix [17–20]. In Eq. (14) both U and U' are unitary matrices that are equivalent to the standard PMNS matrix up to small corrections proportional to the deviations encoded in η and α .

$$\eta = \begin{pmatrix} \eta_{ee} & \eta_{e\mu} & \eta_{e\tau} \\ \eta_{e\mu}^* & \eta_{\mu\mu} & \eta_{\mu\tau} \\ \eta_{e\tau}^* & \eta_{\mu\tau}^* & \eta_{\tau\tau} \end{pmatrix}, \quad \alpha = (1 - T) = \begin{pmatrix} \alpha_{ee} & 0 & 0 \\ \alpha_{\mu e} & \alpha_{\mu\mu} & 0 \\ \alpha_{\tau e} & \alpha_{\tau\mu} & \alpha_{\tau\tau} \end{pmatrix} \quad (15)$$

with $\eta_{\alpha\beta}, \alpha_{\alpha\beta} \ll 1$. Note that we choose to label the α matrix elements with flavour indices for notation ease instead of using numbers as in [19]. Furthermore, in [19] the identity matrix is not singled out from α as in our Eq. (15) so that the diagonal elements α_{ii} in [19] are close to 1 instead of small. Therefore, in practice, $\alpha_{ii} \rightarrow 1 - \alpha_{\alpha\alpha}$. These changes are only cosmetic and the following discussion applies to [19] with the above-mentioned identification.

The deviations from unitarity are directly related to the heavy-active neutrino mixing. For instance, in the hermitian parametrization one can directly identify [21]

$$\eta = \frac{\Theta\Theta^\dagger}{2} \quad (16)$$

where $\Theta = m_D^\dagger M^{-1}$ is the heavy-active mixing given by the ratio of the Dirac over the Majorana mass scales. Thus, $(1 - \eta)$ is just the first term in the cosine series correcting the unitary rotation U' . It is also straightforward to obtain the relation between the heavy-active neutrino mixing and the α parameters in the triangular parametrization, if one considers that the heavy-active mixing can also be encoded by introducing additional complex rotations characterized by new mixing angles θ_{ij} , with $j > 3$. For example,

$$U_{14} = \begin{pmatrix} c_{14} & 0 & 0 & \hat{s}_{14}^* \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -\hat{s}_{14} & 0 & 0 & c_{14} \end{pmatrix}, \quad (17)$$

where $\hat{s}_{ij} = e^{i\delta_{ij}} s_{ij}$, $s_{ij} = \sin\theta_{ij}$ and $c_{ij} = \cos\theta_{ij}$. In the correct order, these extra rotations lead to a lower triangular matrix. For 3 extra neutrinos we can use $U_{36}U_{26}U_{16}U_{35}U_{25}U_{15}U_{34}U_{24}U_{14}$ (where we have not included unphysical rotations among the sterile neutrinos), leading to [17]:

$$\alpha \simeq \begin{pmatrix} \frac{1}{2}(s_{14}^2 + s_{15}^2 + s_{16}^2) & 0 & 0 \\ \hat{s}_{14}\hat{s}_{24}^* + \hat{s}_{15}\hat{s}_{25}^* + \hat{s}_{16}\hat{s}_{26}^* & \frac{1}{2}(s_{24}^2 + s_{25}^2 + s_{26}^2) & 0 \\ \hat{s}_{14}\hat{s}_{34}^* + \hat{s}_{15}\hat{s}_{35}^* + \hat{s}_{16}\hat{s}_{36}^* & \hat{s}_{24}\hat{s}_{34}^* + \hat{s}_{25}\hat{s}_{35}^* + \hat{s}_{26}\hat{s}_{36}^* & \frac{1}{2}(s_{34}^2 + s_{35}^2 + s_{36}^2) \end{pmatrix}, \quad (18)$$

which is accurate to second order in the (small) extra mixing angles.

In principle, the two parametrizations should be equally valid. However, the alternative use of each of them seemingly leads to inconsistent results. As an illustrative example, let us compare the ν_μ disappearance probability in the atmospheric regime in the two parametrizations, obtained at linear order in the non-unitarity parameters and for $\theta_{13} = 0$

$$\begin{aligned}\mathcal{P}_{\mu\mu}^\eta &= 1 - \{ \sin^2 2\theta'_{23} - 2\text{Re}[\eta_{\mu\tau}] \sin 4\theta'_{23} \} \sin^2 \Delta_{31}, \\ \mathcal{P}_{\mu\mu}^\alpha &= 1 - \sin^2 2\theta_{23} \sin^2 \Delta_{31},\end{aligned}\tag{19}$$

where $\Delta_{ij} = \Delta m_{ij}^2 L/4E$. Here, $\mathcal{P}_{\alpha\beta}$ denotes the ‘‘experimental’’ oscillation probability in vacuum including the normalization factors as discussed in Sec. II.

The naive conclusion derived from Eq. (19) is that for the Hermitian parametrization good sensitivity to the non-unitarity parameter $\eta_{\mu\tau}$ is expected in this channel, since it appears at linear order. Conversely, the triangular parametrization does not show this effect. This apparent inconsistency stems from the fact that the unitary matrices U and U' which are identified with the standard unitary PMNS matrix in each parametrization are, in fact, different. As we will show below, the differences between the two are linear in the non-unitarity parameters, and the two matrices can be related to each other. Starting from Eq. (14) a unitary rotation V can be performed to relate U and U'

$$N = (I - \alpha)U = (I - \eta)V V^\dagger U' \tag{20}$$

and therefore

$$I - \alpha = (I - \eta)V \quad \text{and} \quad U = V^\dagger U'. \tag{21}$$

From the first relation in Eq. (21) the elements of V can be identified as

$$V = I - \begin{pmatrix} 0 & -\eta_{e\mu} & -\eta_{e\tau} \\ \eta_{e\mu}^* & 0 & -\eta_{\mu\tau} \\ \eta_{e\tau}^* & \eta_{\mu\tau}^* & 0 \end{pmatrix} \tag{22}$$

at linear order in η . Substituting again in Eq. (21) the relations

$$\begin{pmatrix} \alpha_{ee} & 0 & 0 \\ \alpha_{\mu e} & \alpha_{\mu\mu} & 0 \\ \alpha_{\tau e} & \alpha_{\tau\mu} & \alpha_{\tau\tau} \end{pmatrix} = \begin{pmatrix} \eta_{ee} & 0 & 0 \\ 2\eta_{e\mu}^* & \eta_{\mu\mu} & 0 \\ 2\eta_{e\tau}^* & 2\eta_{\mu\tau}^* & \eta_{\tau\tau} \end{pmatrix} \tag{23}$$

and

$$U = V^\dagger U' = \left(I + \begin{pmatrix} 0 & -\eta_{e\mu} & -\eta_{e\tau} \\ \eta_{e\mu}^* & 0 & -\eta_{\mu\tau} \\ \eta_{e\tau}^* & \eta_{\mu\tau}^* & 0 \end{pmatrix} \right) U' \tag{24}$$

are found. This implies the following mapping between the two sets of mixing angles³ in U'

³ Note that, apart from correcting the PMNS mixing angles and CP-phase δ_{CP} at order η , phase redefinitions of the three charged leptons as well as corrections to the two neutrino Majorana phases are necessary at the same order.

and U :

$$\begin{aligned}
\theta_{12} - \theta'_{12} &= \frac{\text{Re}(s_{23}\eta_{e\tau} - c_{23}\eta_{e\mu})}{c_{13}}, \\
\theta_{13} - \theta'_{13} &= \text{Re}(-s_{23}e^{i\delta_{\text{CP}}}\eta_{e\mu} - c_{23}e^{i\delta_{\text{CP}}}\eta_{e\tau}), \\
\theta_{23} - \theta'_{23} &= -\text{Re}(\eta_{\mu\tau}) + \tan\theta_{13} \text{Re}(c_{23}e^{i\delta_{\text{CP}}}\eta_{e\mu} - s_{23}e^{i\delta_{\text{CP}}}\eta_{e\tau}), \\
\delta_{\text{CP}} - \delta'_{\text{CP}} &= \frac{\cos 2\theta_{12}}{s_{12}c_{12}c_{13}} \text{Im}(s_{23}\eta_{e\tau} - c_{23}\eta_{e\mu}) + \frac{1}{s_{13}c_{13}} \text{Im}(s_{23}e^{i\delta_{\text{CP}}}\eta_{e\mu} + c_{23}e^{i\delta_{\text{CP}}}\eta_{e\tau}) \\
&\quad - \frac{\tan\theta_{13}}{s_{23}c_{23}} \text{Im}\left(c_{23}^3e^{i\delta_{\text{CP}}}\eta_{e\mu} + s_{23}^3e^{i\delta_{\text{CP}}}\eta_{e\tau} + \frac{\eta_{\mu\tau}}{\tan\theta_{13}}\right). \tag{25}
\end{aligned}$$

When the relations given in Eqs. (23) and (25) are taken into account the predictions for the different oscillation channels coincide at leading order in the non-unitarity parameters, as they should. An important conclusion derived from this is that the determination of the mixing angles themselves will generally be affected by non-unitarity corrections. These corrections are parametrization-dependent but, when taken into account and propagated consistently, the predictions derived from both schemes agree.

For neutrino oscillation studies it seems advantageous to adopt the triangular parametrization, since it leads to fewer corrections given its structure. For instance, in the example shown in Eq. (19) there are no corrections coming from non-unitarity for this parametrization, and thus the angle θ_{23} in U can be identified with the angle determined in present global fits to a good approximation. Indeed, this is also the case for θ_{12} and θ_{13} , since the \mathcal{P}_{ee} oscillation probabilities in the solar regime (KamLAND) and in the atmospheric regime (Daya Bay, RENO, Double-Chooz) are also independent of any non-unitarity corrections at linear order when the triangular parametrization is considered and when the appropriate normalization is taken into account, see Sec. II.

Thus, the U matrix from the triangular parametrization corresponds, to a good approximation, with the unitary matrix obtained when determining θ_{12} , θ_{23} and θ_{13} through present (disappearance) neutrino oscillation measurements. Since we are here interested in the impact of non-unitarity and sterile neutrinos on neutrino oscillation phenomenology we will therefore use the triangular parametrization in the remainder of this work.

IV. NSI

Both types of new physics effects in neutrino oscillations discussed above can be described through the Non-Standard Interaction (NSI) formalism, which parametrizes the new physics effects in neutrino production, detection, and propagation processes in a completely model-independent way. Let us first focus on NSI affecting neutrino production and detection. When these effects are included, the oscillation probability is given by

$$P_{\alpha\beta} = \left| \left[(1 + \epsilon^d) U S^0 U^\dagger (1 + \epsilon^s) \right]_{\beta\alpha} \right|^2, \tag{26}$$

where ϵ^s and ϵ^d are general 3×3 complex matrices which represent the NSI modifications to the production and detection diagrams, respectively. S^0 is defined in Eq. (4) with the Hamiltonian H given in Eq. (2). When the NSI effects in propagation are not considered

or negligible, the oscillation probabilities in vacuum in the non-unitarity (Eq. (5)), sterile neutrino (Eq. (10)) and NSI (Eq. (26)) frameworks are equivalent at leading order in the new physics parameters with the identification

$$\epsilon_{\beta\alpha}^{s*} = \epsilon_{\alpha\beta}^d = -\alpha_{\alpha\beta}. \quad (27)$$

This mapping can be easily obtained just considering the triangular parameterization, which can be applied in both the non-unitarity and sterile neutrino cases, in Eqs. (5) or (10) and comparing the result to Eq. (26).

On the other hand, NSI affecting neutrino propagation are usually described through the Hamiltonian

$$H = \frac{1}{2E} \begin{pmatrix} 0 & 0 & 0 \\ 0 & \Delta m_{21}^2 & 0 \\ 0 & 0 & \Delta m_{31}^2 \end{pmatrix} + V_{\text{CC}} U^\dagger \begin{pmatrix} 1 + \epsilon_{ee} & \epsilon_{e\mu} & \epsilon_{e\tau} \\ \epsilon_{e\mu}^* & \epsilon_{\mu\mu} & \epsilon_{\mu\tau} \\ \epsilon_{e\tau}^* & \epsilon_{\mu\tau}^* & \epsilon_{\tau\tau} \end{pmatrix} U, \quad (28)$$

in the mass basis, where U is the standard unitary PMNS matrix, and $\epsilon_{\alpha\beta}$ and $\epsilon_{\alpha\alpha}$ are complex and real parameters respectively. In order to understand how the non-unitarity/sterile neutrino corrected matter effects can be translated to this parametrization, we introduce the triangular parameterization of N into Eq. (2), obtaining the following Hamiltonian at leading order in α

$$H = \frac{1}{2E} \begin{pmatrix} 0 & 0 & 0 \\ 0 & \Delta m_{21}^2 & 0 \\ 0 & 0 & \Delta m_{31}^2 \end{pmatrix} + \frac{V_{\text{CC}}}{2} U^\dagger \begin{pmatrix} 2 - 2\alpha_{ee} & \alpha_{\mu e}^* & \alpha_{\tau e}^* \\ \alpha_{\mu e} & 2\alpha_{\mu\mu} & \alpha_{\tau\mu}^* \\ \alpha_{\tau e} & \alpha_{\tau\mu} & 2\alpha_{\tau\tau} \end{pmatrix} U, \quad (29)$$

where approximately equal densities of electrons n_e and neutrons n_n (for the neutral current contribution) in the Earth have been assumed (see also Ref. [22]). Comparing Eqs. (29) and (28) we find the mapping between the NSI parametrization and the lower triangular parametrization of the non-unitarity and sterile neutrino scenarios

$$\begin{aligned} \epsilon_{ee} &= -\alpha_{ee}, & \epsilon_{\mu\mu} &= \alpha_{\mu\mu}, & \epsilon_{\tau\tau} &= \alpha_{\tau\tau}, \\ \epsilon_{e\mu} &= \frac{1}{2}\alpha_{\mu e}^*, & \epsilon_{e\tau} &= \frac{1}{2}\alpha_{\tau e}^*, & \epsilon_{\mu\tau} &= \frac{1}{2}\alpha_{\tau\mu}^*, \end{aligned} \quad (30)$$

which apply for neutrino oscillation experiments in the Earth with constant matter. Note that, in presence of production and detection NSI, the same normalization as for the non-unitarity case discussed in Section II needs to be taken into account.

V. PRESENT CONSTRAINTS ON DEVIATIONS FROM UNITARITY

The mapping to NSI described above works both for the non-unitarity and the sterile neutrino contributions to neutrino oscillations. However, the present constraints on each contributions from other observables are very different. Indeed, PMNS non-unitarity from very heavy extra neutrinos induces modifications of the W and Z couplings that impact precision electroweak and flavour observables [14, 22–35]. These modification translate into very strong upper limits on the α parameters. These have been taken from Ref. [35] and are listed in the left column in Table I.

	Non-Unitarity	Averaged-out sterile
α_{ee}	$1.3 \cdot 10^{-3}$	$2.6 \cdot 10^{-2}$
$\alpha_{\mu\mu}$	$2.0 \cdot 10^{-4}$	$4.5 \cdot 10^{-2}$
$\alpha_{\tau\tau}$	$2.8 \cdot 10^{-3}$	10^{-1}
$\alpha_{\mu e}$	$6.8 \cdot 10^{-4}$ ($2.4 \cdot 10^{-5}$)	$4.8 \cdot 10^{-2}$
$\alpha_{\tau e}$	$2.7 \cdot 10^{-3}$	$7.2 \cdot 10^{-2}$
$\alpha_{\tau\mu}$	$1.2 \cdot 10^{-3}$	$9.5 \cdot 10^{-2}$

TABLE I: Current upper bounds on the α parameters in the two scenarios considered in this work. The limits are shown at 2σ and 95% CL (1 d.o.f.) for the non-unitarity (left column, taken from Ref. [35]) and averaged-out sterile neutrino (right column, taken from Refs. [36–38]) scenarios, respectively. The second number quoted in parenthesis for the $\alpha_{\mu e}$ element in the non-unitarity case includes the $\mu \rightarrow e\gamma$ observable which can in principle be evaded [39] for heavy neutrino masses close to M_W and some fine-tuning of the parameters. See main text for further details.

For light sterile neutrinos (with masses below the keV scale) these stringent constraints are, however, lost since all mass eigenstates are kinematically available in the observables used to derive the constraints and unitarity is therefore restored. Thus, for the light sterile scenario, much milder constraints stemming from the non-observation of their effects at neutrino oscillation searches apply [36–38], as shown in the right column of Table I. The constraints on the mixing with electrons α_{ee} stem from the BUGEY-3 experiment [40], which enters the averaged out regime for $\Delta m^2 \gtrsim 4 \text{ eV}^2$. Recent competitive constraints on this parameter by the Daya Bay experiment [41] have not been included since we are interested in the averaged-out regime, and Daya Bay constraints dominate for smaller Δm^2 values. Similarly, we quote the limits for the $\alpha_{\mu\mu}$ and $\alpha_{\tau\tau}$ elements from the SK analysis of atmospheric neutrino oscillations [42] also in the averaged-out regime for $\Delta m^2 \gtrsim 0.1 \text{ eV}^2$. MINOS [43] has similar constraints on $\alpha_{\tau\tau}$ while stronger limits were found in Ref. [44] but for $\Delta m^2 = 6 \text{ eV}^2$ and not in the averaged-out limit. Regarding $\alpha_{\mu\mu}$ other competitive limits can be found from the CDHS [45], IceCube [46] and MINOS [47] analyses but again not in the averaged-out limit explored here. Finally, the off-diagonal elements are constrained by the diagonal as $\alpha_{\alpha\beta} \leq 2\sqrt{\alpha_{\alpha\alpha}\alpha_{\beta\beta}}$, from Eq. (18).

VI. DUNE SENSITIVITIES

In this section we present, as an example, the sensitivities that the proposed DUNE experiment would have to PMNS non-unitarity or, equivalently, to averaged-out sterile neutrino oscillations as discussed in Section II. For this analysis we choose the triangular parametrization of the new physics effects since, as argued in Section III, its unitary part can be more directly mapped to the “standard” PMNS matrix as determined by present experiments

through neutrino oscillation disappearance channels. Indeed, production and detection new physics effects in a given channel $P_{\alpha\beta}$ only depend on the elements $\alpha_{\gamma\rho}$ such that $\gamma, \rho \leq \alpha$ or $\gamma, \rho \leq \beta$ when the flavour indices are ordered as $e < \mu < \tau$ [19]. Furthermore, when the new physics affects the near and far detectors in the same manner, the normalization of the oscillation probabilities presented in Eq. (7) has to be applied, which effectively cancels any leading order dependence on the new physics parameters $\alpha_{\beta\beta}$ in disappearance channels in vacuum. The choice of the facility under study is motivated by the strong matter effects that characterize the DUNE setup and that allow to probe not only the source and detector effects induced by the new physics, but also the matter effects which now provide sensitivity to other $\alpha_{\gamma\rho}$ parameters not necessarily satisfying $\gamma, \rho \leq \alpha$ or $\gamma, \rho \leq \beta$.

The simulation of the DUNE setup was performed with the GLoBES software [48, 49] using the DUNE CDR configuration presented in Ref. [50]. The new physics effects have been implemented in GLoBES via the MonteCUBES [51]⁴ plug-in, which has also been used to perform a Markov chain Monte Carlo (MCMC) scan of the 15-dimensional parameter space (the 6 standard oscillation parameters plus the 6 moduli of the $\alpha_{\gamma\rho}$ non-zero entries and the 3 phases of the off-diagonal elements). In the fit, the assumed true values for the standard oscillation parameters are set according to their current best-fits from Ref. [52]. The mixing angles and squared-mass splittings are allowed to vary in the simulations, using a Gaussian prior corresponding to their current experimental uncertainties from Ref. [52] centered around their true values. In the case of θ_{13} and θ_{23} the Gaussian priors are implemented on $\sin^2 2\theta$, which is a more accurate description of the present situation and, in the case of θ_{23} , allows to properly account for the octant degeneracy: $\Delta m_{21}^2 = (7.50 \pm 0.18) \cdot 10^{-5} \text{ eV}^2$, $\Delta m_{31}^2 = (2.457 \pm 0.049) \cdot 10^{-3} \text{ eV}^2$, $\theta_{12} = 33.48^\circ \pm 0.77^\circ$, $\sin^2 2\theta_{13} = 0.085 \pm 0.005$, $\sin^2 2\theta_{23} = 0.991 \pm 0.02$. The CP-violating phase is left completely free during the simulations, and its true value is set to $\delta_{\text{CP}} = -\pi/2$. Finally, a 2 % uncertainty in the PREM matter density profile [53] has also been considered.

We have performed simulations for two distinct new physics scenarios. In the first case (ND averaged) we normalize the oscillation probabilities according to Eq. (7). Indeed, as discussed in Section II, at leading order in the new physics parameters this scenario accurately describes both the effects of PMNS non-unitarity from very heavy neutrinos as well as sterile neutrino oscillations that have been averaged out both at the near detector (ND) and far detector. The second scenario (ND undeveloped) corresponds to the case where sterile neutrino oscillations are averaged out at the far detector but have not yet developed at the near detector. In this case, no extra normalization is needed and the oscillation probability is directly given by Eq. (5). Note that, for the energies and baseline characterizing the DUNE setup, only values of the sterile neutrino masses around $\Delta m^2 \sim 0.1 \text{ eV}^2$ satisfy these conditions. However, we find it instructive to study also this regime in order to remark the differences between the two scenarios and the importance of the normalization in Eq. (7) that will generally apply in most of the parameter space.

Figures 1 and 2 show the expected sensitivities to the new physics parameters. These have been obtained by assuming that the true values of all α entries are zero to obtain the corresponding expected number of events, and fitting for the corresponding parameters while

⁴ A new version of the MonteCUBES software implementing the triangular parametrization will be released shortly after this work.




FIG. 1: Expected frequentist allowed regions at the 1σ , 90% and 2σ C.L. for DUNE. All new physics parameters are assumed to be zero so as to obtain the expected sensitivities. The left panels (ND averaged) correspond to the non-unitarity case, or to the sterile case when the light-heavy oscillations are averaged out in the near and far detectors. The right panels (ND undeveloped) give the sensitivity for the sterile case when the light-heavy oscillations have not yet developed in the near detector, but are averaged out in the far. The solid lines correspond to the analysis of DUNE data alone, while the dashed lines include the present constraints on sterile neutrino mixing from the right column in Tab. I.

marginalizing over all other standard and new physics parameters. The resulting frequentist allowed regions are shown at at 1σ , 90%, and 2σ C.L.

The sensitivities obtained for all parameters fall at least one order of magnitude short of the current bounds on the non-unitarity from heavy neutrino scenario presented in Tab. I. Thus, the standard three-family oscillations explored at DUNE (and the other present and near-future oscillation facilities) will be free from the possible ambiguities that could otherwise be induced by this type of new physics [54]. The situation is quite different if the results are interpreted in terms of an averaged-out sterile neutrino, since present constraints are weaker in this case. We will therefore focus on this scenario for the rest of our discussion and also study the case in which DUNE data is complemented by our present prior constraints on averaged-out sterile neutrino mixing, see Tab. I, since synergies between the data sets can be present. This scenario is depicted with dashed lines in Figs. 1 and 2. As an example of such synergy, the sensitivity to the real part of $\alpha_{\mu\tau}$ improves in the combination with respect to the prior itself.

Another conclusion that can be drawn from Fig. 1 is that the sensitivities to the diagonal parameters α_{ee} and $\alpha_{\mu\mu}$ are significantly stronger for the ND undeveloped (right panels) as compared to the ND averaged scenario (left panels). This was to be expected since the source and detection effects that provide a leading order sensitivity to the diagonal parameters are totally or partially cancelled once the normalization of Eq. (7) is included. In the disappearance channel both effects cancel in the ratio, while for the appearance channel there is a partial cancellation that only allows the experiment to be sensitive to the combination $\alpha_{ee} - \alpha_{\mu\mu}$. This leads to a pronounced correlation among α_{ee} and $\alpha_{\mu\mu}$, seen in the upper left panel of Fig. 1.

From a phenomenological point of view we observe that, if both near and far detectors are affected by the new physics in the same way (as is the case when the sterile neutrino oscillations are averaged out at both detectors, or in the non-unitarity scenario) their effects are more difficult to observe since they cannot be disentangled from the flux and cross section determination at the near detector. Conversely, in the case in which sterile neutrino oscillations have not yet developed at the near detector but are averaged out at the far, the flux determined by both detectors will have a different normalization. Thus, a strong linear sensitivity to α_{ee} (comparable to present constraints) and $\alpha_{\mu\mu}$ (improving over present constraints) is obtained from detector and source effects respectively.

Fig. 1 also shows strong correlations in the middle left panel, involving $\alpha_{\tau\tau}$ and α_{ee} . Indeed, sensitivity to $\alpha_{\tau\tau}$ comes through the matter effects, which only depend on the diagonal entries through their differences $\alpha_{\beta\beta} - \alpha_{\gamma\gamma}$, since a global term in the form of $\alpha_{\gamma\gamma}I$ does not affect neutrino oscillations at leading order in α . In these panels we also observe a large difference between the allowed regions for $\alpha_{\tau\tau}$ once prior constraints on the α parameters are included in the analysis, by comparing the solid and dashed lines. This is due to the lifting of degeneracies involving θ_{23} and the combination $\alpha_{\tau\tau} - \alpha_{\mu\mu}$, and will be discussed in more detail below.

Unlike in the non-unitarity case (where current constraints are too strong to lead to any observable effect), interesting correlations and degeneracies among the standard and new physics parameters can indeed take place in the averaged-out sterile neutrino scenario [38, 55–58]. In our results, even though the true values of the α parameters were set to zero, some very interesting correlations and degeneracies among θ_{23} and the new




FIG. 2: Expected frequentist allowed regions at the 1σ , 90% and 2σ CL for DUNE. All new physics parameters are assumed to be zero so as to obtain the expected sensitivities. The left panels (ND averaged) correspond to the non-unitarity case, or to the sterile case when the light-heavy oscillations are averaged out in the near and far detectors. The right panels (ND undeveloped) give the sensitivity for the sterile case when the light-heavy oscillations have not yet developed in the near detector, but are averaged out in the far. The solid lines correspond to the analysis of DUNE data alone, while the dashed lines include the present constraints on sterile neutrino mixing from the right column in Tab. I.

physics parameters have been recovered. These are shown in Fig. 2, and have been noticed in the context of NSI⁵ in Refs. [59–62]. The first degeneracy appears for the wrong octant of θ_{23} , which would otherwise be correctly determined by the interplay between the appearance and disappearance channels at DUNE (see e.g. Ref. [63]). We have checked that this degeneracy is characterized by non zero values of $\alpha_{\tau e}$ with a non-trivial phase around π . At the same time, positive values of $\alpha_{\mu\mu} - \alpha_{\tau\tau}$ are slightly preferred. From Ref. [62] this degeneracy was expected for the phase of $\phi_{\tau e} = \arg(\alpha_{\tau e}) \sim \pi$ since $\delta_{\text{CP}} = -\pi/2$ and strong correlations between these two parameters are required in order to reproduce this degeneracy. Note that this degeneracy is partially lifted in the ND undeveloped scenario (right panels). Indeed, the strong sensitivity that this scenario presents to $\alpha_{\mu\mu}$ translates into very stringent bounds that do not allow the preferred positive values of $\alpha_{\mu\mu} - \alpha_{\tau\tau}$ seen in the left panels for the ND averaged case since the diagonal elements of α are positive (see Eq. (18)).

Interestingly, the second degeneracy involves values of $\theta_{23} \sim \pi/4$, so that it could potentially compromise the capabilities of DUNE to determine the maximality of this mixing angle. This degeneracy takes place for $\alpha_{\mu\mu} - \alpha_{\tau\tau} \sim -0.6$, and large values of $\alpha_{\mu e}$ and $\alpha_{\tau e}$ are also needed. Fortunately, present constraints on these parameters are already strong enough to rule out this possibility (see Tab. I), so that a clean determination of the maximality of θ_{23} should be possible at DUNE. Indeed, when the current bounds from the right column in Tab. I are added as priors to the simulations, the sensitivity to $\alpha_{\mu e}$ is considerably increased and the allowed region around $\theta_{23} \sim \pi/4$ is ruled out. This example shows explicitly the complementarity between current constraints and DUNE sensitivities.

All in all, we find that the combination of DUNE with present constraints would improve over current constraints for $\alpha_{\mu e}$ as well as for the real part of $\alpha_{\tau\mu}$. In the ND undeveloped case the sensitivity to α_{ee} is comparable while that to $\alpha_{\mu\mu}$ improves over the present constraints.

VII. CONCLUSIONS

The simplest and most natural extension of the Standard Model that can account for our present evidence for neutrino masses and mixings is the addition of right-handed neutrinos to the Standard Model (SM) particle content. Gauge and Lorentz invariance then imply the possible existence of a Majorana mass for these new particles at a scale to be determined by observations. In this work we have studied the impact that two limiting regimes for this new physics scale can have in neutrino oscillation experiments. For very high Majorana masses, beyond the kinematic reach of our experiments, the imprint of these new degrees of freedom at low energies takes the form of unitarity deviations of the PMNS mixing matrix. In the opposite limit, for small Majorana masses, these extra sterile neutrinos are produced and can participate in neutrino oscillations. However, it should be kept in mind that the neutrino oscillation phenomenology discussed here applies to any theory beyond the SM inducing unitarity deviations for the PMNS mixing matrix. This includes any model in which heavy fermions mix with the SM neutrinos or charged leptons, as for instance the type-I/type-III seesaw, Left-Right symmetric models, and models with kinematically accessible sterile neutrinos such as those invoked to explain the LSND or reactor anomalies.

⁵ Note the correspondence between NSI, steriles, and non-unitarity presented in Section IV.

Despite being sourced by different underlying physics, we have seen that, when the sterile neutrino oscillations are averaged out (and at leading order in the small heavy-active mixing angles) both limits lead to the same modifications in the neutrino oscillation probabilities. Namely, a modification of the interactions in the source and detector which implies short-distance effects as well as modified matter effects which, contrary to the standard scenario, also involve neutral current interactions. However, the present constraints that apply to these two scenarios are very different. Indeed, PMNS non-unitarity is bounded at the per mille level, or even better for some elements, through precision electroweak and flavour observables, while sterile neutrino mixing in the averaged-out regime is allowed at the percent level since it can only be probed via oscillation experiments themselves. Thus, PMNS non-unitarity can have no impact in present or near-future oscillation facilities while sterile neutrino mixing could potentially be discovered by them.

We have also noted apparently conflicting results depending on the parametrization used to encode these new physics effects. The source of this apparent inconsistency was found to be the different quantities that are commonly identified with the standard PMNS matrix in each parametrization. The conflict was solved by providing a mapping between the two sets of parameters and by identifying the parametrization for which these PMNS parameters correspond to what is determined experimentally.

The role of the near detector was also explored in depth. Indeed, since present and near future oscillation experiments constrain their fluxes and detection cross sections using near detector data it is important to consider if the new physics affects the near and far detector measurements in the same way. If this is the case, the source and detector short-distance effects cancel to a large extent, since there is no additional handle to separate them from flux and cross-section uncertainties. This is always the case in the non-unitarity scenario and when sterile neutrino oscillations are averaged out both at the near and far detectors. Conversely, if sterile neutrino oscillations have not developed yet at the near detector, the determination of the flux and cross section is free from new physics ambiguities and, when compared with the far detector data, a greater sensitivity to the new physics is obtained. This crucial difference is sometimes overlooked in the present literature. Finally we also provided a mapping of these new physics effects in the popular non-standard interaction (NSI) formalism.

These effects were exemplified through numerical simulations of the proposed DUNE neutrino oscillation experiment. Our simulations confirm that PMNS non-unitarity is indeed beyond the reach of high precision experiments such as DUNE, but that sterile neutrino oscillations could manifest in several possible interesting ways. Indeed, degeneracies between θ_{23} and the new physics parameters, previously identified in the context of NSI, have been found in our simulations. These degeneracies could potentially compromise the capability of DUNE to determine the maximality of θ_{23} as well as its ability to discern its correct octant. We find that current bounds on the new physics parameters are able to lift the degeneracies around $\theta_{23} \sim \pi/4$, although the octant degeneracies would remain at the 2σ level for the value of θ_{23} analysed in this work.

Through these simulations the importance of correctly accounting for the impact of the near detector was made evident. Indeed, a very significant increase in the sensitivity to the new physics parameters was found for the case in which the near detector is not affected in the same way as the far. This would be the case of sterile neutrino oscillations that are

undeveloped at the near detector but averaged out at the far. However, the parameter space for this situation to take place is rather small (for $\Delta m^2 \sim 0.1 \text{ eV}^2$). The most common situation would rather be that in which sterile neutrino oscillations are averaged out at both near and far detectors. However, this fact has been usually overlooked in previous literature.

The origin of neutrino masses remains one of our best windows to explore the new physics underlying the open problems of the SM. Its simplest extension to accommodate neutrino masses and mixings offers a multitude of phenomenological consequences that vary depending of the new physics scale introduced and that should be thoroughly explored by future searches. In this work we have seen that neutrino oscillation facilities can probe some parts of this largely unexplored parameter space that holds the answer to fundamental open puzzles.

Acknowledgements

The authors acknowledge support from the EU through grants H2020-MSCA-ITN-2015/674896-Elusives and H2020-MSCA-RISE-2015/690575-InvisiblesPlus. This work was supported by the Göran Gustafsson Foundation (MB). EFM and JHG also acknowledge support from the EU FP7 Marie Curie Actions CIG NeuProbes (PCIG11-GA-2012-321582) and the Spanish MINECO through the “Ramón y Cajal” programme (RYC2011-07710), the project FPA2012-31880 and through the Centro de Excelencia Severo Ochoa Program under grant SEV-2012-0249. Fermilab is operated by Fermi Research Alliance, LLC under Contract No. DE-AC02-07CH11359 with the United States Department of Energy. EFM and JHG also warmly thank Fermilab for its hospitality during their InvisiblesPlus secondment where this project was initiated.

-
- [1] M. Fukugita and T. Yanagida, Phys. Lett. **B174**, 45 (1986).
 - [2] P. Minkowski, Phys. Lett. **B67**, 421 (1977).
 - [3] R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980).
 - [4] T. Yanagida (1979), in Proceedings of the Workshop on the Baryon Number of the Universe and Unified Theories, Tsukuba, Japan.
 - [5] M. Gell-Mann, P. Ramond, and R. Slansky, *Complex Spinors and Unified Theories*, Print-80-0576 (CERN).
 - [6] J. A. Casas, J. R. Espinosa, and I. Hidalgo, JHEP **11**, 057 (2004), hep-ph/0410298.
 - [7] R. Mohapatra and J. Valle, Phys.Rev. **D34**, 1642 (1986).
 - [8] J. Bernabeu, A. Santamaria, J. Vidal, A. Mendez, and J. Valle, Phys.Lett. **B187**, 303 (1987).
 - [9] G. C. Branco, W. Grimus, and L. Lavoura, Nucl. Phys. **B312**, 492 (1989).
 - [10] W. Buchmuller and D. Wyler, Phys.Lett. **B249**, 458 (1990).
 - [11] A. Pilaftsis, Z. Phys. **C55**, 275 (1992), hep-ph/9901206.
 - [12] J. Kersten and A. Y. Smirnov, Phys.Rev. **D76**, 073005 (2007), 0705.3221.
 - [13] A. Abada, C. Biggio, F. Bonnet, M. B. Gavela, and T. Hambye, JHEP **12**, 061 (2007), 0707.4058.
 - [14] S. Antusch, C. Biggio, E. Fernandez-Martinez, M. Gavela, and J. Lopez-Pavon, JHEP **0610**, 084 (2006), hep-ph/0607020.

- [15] A. Broncano, M. B. Gavela, and E. E. Jenkins, Phys. Lett. **B552**, 177 (2003), hep-ph/0210271.
- [16] E. Fernandez-Martinez, M. B. Gavela, J. Lopez-Pavon, and O. Yasuda, Phys. Lett. **B649**, 427 (2007), hep-ph/0703098.
- [17] Z.-z. Xing, Phys. Lett. **B660**, 515 (2008), 0709.2220.
- [18] Z.-z. Xing, Phys. Rev. **D85**, 013008 (2012), 1110.0083.
- [19] F. J. Escrihuela, D. V. Forero, O. G. Miranda, M. Tortola, and J. W. F. Valle, Phys. Rev. **D92**, 053009 (2015), 1503.08879.
- [20] Y.-F. Li and S. Luo, Phys. Rev. **D93**, 033008 (2016), 1508.00052.
- [21] S. Antusch, M. Blennow, E. Fernandez-Martinez, and J. Lopez-Pavon, Phys. Rev. **D80**, 033002 (2009), 0903.3986.
- [22] S. Antusch, J. P. Baumann, and E. Fernandez-Martinez, Nucl.Phys. **B810**, 369 (2009), 0807.1003.
- [23] R. E. Shrock, Phys. Lett. **B96**, 159 (1980).
- [24] J. Schechter and J. W. F. Valle, Phys. Rev. **D22**, 2227 (1980).
- [25] R. E. Shrock, Phys. Rev. **D24**, 1232 (1981).
- [26] R. E. Shrock, Phys. Rev. **D24**, 1275 (1981).
- [27] P. Langacker and D. London, Phys.Rev. **D38**, 886 (1988).
- [28] S. M. Bilenky and C. Giunti, Phys.Lett. **B300**, 137 (1993), hep-ph/9211269.
- [29] E. Nardi, E. Roulet, and D. Tommasini, Phys.Lett. **B327**, 319 (1994), hep-ph/9402224.
- [30] D. Tommasini, G. Barenboim, J. Bernabeu, and C. Jarlskog, Nucl.Phys. **B444**, 451 (1995), hep-ph/9503228.
- [31] C. Biggio, Phys. Lett. **B668**, 378 (2008), 0806.2558.
- [32] R. Alonso, M. Dhen, M. Gavela, and T. Hambye, JHEP **1301**, 118 (2013), 1209.2679.
- [33] S. Antusch and O. Fischer, JHEP **1410**, 94 (2014), 1407.6607.
- [34] A. Abada and T. Toma, JHEP **02**, 174 (2016), 1511.03265.
- [35] E. Fernandez-Martinez, J. Hernandez-Garcia, and J. Lopez-Pavon, JHEP **08**, 033 (2016), 1605.08774.
- [36] J. Kopp, P. A. N. Machado, M. Maltoni, and T. Schwetz, JHEP **05**, 050 (2013), 1303.3011.
- [37] S. Parke and M. Ross-Lonergan (2015), 1508.05095.
- [38] D. Dutta, R. Gandhi, B. Kayser, M. Masud, and S. Prakash (2016), 1607.02152.
- [39] D. V. Forero, S. Morisi, M. Tortola, and J. W. F. Valle, JHEP **09**, 142 (2011), 1107.6009.
- [40] Y. Declais, J. Favier, A. Metref, H. Pessard, B. Achkar, et al., Nucl.Phys. **B434**, 503 (1995).
- [41] F. P. An et al. (Daya Bay), Phys. Rev. Lett. **113**, 141802 (2014), 1407.7259.
- [42] K. Abe et al. (Super-Kamiokande), Phys. Rev. **D91**, 052019 (2015), 1410.2008.
- [43] P. Adamson et al. (MINOS), Phys. Rev. Lett. **107**, 011802 (2011), 1104.3922.
- [44] G. H. Collin, C. A. Argelles, J. M. Conrad, and M. H. Shaevitz (2016), 1607.00011.
- [45] F. Dydak et al., Phys. Lett. **B134**, 281 (1984).
- [46] M. G. Aartsen et al. (IceCube), Phys. Rev. Lett. **117**, 071801 (2016), 1605.01990.
- [47] P. Adamson et al. (MINOS), Submitted to: Phys. Rev. Lett. (2016), 1607.01176.
- [48] P. Huber, M. Lindner, and W. Winter, Comput. Phys. Commun. **167**, 195 (2005), hep-ph/0407333.
- [49] P. Huber, J. Kopp, M. Lindner, M. Rolinec, and W. Winter, Comput. Phys. Commun. **177**, 432 (2007), hep-ph/0701187.
- [50] T. Alion et al. (DUNE) (2016), 1606.09550.

- [51] M. Blennow and E. Fernandez-Martinez, *Comput. Phys. Commun.* **181**, 227 (2010), 0903.3985.
- [52] M. Gonzalez-Garcia, M. Maltoni, and T. Schwetz, *JHEP* **1411**, 052 (2014), 1409.5439.
- [53] A. M. Dziewonski and D. L. Anderson, *Phys. Earth Planet. Interiors* **25**, 297 (1981).
- [54] O. G. Miranda, M. Tortola, and J. W. F. Valle (2016), 1604.05690.
- [55] J. M. Berryman, A. de Gouvea, K. J. Kelly, and A. Kobach, *Phys. Rev.* **D92**, 073012 (2015), 1507.03986.
- [56] S. K. Agarwalla, S. S. Chatterjee, and A. Palazzo (2016), 1605.04299.
- [57] S. Verma and S. Bhardwaj (2016), 1609.06412.
- [58] D. Dutta, P. Ghoshal, and S. Roy (2016), 1609.07094.
- [59] P. Coloma, *JHEP* **03**, 016 (2016), 1511.06357.
- [60] A. de Gouvea and K. J. Kelly, *Nucl. Phys.* **B908**, 318 (2016), 1511.05562.
- [61] M. Blennow, S. Choubey, T. Ohlsson, D. Pramanik, and S. K. Raut, *JHEP* **08**, 090 (2016), 1606.08851.
- [62] S. K. Agarwalla, S. S. Chatterjee, and A. Palazzo (2016), 1607.01745.
- [63] V. De Romeri, E. Fernandez-Martinez, and M. Sorel (2016), 1607.00293.