




Electrons for Neutrinos




This document was prepared by [MicroBooNE Collaboration] using the resources of the Fermi National Accelerator Laboratory (Fermilab), a U.S. Department of Energy, Office of Science, HEP User Facility. Fermilab is managed by Fermi Research Alliance, LLC (FRA), acting under Contract No. DE-AC02-07CH11359.


Afroditi Papadopoulos
New Perspectives 2018
Fermilab, IL
6/18/18



KamLAND, PRL 100, 221803 (2008)



KamLAND, PRL 100, 221803 (2008)



KamLAND, PRL 100, 221803 (2008)



Genie

New Strategy

Electrons for Neutrinos!

- Use *e*-scattering data to constraint *v*-data.



Ask
your
question

Why ?




- *e* and *v* share many common aspects of the interaction (isovector part).
- Beam energy and EM interaction well known.
- CLAS@JLab: Large number of *e*-scattering data in a wide phase-space.

Target	Beam energy (# triggers)		
	1.161 GeV	2.261 GeV	4.461 GeV
³ He	141	217	186
⁴ He	-	333	445
¹² C	62	238	310
⁵⁶ Fe	-	23	30
CH2	10	35	21
Empty cell	19	69	33



e2a Data Analysis Strategy

- *Select QE-like ($e, e'p$) events.*
- *Reweight by $e\text{-}N / \nu\text{-}N$ cross-section ratio.*
- *Analyze them as “neutrino data”.*
- *Study kinematic quantities and beam energy reconstruction methods.*
- *Compare to Neutrino Event Generator predictions.*
- *Identify parts in phase-space with good agreement.*



GENIE is a Universal Neutrino Event Generator




Generates

Events for

Neutrino

Interaction

Experiments





Includes:

- *Flux Drivers.*
- *Detector Geometry Analysers for standard detector descriptions.*
- *Interfaces with GEANT for feeding it with GENIE events.*



Event Selection



Analysis on ^{12}C @ $E = 2.261 \text{ GeV}$

- *1 proton with $p_p > 300 \text{ MeV}/c$.*
- *Fiducial cuts.*
- *π^0 corrections.*
- *2-proton & charged pion corrections.*
- *$Q^2 \geq 0.5 \text{ GeV}^2/c^2$*
- *$W < 2 \text{ GeV}/c^2$*
- *$|x_B - 1| < 0.2$*
- *Simulation: Only QE & MEC events.*



Phase Space







$^{12}\text{C}(\text{e},\text{e}'\text{p})$ @ $E = 2.261$ GeV

$Q^2 \geq 0.5 \text{ GeV}^2/c^2$, $|x_B - 1| < 0.2$, $W < 2 \text{ GeV}/c^2$



\blacktriangle Data \blacktriangle GENIE hN
 \blacktriangle GENIE hA \blacktriangle GENIE hN2014



$^{12}\text{C}(\text{e},\text{e}'\text{p})$ @ $E = 2.261$ GeV

$Q^2 \geq 0.5 \text{ GeV}^2/c^2$, $|x_B - 1| < 0.2$, $W < 2 \text{ GeV}/c^2$



\blacktriangle Data \blacktriangle GENIE hN
 \blacktriangle GENIE hA \blacktriangle GENIE hN2014



$^{12}\text{C}(\text{e},\text{e}'\text{p})$ @ $E = 2.261$ GeV

$Q^2 \geq 0.5 \text{ GeV}^2/c^2$, $|x_B - 1| < 0.2$, $W < 2 \text{ GeV}/c^2$



\blacktriangle Data \blacktriangle GENIE hN
 \blacktriangle GENIE hA \blacktriangle GENIE hN2014



$^{12}\text{C}(\text{e},\text{e}'\text{p})$ @ $E = 2.261$ GeV

$Q^2 \geq 0.5 \text{ GeV}^2/c^2$, $|x_B - 1| < 0.2$, $W < 2 \text{ GeV}/c^2$



\blacktriangle Data \blacktriangle GENIE hN
 \blacktriangle GENIE hA \blacktriangle GENIE hN2014



$^{12}\text{C}(\text{e},\text{e}'\text{p})$ @ $E = 2.261$ GeV

$Q^2 \geq 0.5 \text{ GeV}^2/c^2$, $|x_B - 1| < 0.2$, $W < 2 \text{ GeV}/c^2$



\blacktriangle Data \blacktriangle GENIE hN
 \blacktriangle GENIE hA \blacktriangle GENIE hN2014



$^{12}\text{C}(\text{e},\text{e}'\text{p})$ @ $E = 2.261$ GeV

$Q^2 \geq 0.5 \text{ GeV}^2/c^2$, $|x_B - 1| < 0.2$, $W < 2 \text{ GeV}/c^2$

\blacktriangle Data \blacktriangle GENIE hN
 \blacktriangle GENIE hA \blacktriangle GENIE hN2014





$^{12}\text{C}(e,e'p) @ E = 2.261 \text{ GeV}$


$Q^2 \geq 0.5 \text{ GeV}^2/c^2, |x_B - 1| < 0.2, W < 2 \text{ GeV}/c^2$

\blacktriangleleft Data + GENIE hN
 + GENIE hA + GENIE hN2014



$$\begin{matrix} \rightarrow \text{miss} & \rightarrow e' & \rightarrow p \\ P_{\perp} = P_{\perp} + P_{\perp} \end{matrix}$$



Energy Reconstruction



$$Q^2 \geq 0.5 \text{ GeV}^2/c^2, |x_B - 1| < 0.2, W < 2 \text{ GeV}/c^2$$

 ^{12}C

 ^{12}C


Calorimetric Reconstruction

(sum over all particles)

$$E_v^{\text{cal}} = T_h + E_l + BE$$

Benchmarking the GENIE Neutrino Event Generator Against Electron Scattering Data

Ongoing Efforts

- *Implementation of the MEC, Resonance and DIS into GENIE.*
- *Expand to other generators / reactions / nuclei / energies.*
- *Study impact on bias in oscillation analyses.*





**Mariana
Khachatryan
(ODU)**



**Afroditi
Papadopoulou
(MIT)**





**Adi
Ashkenazi
(MIT)**

+ Larry Weinstein (ODU),
Or Hen, Adrian Silva (MIT),
Kendall Mahn (MSU),

Steve Dytman (Pittsburg),
Eli Piasetzky, Erez Cohen (TAU)
Minerba Betancourt (FNAL)

Backup Slides

$$Q^2 \geq 0.5 \text{ GeV}^2/c^2, |x_B - 1| < 0.2, W < 2 \text{ GeV}/c^2$$

 ^{12}C

 ^{12}C


Leptonic Reconstruction

(only scattered lepton)

$$E_v^{\text{kin}} = \frac{2M\varepsilon + 2ME_1 - m_l^2}{2(M - E_1 + |k_l|\cos\theta)}$$

