

Beauty at High { Precision Sensitivity

Chris Quigg*

Theoretical Physics Department, Fermi National Accelerator LaboratoryP.O. Box 500, Batavia, Illinois 60510 USAE-mail: quigg@fnal.govORCID: 0000-0002-2728-2445

Opening lecture at Beauty2019.

FERMILAB-CONF-20-071-T

Origins of contemporary *B*-physics. Mesons with beauty and charm. Stable tetraquarks? Flavor and the problem of identity. Top matters. Electroweak symmetry breaking and the Higgs sector. Future instruments. Slides available at zenodo.3468909.

18th International Conference on B-Physics at Frontier Machines - Beauty2019 -29 September / 4 October, 2019 Ljubljana, Slovenia

*Speaker.

http://pos.sissa.it/

This manuscript has been authored by Fermi Research Alliance, LLC under Contract No. DE-AC02-07CH11359 with the U.S. Department of Energy, Office of Science, Office of High Energy Physics.



[©] Copyright owned by the author(s) under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND 4.0).

1. Introduction

As we open Beauty2019, I note with pleasure the large number of young scientists among the participants. Since many of you were not yet living when *B* physics was born, I want to begin with a short review of our Origin Story. I will next touch on two topics in hadron spectroscopy that have been particularly interesting to me recently: next steps in the investigation of the B_c spectrum and the likely existence of doubly heavy tetraquarks that are stable, or nearly stable, against strong decay. Then I will speak more generally to the future of our subject, posing questions about flavor physics, the top quark, and electroweak symmetry breaking and the Higgs sector. In anticipation of the European Strategy Update for Particle Physics [1], I will close by inviting you to consider the relative merits of future accelerator projects.

2. Origin Story

The first experimental evidence for the existence of the fifth quark came in the summer of 1977, with the discovery of a strong enhancement at 9.5 GeV in the mass spectrum of dimuons



FIG. 2. Excess of the data over the continuum fit of

Eq. (joure 19 Echopanel: Strictline under the dashed curve is the three-peak fit; the dashed curve is the two-peak fit. The solid curve is the two-peak fit; the dashed curve is the two-peak fit; the dashed curve is the two-peak fit; the dashed curve is the two-peak fit. The solid curve is the two-peak fit; the dashed curve is the two

cise form of the continuum. The first test is to (iii) There is evidence for a third peak $\Upsilon''(10.4)$

value and construction in the function of the parameter fit. The correlations of the matrix grant to prove the multiparameter fit. The correlations of the parameter fit for the construction of the parameter fit. The correlations of the parameter fit for the construction of the parameter fit. The correlations of the parameter fit for the construction of the parameter fit. The correlations of the parameter fit for the construction of the parameter fit. The correlations of the parameter fit for the construction of the parameter fit. The correlations of the parameter fit for the construction of the parameter fit. The correlations of the parameter fit for the construction of the parameter fit. The correlations of the parameter fit for the construction of the parameter fit. The correlations of the parameter fit for the construction of the parameter fit. The correlations of the parameter fit for the construction of the parameter fit. The correlations of the parameter fit for the construction of the parameter fit. The correlations of the parameter fit for the construction of the parameter fit for the parameter fit. The correlations of the parameter fit for the parameter for the paramete

< 15/6 mised to search for structures in the dilepton spectrum structures in the dilepton spectrum and the second the second the second the second se Further uncertainties in the results presented ± 0.06 GeV. (above since for the continuous the continuous the resonance of the resonance of the continuous the resonance of the reson ± 0.06 GeV is douided to have been and the section of the sect could provide reasonable departures from Eq. (1) two- and three-peak hypotheses and compares Atthough Makoto Kohayashi and Toshihide Maskawa's insight of that three generations of quarks above this thas. These issues must wait for a large with the set of ly above ~ 11 GeV. However, the primary conclu-number of the move of the device of the primary conclu-sions are independent of these uncertainties and the bound state of a new quark (4) and its an may be summarized as follows: (i) The structure antiquark (\overline{q}) .³⁻¹¹ Eichten and Gottfried³ have calcontains at least two narrow peaks: $\Upsilon(9.4)$ and culated the energy spacing to be expected from $\Upsilon'(10.0)$. (ii) The cross section for $\Upsilon(9.4)$, (Bd σ / thelpotential model used in their accounting for $dy)|_{y=0}$, is² 0.18±0.07 pb/nucleon. (The error inthe energy levels in charmonium. Their potential cludes our $\pm 25\%$ absolute normalization uncertain-· · /) 4 . () /. 1 2

$$V(r) = -\frac{\pi}{3}\alpha_s(m_q)/r + r/a^2$$

(2)

TABLE I. Resonance fit parameters. Continuum subtraction is given by Eq. (1). Errors are statistical only.

predicts line spacings and leptonic widths. The level spacings [Table III(a)] suggest that the shape of the potential may be oversimplified; we note that $M(\Upsilon') - M(\Upsilon)$ is remarkably close to $M(\psi')$ $- M(\psi)$.¹²

28 April 1980

The evidence for three narrow peaks was in accord with what Eichten & Gottfried [10] had anticipated within a Coulomb + linear potential model, in their preparations for the Cornell Electron Storage Ring Proposal in November 1976. They calculated quarkonium spectra over a range of quark masses $m_Q \gtrsim m_c$. At the 5-GeV nominal beam energy of CESR, they foresaw three narrow levels, as observed, but predicted a level spacing $E(2S) - E(1S) \approx 420$ MeV. It soon came to light that for a very general class of potentials, the number of narrow ${}^{3}S_{1} Q \bar{Q}$ levels that lie below the threshold for Zweig-allowed decay grows as $N = a \sqrt{m_0/m_c}$ [11]. Since N = 2 for charmonium, it is a general result that three or perhaps 4 narrow Υ levels should be seen, depending on the ratio of quark masses. Combining information from the J/ψ and Υ families, we would come to learn much more about the interquark potential than we could from either family alone.

Why choose $m_Q = 5$ GeV? Fermilab experiment E1A had reported an excess of events at high values of the inelasticity parameter $y = \int F$ $F \setminus F$ in the reaction $\overline{v} = M - \langle u^+ \rangle$ anything [17]

The excess was dubbed the hi expect the behavior $d\sigma(\bar{v}q)/$ (mostly) of quarks, in contra The excess events could be ex That was not to be. In an in discovery at the 1977 Europe by Jack Steinberger's report 1

Despite the vanishing of ring at DESY had plenty to st of Υ , measurements of the Υ







orbit. Although CESR energy settings were found by repeated resonance scans to be reproducible to better than 0.01% accuracy, there is at present an uncertainty in the overall calibration scale factor amounting to about 0.3%.

The resonances near 9.4 and 10.0 GeV match the Υ and Υ' observed first by Herb *et al*.¹ and

VOLUME 44, NUMBER 17

PHYSICAL REVIEW LETTERS

all signals were digitized and recorded on tape. This trigger gave an event rate of 0.3 Hz for a luminosity of $1 \ \mu b^{-1} s^{-1}$. A typical fill of CESR lasts 3 to 5 hours yielding an integrated luminosity of up to ~15 mb^{-1} . The integrated luminos ity for each run was measured by detecting and counting small-angle (40 to 80 mrad) collinear Bhabha scatters with lead-scintillator sandwich shower detectors. The long-term stability of the luminosity monitor is confirmed by the yield of large-angle Bhabha scattering events in the NaI

Because of the limited solid angle of the NaI array as used, a major fraction of the hadronic e^+e^- annihilations gave very few particles in the detector. Rather than trying to identify all hadronic events, which would result in an unaccept-able amount of background, our aim in the analy sis was to obtain a clean sample through the use of strict event-selection criteria. Fundamental in all criteria used was the identification of mini-mum-ionizing hadrons. At normal incidence, minimum-ionizing particles deposit 15 MeV in the first four NaI layers and ~68 MeV in the last layer of a single sector. In all scans one unambiguous and isolated minimum-ionizing track plus at least two other tracks or showers were required. All data were scanned by physicists and with computer programs. The acceptance criteria for data presented were determined by maximizing detection efficiency while maintaining the background level well below 10% of the continuum cross section. The overall efficiencies for detecting continuum and Υ events are, respectively, 28% and 37%. These values are of tained by use of the cross sections measured at oh DORIS^{2,3} (σ_{cont} = 3.8 nb at 9.4 GeV, σ_{Tpeak} = 18.5 nb after correcting for the difference in beam en ergy spread at CESR and DORIS). Absolute normalization was obtained by use of large-angle Bhabha-scattering data. The difference in efficiencies is due to the fact that T decays have higher multiplicity and sphericity than continuum revents.³ The actual number of T, T', and T''events detected above continuum were, respec-tively, 214, 53, and 133. From the continuum around the three T's we collected 272 events.

The major sources of background were (i) far single beam-wall and beam-gas interactions, (ii) close beam-wall interactions. (iii) close beam-gas interactions, and (iv) cosmic rays. Case (i) was trivially removed by the requirement of an isolated track. Cases (ii) and (iii) occur with very small probability of producing pene-trating hadrons at $\theta = 90^\circ \pm 30^\circ$ with 5-GeV electrons. Case (ii), which is more frequent, is also recognizable by tracks crossing azimuthal sector boundaries. Case (iv) was rejected by the re-quirement of three tracks. We point out that the minimal residual background does not affect the results presented here.

The hadronic yield is presented in Fig. 2, plotted in arbitrary units proportional to the ratio of detected events to small-angle Bhabha yield. In this way, the energy dependence (~ $1/E^2$) of the single-photon processes is removed. The hori-



1113

with those reported by Böhringer et al.⁷ On the T and T' our results agree with those from by the CloEcheanals difference on the CESR, Ref. [16]. Γ_{ee} ratio. Because of rather large uncertainties in the contribution of background processes such as τ production and two-photon collisions, we do not regard our present measurement of the continuum cross section as definitive.

Mass differences have been predicted by assuming that the Υ , Υ' , and Υ'' are the triplet 1S, 2S, and 3S states of a $b\overline{b}$ quark pair bound in a phenomenological potential, essentially the same as that responsible for the psion spectrum. When the potential is adjusted to fit masses in the psion region and earlier measurements of the

 $\Upsilon'-\Upsilon$ difference, the predictions for the $\Upsilon''-\Upsilon$

In 1981, studies by the CLEO experiment [17] at the $\Upsilon(4S)$ resonance launched *B* physics as we know it today. They observed an enhancement in the inclusive cross section for directly produced single electrons on the resonance. This discovery was evidence for a new weakly decaying particle, the *B* meson, with a semileptonic branching ratio $\mathscr{B}(B \to Xev) \approx 13\%$.

Although the Particle Data Group's summary tables for $(b\bar{b})$ mesons run to seventeen pages, a rich spectrum of levels is still to be explored—fourteen ordinary $(b\bar{b})$ states below flavor threshold, as indicated by the display in Figure 3.



Figure 3: Predicted and observed levels of the $(b\bar{b})$ quarkonium family [18]. Two-meson flavor thresholds are shown as dashed lines to the right of the figure.

And that is sure to be far from the whole story. We have encountered many states associated with charmonium that seem not to be pure $(c\bar{c})$ configurations [19], some indicated in Figure 4.



Figure 4: Some states associated with charmonium that may entail body plans beyond $(c\bar{c})$ [18]. Neutral states not identified as $J^{PC} = 1^{--}$ are labeled by *X*, charged states by *Z*.

Beginning with the famous X(3872), now designated $\chi_{c1}(3872)$ without prejudice to its composition, these above-flavor-threshold states are mostly narrow, and are seen in hadronic transitions or decays. Many lie in close proximity to two-meson thresholds, which may influence their properties, along with communication with open-flavor channels that must be considered for all states near or above threshold. Bearing in mind that the novel states are likely to be quantum-mechanical superpositions of several components, the response to "What are they?" may contain components of $(c\bar{c})$ quarkonium and of new body plans, including quarkonium hybrids $(q\bar{q}g)$ and $(qq\bar{q}\bar{q})$ states. In the latter category, we may include dimeson "molecules," tetraquark mesons, diquarkonium, and hadroquarkonium. When can we find $(b\bar{b})$ analogues in similar profusion?

Prescient papers in 1980–1981 by Carter & Sanda [20] and Bigi & Sanda [21] emphasized that CP violation might be large and observable in *B* decays.

B mesons were first reconstructed by the CLEO Collaboration in 1983 in final states containing D^0 or $D^{*\pm}$ and one or two charged pions [22]. I call your attention to the low statistics shown in Figure 5; what a contrast to the enormous samples available today! CLEO estimated the charged-*B* mass as $5270.8 \pm 2.3 \pm 2.0$ MeV and the neutral-*B* mass as $5274.2 \pm 1.9 \pm 2.0$ MeV, a few MeV lower than the current world averages, $M(B^{\pm}) = 5279.33 \pm 0.13$ MeV and $M(B^0) = 5279.64 \pm 0.13$ MeV [23]. According to the Particle Data Group, the quantum numbers I, J, P of B^{\pm} and B^0 still require confirmation.



Figure 5: Mass distribution of *B*-meson candidates from the CLEO experiment, Ref. [22]. The labels imply charge-conjugate final states as well.

In the same year, the MAC [24] and Mark II [25] experiments operating at the PEP e^+e^- collider at SLAC established an unexpectedly long *b*-hadron lifetime (see Figure 6), which implied a small value for the quark-mixing matrix element $|V_{cb}| \approx 0.05$.

In 1987, the UA1 experiment reported an excess of same-sign dimuons [26] in $\sqrt{s} = 630 \text{ GeV}$ collisions at the $S\bar{p}pS$ Collider at CERN. The excess could be taken as evidence for $B^0-\bar{B}^0$ or $B_s^0-\bar{B}_s^0$ oscillations. A golden event (Figure 7) recorded in the ARGUS detector operating at DESY's DORIS II storage ring [27] unambiguously demonstrated $B^0-\bar{B}^0$ oscillations. Statistical evidence from like-sign dilepton events and events containing one reconstructed B^0 and an additional fast charged lepton indicated that B^0 mixing was substantial.



Figure 6: Contours of equal likelihood for charm and beauty lifetimes from the Mark II Experiment [25] are shown together with the 2019 world-average lifetimes for hadrons containing c or b quarks [23].



Figure 7: An explicitly mixed $\Upsilon(4S) \rightarrow B^0 B^0$ event, with fully reconstructed B^0 -mesons, from Ref. [27].

Once we had established that the bottom quark carries charge $Q_b = -\frac{1}{3}$, it was natural to presume that it formed a weak-isospin doublet with an as yet undiscovered top quark. As experimental information accumulated about the neutral-current couplings of *b*, it became possible to back up that presumption [28]. Working within a V - A framework, we may generalize the left- and righthanded chiral couplings to $L_b \equiv 2I_{3L} - 2Q_b \sin^2 \theta_W$, $R_b \equiv 2I_{3R} - 2Q_b \sin^2 \theta_W$, where $\sin^2 \theta_W$ is the weak mixing parameter. Then we may deduce constraints on the chiral couplings, using these relations: The partial width $\Gamma(Z^0 \to b\bar{b})$ measures $(L_b^2 + R_b^2)$, the forward–backward asymmetry $A_{b\bar{b}}^{(\text{peak})}$ in the reaction $e^+e^- \to b\bar{b}$ on the Z^0 peak is sensitive to $(L_b^2 - R_b^2)/(L_b^2 + R_b^2)$, and the forwardbackward asymmetry in the γ -Z interference regime $A_{b\bar{b}}$ is proportional to $(R_b - L_b)$. A graphical representation of these constraints is given in Figure 8. The three constraints overlap in a small region consistent with the standard-model expectation $(I_{3L} = -\frac{1}{2}, I_{3R} = 0)$, indicating that the *b*-quark is indeed the lower member of a weak-isospin doublet.



Figure 8: Determination of the weak isospin of the *b* quark (after Ref. [28]).

Expectations of appreciable CP violation in B^0 decays were fulfilled in 2001 in reports from BABAR [29] (sin $2\beta \approx 0.59$) and Belle [30] (sin $2\phi_1 \equiv \sin 2\beta \approx 0.99$). CP-violating asymmetries have now been observed in dozens of decay modes, and the mixing parameter has settled to sin $2\beta =$ 0.699 ± 0.017 [31]. In 2006, the CDF Collaboration reported the definitive observation of timedependent $B_s^0 \cdot \overline{B}_s^0$ mixing, fixing the (very rapid) oscillation frequency at $\Delta m_{B_s^0} = 17.77 \pm 0.10 \pm$ $0.07 \hbar \text{ ps}^{-1}$ [32], within one standard deviation of the current world average. Five years later, CDF reported the first detection of CP violation in charmless decays of B_s [33]. Thanks to decades of experimentation—in dialogue with theory—we now have a library of highly constraining precision tests of the Cabibbo–Kobayashi–Maskawa quark-mixing paradigm [34], including the influence of penguin diagrams [35].

3. Mesons with Beauty and Charm

The B_c family of mesons has attracted theoretical interest for decades because of its intermediate position between the J/ψ and Υ families, and because the unequal c and b quark masses make it sensitive to relativistic effects [36]. Since B_c is composed of two different quarks, the annihilations into two- or three-gluon intermediate states that characterize $(c\bar{c})$ and $(b\bar{b})$ states cannot occur, so all excited states cascade to the ground state. The B_c itself decays only through weak transitions $b \rightarrow c, c \rightarrow s$, and $b\bar{c} \rightarrow W^-$. It took a decade after the early theoretical studies until the CDF Collaboration was able to announce the reconstruction of $B_c \rightarrow J/\psi \pi^{\pm}$, shown in Figure 9 [37]. The long wait meant that the reconstructed mass, $M(B_c) = 6274.9 \pm 0.8$ MeV, tested a precise prediction from lattice QCD [38].

Until recently, the only evidence reported for a $(c\bar{b})$ excited state was presented by the ATLAS Collaboration [39] in *pp* collisions at 7 and 8 TeV, in samples of 4.9 and 19.2 fb⁻¹. They observed a new state at 6842 ± 7 MeV in the $M(B_c^{\pm}\pi^{+}\pi^{-}) - M(B_c^{\pm}) - 2M(\pi^{\pm})$ mass difference, with B_c^{\pm} detected in the $J/\psi\pi^{\pm}$ mode. It was plausible to guess that ATLAS might have observed the transition $B_c^*(2S) \rightarrow B_c^*(1S)\pi^{+}\pi^{-}$, missing the low-energy photon from the subsequent $B_c^* \rightarrow B_c\gamma$ decay, and that the signal is an unresolved combination of 2^3S_1 and 2^1S_0 peaks. A search by the LHC*b* collaboration in 2 fb⁻¹ of 8-TeV *pp* data yielded no evidence for either $B_c(2S)$ state [40].



Figure 9: Invariant mass distribution of the $J/\psi\pi^{\pm}$ candidates. The broad enhancement below 6.2 GeV is attributable to partially reconstructed B_c mesons (from Ref. [37]).

The unsettled experimental situation and the advent of large new data sets motivated us to take a new look at the prospects for filling in the B_c spectrum [41]. The spectrum shown in Figure 10, calculated in a nonrelativistic potential model, indicates that 12–15 narrow $c\bar{b}$ levels might be found. We calculated hadronic and electromagnetic cascade decay rates for the expected narrow



Figure 10: Calculated $c\bar{b}$ spectrum, with (spin-singlet, spin-triplet) states on the (left [red], right [blue]) for each orbital-angular-momentum family; dashed lines mark two-body open-flavor thresholds (Ref. [41]).

states, and computed differential and integrated cross sections for the narrow B_c levels in pp collisions at the LHC. Putting all these elements together, we discussed how to unravel the 2S levels and explored how higher levels might be observed.

The CMS Collaboration [42] beat us to the arXiv, followed not long after by LHCb [43]. These experiments published striking evidence for both $B_c(2S)$ levels, in the form of well-separated peaks in the $B_c \pi^+ \pi^-$ invariant mass distribution, closely matching the theoretical template. The difference of mass differences, $\Delta_{21} \equiv [M(B_c^{*\prime}) - M(B_c')] - [M(B_c^{*}) - M(B_c)]$, sets the splitting between the peaks. We estimate -23 MeV, whereas CMS measures -29 MeV and LHCb, -31 MeV. The estimate depends on the not-yet-measured hyperfine splitting of the ground state, $[M(B_c^{*}) - M(B_c)]$, for which we assumed 54 MeV, the consensus value of modern lattice QCD calculations. We look



Figure 11: (Shifted) $B_c \pi^+ \pi^-$ invariant mass distributions reported by the CMS [42] (left panel) and LHC*b* [43] (right panel) Collaborations. The (lower, upper) peak is interpreted as $(B_c^*(2S), B_c(2S))$ since the photon energy in the $B_c^*(1S) \rightarrow B_c(1S) + \gamma$ transition goes undetected.

forward to further experimental analysis to determine the relative weights of the two peaks, which can test calculations of production and decay rates, and to studies of the $\pi^+\pi^-$ invariant mass distribution as next steps in B_c spectroscopy.

Our calculations indicate that the 3S levels will lie above flavor threshold. The 3^1S_0 state can decay into the final state B^*D and the 3^3S_1 level has decays into both the *BD* and B^*D final states. However, it is conceivable that coupled-channel effects might push one or both states lower in mass. It is therefore worth examining the $B_c \pi^+ \pi^-$ mass spectrum up through 7200 MeV for indications of $3^1S_0 \rightarrow B_c \pi^+ \pi^-$ and $3^3S_1 \rightarrow B_c^* \pi^+ \pi^-$ lines.

The $3^{3}P_{2}(c\bar{b})$ state might be observed as a very narrow (*d*-wave) *BD* line near open-flavor threshold, in the spirit of the LHC*b* candidate [44] for the charmonium level $3^{3}D_{3} \equiv \psi_{3}(3D) \rightarrow D\bar{D}$ at mass $M = 3842.72 \pm 0.16 \pm 0.12$ MeV and natural width $\Gamma = 2.79 \pm 0.51 \pm 0.35$ MeV. We anticipate that one more narrow charmonium state, $4^{3}F_{4}$, is to be found above flavor threshold, perhaps near 4054 MeV [45].

It may in time become possible for experiments to detect some of the more energetic E1transition photons that appear in $(c\bar{b})$ cascades. A particularly attractive target for experiment is the $2^{3}P_{2}(6750) \rightarrow B_{c}^{*}\gamma$ line, because of the favorable production cross section, branching fraction, and 409-MeV photon energy. The $3^{3}P_{2}(7154) \rightarrow B_{c}^{*}\gamma(777 \text{ MeV})$ line is favored for its high photon energy, which may be a decisive advantage for detection.

4. Tetraquarks stable against strong decays

The proposition that stable or nearly stable multiquark states containing heavy flavors might exist goes back nearly four decades [46]. The discovery of the doubly-charmed baryon $\Xi_{cc}^{++}(3621)$ in the LHCb experiment [47] has provoked a new wave of interest in exotic mesons containing two heavy quarks. Estia Eichten and I have examined the possibility of unconventional tetraquark configurations for which all strong decays are kinematically forbidden [48]. In the heavy-quark limit, stable—hence exceedingly narrow— $Q_i Q_j \bar{q}_k \bar{q}_l$ mesons must exist. To apply this insight, we take into account corrections for finite heavy-quark masses to deduce which tetraquark states

8

containing b or c quarks might be stable. The most promising candidate is a $J^P = 1^+$ isoscalar double-b meson, $\mathscr{T}_{[\bar{u}\bar{d}]}^{\{bb\}-}$.

Let us start with the observation that one-gluon-exchange between a pair of color-triplet heavy quarks is attractive for (QQ) in a color- $\overline{\mathbf{3}}$ configuration and repulsive for the color- $\mathbf{6}$ configuration. The strength of the $\overline{\mathbf{3}}$ attraction is half that of the corresponding $(Q\overline{Q})$ in a color- $\mathbf{1}$. This means that in the limit of very heavy quarks, we may idealize the color-antitriplet (QQ) diquark as a stationary, structureless color charge, as depicted in the leftmost panel in Figure 12. In that



Figure 12: Schematic evolution of a $Q_i Q_j \bar{q}_k \bar{q}_l$ state as the heavy-quark masses decrease (and the mean separation between the heavy quarks increases) from left to right.

case, we can separate the strong dynamics binding the diquark from the long-range color interaction by which the light antiquarks interact with each other and are bound to the diquark "nucleus." For sufficiently heavy quarks Q, a $Q_i Q_j \bar{q}_k \bar{q}_l$ tetraquark meson is stable against strong decays, as we can show by considering possible decay modes. First, we note that dissociation into two heavy–light mesons is kinematically forbidden. The \mathscr{Q} value for the decay is $\mathscr{Q} \equiv m(Q_i Q_j \bar{q}_k \bar{q}_l) - [m(Q_i \bar{q}_k) + m(Q_j \bar{q}_l)] = \Delta(q_k, q_l) - \frac{1}{2}(\frac{2}{3}\alpha_s)^2[1 + O(v^2)]\overline{M} + O(1/\overline{M})$, where $\Delta(q_k, q_l)$, the contribution due to light dynamics, becomes independent of the heavy-quark masses, $\overline{M} \equiv (1/m_{Q_i} + 1/m_{Q_j})^{-1}$ is the reduced mass of Q_i and Q_j , and α_s is the strong coupling. For large enough values of \overline{M} , the middle term on the right-hand side dominates, so the tetraquark is stable against decay into two heavy-light mesons.

What of the other possible decay channel, a doubly heavy baryon plus a light antibaryon, $(Q_i Q_j \bar{q}_k \bar{q}_l) \rightarrow (Q_i Q_j q_m) + (\bar{q}_k \bar{q}_l \bar{q}_m)$? For very heavy quarks, the contributions of Q motion and spin to the tetraquark mass are negligible. Since the (QQ) diquark is a color-antitriplet, heavyquark symmetry tells us that

$$m(Q_i Q_j \bar{q}_k \bar{q}_l) - m(Q_i Q_j q_m) = m(Q_x q_k q_l) - m(Q_x \bar{q}_m).$$
(4.1)

Taking into account finite-mass corrections prescribed by heavy-quark symmetry, we can use measured masses to show that the right-hand side is in every case smaller than the mass of the lightest antibaryon, \bar{p} , so no decay to a doubly heavy baryon and a light antibaryon is kinematically allowed. With no open channels in the heavy-quark limit, stable $Q_i Q_j \bar{q}_k \bar{q}_l$ mesons must exist.

To assess the implications for the real world, we must first test whether it makes sense to idealize the (QQ) diquark as a tiny, structureless, color-antitriplet color source. As the separation between the heavy quarks increases, the light-antiquark cloud screens the Q_iQ_j interaction, altering the $\mathbf{\bar{3}}, \mathbf{6}$ mix, and eventually leading to the fission of the $(Q_iQ_j\bar{q}_k\bar{q}_l)$ state into a pair of heavy–light mesons. These changes are indicated in the progression from left to right in Figure 12. Using a half-strength Coulomb+linear quarkonium potential, we verified that the rms core radii are small on the

expected tetraquark scale: $\langle r^2 \rangle^{1/2} = 0.19 \text{ fm}(bb)$; 0.24 fm(bc); 0.28 fm(cc). This assessment is consistent with other results [49].

If we had access to measured masses for all of the doubly heavy baryons, Eqn. (4.1), with controlled finite-mass corrections, would yield predictions of tetraquark masses, without the intervention of any models. That is not yet the case; only one example, Ξ_{cc}^{++} , can be considered established. While waiting for experiments to provide more comprehensive information, we use as inputs the model calculations of doubly-heavy baryon masses by Karliner & Rosner [50].

We find two real-world candidates for stable tetraquarks: the axial vector $\{bb\}[\bar{u}\bar{d}]$ meson, $\mathscr{T}_{[\bar{u}\bar{d}]}^{\{bb\}^-}$ bound by 121 MeV against strong decays, and the axial vector $\{bb\}[\bar{u}\bar{s}]$ and $\{bb\}[\bar{d}\bar{s}]$ mesons bound by 48 MeV. Given the provisional doubly heavy baryon masses, we expect all the other $Q_i Q_j \bar{q}_k \bar{q}_l$ tetraquarks to lie at least 78 MeV above the corresponding thresholds for strong decay. Promising final states in which to search for stable tetraquarks include $\mathscr{T}_{[\bar{u}\bar{d}]}^{\{bb\}}(10482)^- \rightarrow$ $\Xi_{bc}^0 \bar{p}, B^- D^+ \pi^-$, and $B^- D^+ \ell^- \bar{\nu}$ (which establishes a weak decay), $\mathscr{T}_{[\bar{u}\bar{s}]}^{\{bb\}}(10643)^- \rightarrow \Xi_{bc}^0 \bar{\Sigma}^-$, $\mathscr{T}_{[\bar{d}\bar{s}]}^{\{bb\}}(10643)^0 \rightarrow \Xi_{bc}^0(\bar{\Lambda}, \bar{\Sigma}^0)$, and so on. Observing a weakly decaying double-beauty state would establish the existence of tetraquarks and illuminate the role of heavy color- $\bar{\mathbf{3}}$ diquarks as hadron constituents.

Heavier $bb\bar{q}_k\bar{q}_l$ states, as well as $bc\bar{q}_k\bar{q}_l$ and $cc\bar{q}_k\bar{q}_l$ might be seen as "wrong-sign" $Q\bar{q} + Q\bar{q}$ resonances in double-flavor DD, DB, BB combinations near threshold. (In their model calculations, Karliner and Rosner [51] estimate somewhat deeper binding, and so point to additional bc and cccandidates.) For example, the double-charge, double charm $J^P = 1^+ \mathscr{T}_{[\bar{d}\bar{s}]}^{\{cc\}++}(4156) \rightarrow D^+D_s^{*+}$ would constitute prima facie evidence for a non- $q\bar{q}$ level. Other promising cases include the $1^+ \mathscr{T}_{\{\bar{q}_k\bar{q}_l\}}^{\{bb\}}(10681)^{0,-,--}, 1^+ \mathscr{T}_{[\bar{u}\bar{d}]}^{\{bc\}}(7272)^0, 0^+ \mathscr{T}_{[\bar{u}\bar{d}]}^{[bc]}(7229)^0$, and $1^+ \mathscr{T}_{[\bar{u}\bar{d}]}^{\{cc\}}(3978)^+$. None of this will be easy, but both experiment and theory have much to do along the way.

- $\mathcal{T}1$. Look for double-flavor resonances near threshold.
- $\mathscr{T}2$. Measure cross sections for final states containing 4 heavies: $Q_i \bar{Q}_i Q_j \bar{Q}_j$.
- \mathscr{T} 3. Discover and determine masses of doubly-heavy baryons. We need this information to implement the heavy-quark-symmetry calculation of tetraquark masses embodied in Eqn. (4.1). The doubly heavy baryons are also of intrinsic interest: in the heavy-quark limit, they resemble heavy–light mesons, with the added possibility of (QQ) core excitations.
- $\mathscr{T}4$. Resolve the uncertainty surrounding Ξ_{cc}^+ between LHCb [52] and SELEX [53].
- \mathscr{T} 5. Find stable tetraquarks through weak decays. A rough guess for the lifetimes is ~ 1 ps.
- $\mathcal{T}6.$ Develop expectations for production (cf. Ref. [54]).
- \mathcal{T} 7. Refine lifetime estimates for stable states.
- \mathscr{T} 8. Understand how color configurations evolve with QQ (and $\bar{q}\bar{q}$) masses.
- \mathscr{T} 9. Does it make sense to consider body plans such as $(Q_iQ_j)(Q_kQ_l)(Q_mQ_n)$ in the heavy-quark limit? This is an example of a six-quark state with baryon number B = 2, but a $Q_pQ_qQ_r$ color structure, if the three-diquark configuration should dominate.

5. Flavor: the problem of identity

The essence of the problem is easy to state, but not so easy to answer: What makes an electron an electron, a top quark a top quark, a neutrino a neutrino? We do not have a clear view of how to approach the diverse character of the constituents of matter. The CKM paradigm is an extraordinarily fruitful framework in the hadron sector, but there are many parameters. We have no clue what determines them, nor at what energy scale they are set. Even if the Higgs mechanism explains *how* masses and mixing angles arise, we do not know *why* they have the values we observe. In this sense, all fermion masses—beginning with the electron mass—cry out for physics beyond the standard model!

Our ability to calculate within the standard model rests on knowing the values of many parameters: 3 Coupling parameters, α_s , α_{em} , $\sin^2 \theta_W$; 2 Parameters of the Higgs potential; 1 Vacuum phase (QCD); 6 Quark masses, 3 Quark mixing angles, 1 CP-violating phase; 3 Charged-lepton masses, 3 Neutrino masses, 3 Leptonic mixing angles, 1 Leptonic CP-violating phase (probably plus two Majorana phases), for a total of 26⁺ seemingly arbitrary parameters. Of these, twenty concern the flavor sector. The problem of identity is rich in questions:

- F1. Can we find evidence of right-handed charged-current interactions? Is nature built on a fundamentally asymmetrical plan, or are the right-handed weak interactions simply too feeble for us to have observed until now, reflecting an underlying hidden symmetry?
- F2. What is the relationship of left-handed and right-handed fermions?
- F3. Are there additional electroweak gauge bosons, beyond W^{\pm} and Z?
- F4. Are there additional kinds of matter?
- F5. Is charged-current universality exact? What about lepton-flavor universality?
- F6. Where are flavor-changing neutral currents in quark transitions? In the standard model, these are absent at tree level and highly suppressed by the Glashow–Iliopoulos–Maiani mechanism [55]. They arise generically in proposals for physics beyond the standard model, and need to be controlled. And yet we have made no sightings! Why not? The focus for now is on studies of $B_{s,d} \rightarrow \mu^+\mu^-$ and $K^+ \rightarrow \pi^+\nu\bar{\nu}$.
- F7. How well can we test the standard-model correlation among the quark-mixing matrix parameter γ , $\mathscr{B}(K^+ \to \pi^+ v \bar{v})$, and $\mathscr{B}(B_s \to \mu^+ \mu^-)$?
- F8. Have we found the "periodic table" of elementary particles?
- F9. What do generations mean? Is there a family symmetry?
- F10. Why are there three families of quarks and leptons? (Is it so?)
- F11. Are there new species of quarks and leptons, possibly carrying exotic charges?
- F12. Is there any link to a dark sector?
- F13. What will resolve the disparate values of $|V_{ub}|$ and $|V_{cb}|$ measured in inclusive and exclusive decays?
- F14. Is the 3×3 (CKM) quark-mixing matrix unitary?
- F15. Why is isospin a good symmetry? What does it mean?
- F16. Can we find evidence for charged-lepton flavor violation in lepton decays?
- F17. Will we establish and diagnose a break in the standard model?
- F18. Do flavor parameters *mean* anything at all? Contrast the landscape perspective.
- F19. If flavor parameters have meaning (beyond engineering information), what is the metaquestion?

6. Top matters

The top quark touches many topics in particle physics and presents us with many questions.

- t1. How stringently will refined measurements of M_W and m_t test the electroweak-theory prediction for M_H ?
- t2. How much can we tighten the m_t - M_W - M_H constraints?
- *t*3. Does top's large $Ht\bar{t}$ (Yukawa) coupling imply a special role in electroweak symmetry breaking? How does it influence $t\bar{t}$ dynamics? Does the large value of m_t make top an outlier or the only normal fermion?
- t4. How well can we constrain V_{tb} in single-top production, and elsewhere?
- t5. How much can we refine our knowledge of V_{td} and V_{ts} ?
- t6. How complete is our understanding of $t\bar{t}$ production in QCD: total and differential cross sections, charge asymmetry, spin correlations, etc.?
- t7. What might we learn from "dead-cone" studies using boosted tops [56]?
- *t*8. How well can we constrain the top-quark lifetime [57]? How free is *t*?
- t9. Are there (vestiges of) $t\bar{t}$ resonances?
- t10. Can we find evidence of flavor-changing top decays $t \to (Z, \gamma)(c, u)$?

7. Electroweak symmetry breaking and the Higgs sector

In a few short years, the LHC experiments have given us a wealth of information about the Higgs boson [58]. We can summarize by saying that the evidence is developing as it would if H(125 GeV) were the textbook scalar responsible for electroweak symmetry breaking and the generation of fermion masses and mixings. Here is a list of questions about electroweak symmetry breaking and the Higgs sector that we must answer to approach a final verdict about how closely H(125) matches the textbook Higgs boson [59].

- *H*1. Is H(125) the only member of its clan? Might there be others—charged or neutral—at higher or lower masses?
- H2. Does H(125) fully account for electroweak symmetry breaking? Does it match standardmodel branching fractions to gauge bosons? Are absolute couplings to W and Z as expected in the standard model?
- H3. Are all production rates as expected? Any surprise sources of H(125)?
- H4. What accounts for the immense range of fermion masses?
- *H5.* Is the Higgs field the only source of fermion masses? Are fermion couplings proportional to fermion masses? $\mu^+\mu^-$ soon? How can we detect $H \to c\bar{c}$? What about $H \to e^+e^-$, which would give new insight into the finiteness of the Bohr radius and the origins of valence bonding?
- H6. What role does the Higgs field play in generating neutrino masses?
- *H*7. Can we establish or exclude decays to new particles? Does H(125) act as a portal to hidden sectors? When can we measure Γ_H ?
- *H*8. Can we detect flavor-violating decays $(\tau^{\pm}\mu^{\mp},...)$?
- H9. Do loop-induced decays $(gg, \gamma\gamma, \gamma Z)$ occur at standard-model rates?

- *H*10. What can we learn from rare decays $(J/\psi \gamma, \Upsilon \gamma, ...)$?
- H11. Does the electroweak vacuum seem stable, or suggest a new physics scale?
- H12. Can we find signs of new strong dynamics or (partial) compositeness?
- H13. Can we establish the HHH trilinear self-coupling?
- *H*14. How well can we test the notion that *H* regulates Higgs–Goldstone scattering, i.e., tames the high-energy behavior of *WW* scattering?
- H15. Is the electroweak phase transition first-order?

8. Future instruments

Our experimental future demands both diversity and scale diversity, but an important driver of progress will be the next great accelerator. Every one of us should take time to explore the possibilities, form an opinion, and communicate it to our colleagues in particle physics and beyond. I have given an inventory of frontier machines in a recent essay [60] and series of seminars [61], where you may find pointers to detailed information. Now, the great question: How do *you* assess the scientific potential *for Beauty and in general* of

- (a) The High-Luminosity LHC?
- (b) The High-Energy LHC?
- (c) A 100-TeV *pp* Collider (FCC-hh or SppC)?
- (d) A 250-GeV ILC?
- (e) A circular Higgs factory (FCC-ee or CEPC)?
- (f) A 380-GeV CLIC?
- (g) A $\mu^+\mu^- \rightarrow H$ Higgs factory?
- (h) LHeC / FCC-eh? (or an electron-ion collider?)
- (i) A muon-storage-ring neutrino factory?
- (j) A multi-TeV muon collider?
- (k) The instrument of your dreams?

Acknowledgments

I thank the Beauty2019 Organizing Committee, particularly Robert Fleischer and Guy Wilkinson, for their kind invitation to present the opening lecture. Congratulations to our Ljubljana hosts, led by Bostjan Golob, for their flawless preparations and peerless hospitality. I join all participants in looking forward to Beauty2020, where we anticipate the first wave of results from Belle II.

This work was supported by Fermi Research Alliance, LLC under Contract No. DE-AC02-07CH11359 with the U.S. Department of Energy, Office of Science, Office of High Energy Physics, and by the Munich Institute for Astro- and Particle Physics (MIAPP), which is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC-2094 – 390783311. I am grateful to the Visiting Professor Program of the Bavarian State Ministry for Science, Research, and the Arts and also the Institute for Advanced Study at Technical University Munich where much of this talk was prepared. I thank Estia Eichten for a long and enlightening collaboration.

References

- [1] Information about the European Strategy for Particle Physics may be found at https://europeanstrategyupdate.web.cern.ch. See in particular R. K. Ellis *et al.*, "Physics Briefing Book : Input for the European Strategy for Particle Physics Update 2020," arXiv:1910.11775.
- [2] S. W. Herb *et al.*, "Observation of a Dimuon Resonance at 9.5-GeV in 400-GeV Proton-Nucleus Collisions," Phys. Rev. Lett. **39** (1977) 252. Leon Lederman's announcement at the 1977 European Physical Society conference in Budapest is recounted in "The discovery of Upsilon," CERN Courier **17** number 7–8 (July–August 1977) 223.
- [3] W. R. Innes *et al.*, "Observation of Structure in the Υ Region," Phys. Rev. Lett. **39** (1977) 1240; Erratum: [Phys. Rev. Lett. **39** (1977) 1640].
- [4] J. A. Appel *et al.*, "A Study of Di-Lepton Production in Proton Collisions at NAL," NAL Proposal #288 (February 1974). For color commentary, see T. Toohig, "Leon and the Lady," CERN *Courier* 17 Number 1–2 (1977) 25.
- [5] J. J. Aubert *et al.*, "Experimental Observation of a Heavy Particle *J*," Phys. Rev. Lett. 33 (1974) 1404. For a perspective, see S. C. C. Ting, "The Discovery of the *J* Particle: A Personal Recollection," Rev. Mod. Phys. 49 (1977) 235.
- [6] J. E. Augustin *et al.* [SLAC-SP-017 Collaboration], "Discovery of a Narrow Resonance in e⁺e⁻ Annihilation," Phys. Rev. Lett. **33** (1974) 1406. See also G. S. Abrams *et al.*, "The Discovery of a Second Narrow Resonance in e⁺e⁻ Annihilation," Phys. Rev. Lett. **33** (1974) 1453. For additional perspective, consult B. Richter, "From the psi to Charm: The Experiments of 1975 and 1976," Rev. Mod. Phys. **49** (1977) 251.
- [7] M. L. Perl *et al.*, "Evidence for Anomalous Lepton Production in e⁺e⁻ Annihilation," Phys. Rev. Lett. 35 (1975) 1489. See also M. L. Perl, "Reflections on the Discovery of the Tau Lepton," in G. Ekspong (ed.), *Nobel lectures in physics 1991-1995* (World Scientific, Singapore, 1997), pp. 168–195.
- [8] "FERMILAB Dileptomania," CERN Courier 17 Number 1-2 (1977) 3-5.
- M. Kobayashi and T. Maskawa, "CP Violation in the Renormalizable Theory of Weak Interaction," Prog. Theor. Phys. 49 (1973) 652.
- [10] E. Eichten and K. Gottfried, "Heavy Quarks in e^+e^- Annihilation," Phys. Lett. **66B** (1977) 286.
- [11] C. Quigg and J. L. Rosner, "Counting Narrow Levels of Quarkonium," Phys. Lett. 72B (1978) 462.
- [12] A. C. Benvenuti *et al.*, "Further Data on the High y Anomaly in Inelastic anti-neutrino Scattering," Phys. Rev. Lett. **36** (1976) 1478.
- [13] R. M. Barnett, "Evidence in Neutrino Scattering for Righthanded Currents Associated with Heavy Quarks," Phys. Rev. D 14 (1976) 70.
- [14] M. Holder *et al.*, "Is There a High y Anomaly in anti-neutrino Interactions?" Phys. Rev. Lett. **39** (1977) 433.
- [15] See the discussion surrounding Figure 3 of J. D. Jackson, C. Quigg and J. L. Rosner, "New Particles, Theoretical," in S. Homma, M. Kawaguchi and H. Miyazawa, *High energy physics. Proceedings, 19th International Conference, ICHEP 1978, Tokyo,* pp. 391–408: Conf. Proc. C 780823 (1978) 391.
- [16] D. Andrews *et al.* [CLEO Collaboration], "Observation of Three Upsilon States," Phys. Rev. Lett. 44 (1980) 1108; T. Bohringer *et al.*, "Observation of Υ, Υ', and Υ" at the Cornell Electron Storage Ring," Phys. Rev. Lett. 44 (1980) 1111.

- [17] C. Bebek *et al.* [CLEO Collaboration], "Evidence for New Flavor Production at the $\Upsilon(4S)$," Phys. Rev. Lett. **46** (1981) 84.
- [18] Estia J. Eichten, private communication.
- [19] For a recent review containing extensive lists of nonstandard hadron candidates, see S. L. Olsen,
 T. Skwarnicki and D. Zieminska, "Nonstandard heavy mesons and baryons: Experimental evidence," Rev. Mod. Phys. 90 (2018) 015003 [arXiv:1708.04012].
- [20] A. B. Carter and A. I. Sanda, "CP Violation in Cascade Decays of *B* Mesons," Phys. Rev. Lett. 45 (1980) 952;
- [21] I. I. Y. Bigi and A. I. Sanda, "Notes on the Observability of CP Violations in *B* Decays," Nucl. Phys. B 193 (1981) 85.
- [22] S. Behrends *et al.* [CLEO Collaboration], "Observation of Exclusive Decay Modes of *B* Flavored Mesons," Phys. Rev. Lett. **50** (1983) 881.
- [23] M. Tanabashi *et al.* [Particle Data Group], "Review of Particle Physics," Phys. Rev. D 98 (2018) 030001, and 2019 update.
- [24] E. Fernandez et al., "Lifetime of Particles Containing B Quarks," Phys. Rev. Lett. 51 (1983) 1022.
- [25] N. Lockyer *et al.*, "Measurement of the Lifetime of Bottom Hadrons," Phys. Rev. Lett. **51** (1983) 1316.
- [26] C. Albajar *et al.* [UA1 Collaboration], "Search for $B^0 \overline{B}^0$ Oscillations at the CERN Proton anti-Proton Collider. 2.," Phys. Lett. B **186** (1987) 247, Erratum: [Phys. Lett. B **197** (1987) 565].
- [27] H. Albrecht *et al.* [ARGUS Collaboration], "Observation of $B^0 \overline{B}^0$ Mixing," Phys. Lett. B **192** (1987) 245.
- [28] D. Schaile and P. M. Zerwas, "Measuring the weak isospin of b quarks," Phys. Rev. D 45 (1992) 3262.
- [29] B. Aubert *et al.* [BaBar Collaboration], "Observation of CP violation in the B⁰ meson system," Phys. Rev. Lett. 87 (2001) 091801 [arXiv:hep-ex/0107013].
- [30] K. Abe *et al.* [Belle Collaboration], "Observation of large CP violation in the neutral *B* meson system," Phys. Rev. Lett. 87 (2001) 091802 [arXiv:hep-ex/0107061].
- [31] Y. S. Amhis *et al.* [HFLAV Collaboration], "Averages of *b*-hadron, *c*-hadron, and *τ*-lepton properties as of 2018," arXiv:1909.12524.
- [32] A. Abulencia *et al.* [CDF Collaboration], "Observation of $B_s^0 \bar{B}_s^0$ Oscillations," Phys. Rev. Lett. **97** (2006) 242003 [arXiv:hep-ex/0609040].
- [33] T. Aaltonen *et al.* [CDF Collaboration], "Measurements of Direct CP Violating Asymmetries in Charmless Decays of Strange Bottom Mesons and Bottom Baryons," Phys. Rev. Lett. **106** (2011) 181802 [arxiv:1103.5762].
- [34] For recent compilations and critical evaluations, see the websites of the UTfit Collaboration: M. Bona *et al.*, "The Unitarity Triangle Fit in the Standard Model and Hadronic Parameters from Lattice QCD: A Reappraisal after the Measurements of Δm_s and ℬ(B → τν_τ)," JHEP 0610 (2006) 081 [arXiv:hep-ph/0606167]; and CKMfitter Group: J. Charles *et al.*, "CP violation and the CKM matrix: Assessing the impact of the asymmetric *B* factories," Eur. Phys. J. C 41 (2005) 1 [arXiv:hep-ph/0406184]. For perspective, see M. Kobayashi, "Nobel Lecture: CP violation and flavor mixing," Int. J. Mod. Phys. A 24 (2009) 2379 [Rev. Mod. Phys. 81 (2009) 1019]; T. Maskawa, "Nobel Lecture: What does CP violation tell us?" Int. J. Mod. Phys. A 24 (2009) 2393 [Rev. Mod. Phys. 81

(2009) 1027]. And keep in mind N. Cabibbo, "Unitary Symmetry and Leptonic Decays," Phys. Rev. Lett. **10** (1963) 531.

- [35] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, "Light Quarks and the Origin of the $\Delta I = 1/2$ Rule in the Nonleptonic Decays of Strange Particles," Nucl. Phys. B **120** (1977) 316. The origin of the penguin monicker is the stuff of legend, recounted (for example) in §1.1 of M. A. Shifman, "Foreword to ITEP lectures in particle physics," in *ITEP Lectures in Particle Physics and Field Theory*, edited by M. Shifman (World Scientific, Singapore, 1999) Vol. 1, pp. v–xi [arXiv:hep-ph/9510397].
- [36] My own interest dates back to E. J. Eichten and C. Quigg, "Mesons with beauty and charm: Spectroscopy," Phys. Rev. D 49 (1994) 5845 [arXiv:hep-ph/9402210].
- [37] A. Abulencia *et al.* [CDF Collaboration], "Evidence for the exclusive decay $B_c^{\pm} \rightarrow J/\psi \pi^{\pm}$ and measurement of the mass of the B_c meson," Phys. Rev. Lett. **96** (2006) 082002 [arXiv:hep-ex/0505076].
- [38] I. F. Allison *et al.* [HPQCD and Fermilab Lattice and UKQCD Collaborations], "Mass of the B_c meson in three-flavor lattice QCD," Phys. Rev. Lett. 94 (2005) 172001 [arXiv:hep-lat/0411027].
- [39] G. Aad *et al.* [ATLAS Collaboration], "Observation of an Excited B_c^{\pm} Meson State with the ATLAS Detector," Phys. Rev. Lett. **113** (2014) 212004 [arXiv:1407.1032].
- [40] R. Aaij *et al.* [LHCb Collaboration], "Search for excited B_c^+ states," JHEP **1801** (2018) 138 [arXiv:1712.04094].
- [41] E. J. Eichten and C. Quigg, "Mesons with Beauty and Charm: New Horizons in Spectroscopy," Phys. Rev. D 99 (2019) 054025 [arXiv:1902.09735].
- [42] A. M. Sirunyan *et al.* [CMS Collaboration], "Observation of Two Excited B_c^+ States and Measurement of the $B_c^+(2S)$ Mass in *pp* Collisions at $\sqrt{s} = 13$ TeV," Phys. Rev. Lett. **122** (2019) 132001 [arXiv:1902.00571].
- [43] R. Aaij *et al.* [LHCb Collaboration], "Observation of an excited B_c^+ state," Phys. Rev. Lett. **122** (2019) 232001 [arXiv:1904.00081].
- [44] R. Aaij et al. [LHCb Collaboration], "Near-threshold DD spectroscopy and observation of a new charmonium state," JHEP 1907 (2019) 035 [arXiv:1903.12240].
- [45] E. J. Eichten, K. Lane and C. Quigg, "Charmonium levels near threshold and the narrow state $X(3872) \rightarrow \pi^+\pi^- J/\psi$," Phys. Rev. D **69** (2004) 094019 [arXiv:hep-ph/0401210].
- [46] J. P. Ader, J. M. Richard and P. Taxil, "Do Narrow Heavy Multi Quark States Exist?," Phys. Rev. D 25 (1982) 2370. An exemplary bibliography appears in J. M. Richard, A. Valcarce and J. Vijande, "Few-body quark dynamics for doubly heavy baryons and tetraquarks," Phys. Rev. C 97 (2018) 035211 [arXiv:1803.06155].
- [47] R. Aaij *et al.* [LHCb Collaboration], "Observation of the doubly charmed baryon Ξ_{cc}^{++} ," Phys. Rev. Lett. **119** (2017) 112001 [arXiv:1707.01621].
- [48] E. J. Eichten and C. Quigg, "Heavy-quark symmetry implies stable heavy tetraquark mesons $Q_i Q_j \bar{q}_k \bar{q}_l$," Phys. Rev. Lett. **119** (2017) 202002 [arXiv:1707.0957]
- [49] For investigations using a variety of techniques, see Richard–Valcarce–Vijande, in Ref. [46];
 A. Czarnecki, B. Leng and M. B. Voloshin, "Stability of tetrons," Phys. Lett. B 778 (2018) 233
 [arXiv:1708.04594]; C. Hughes, E. Eichten and C. T. H. Davies, "Searching for beauty-fully bound tetraquarks using lattice nonrelativistic QCD," Phys. Rev. D 97 (2018) 054505 [arXiv:1710.03236].
 P. Bicudo, M. Cardoso, A. Peters, M. Pflaumer and M. Wagner, "Doubly heavy tetraquark resonances in lattice QCD," J. Phys. Conf. Ser. 1137 (2019) 012039.

- [50] M. Karliner and J. L. Rosner, "Baryons with two heavy quarks: Masses, production, decays, and detection," Phys. Rev. D 90 (2014) 094007 [arXiv:1408.5877].
- [51] M. Karliner and J. L. Rosner, "Discovery of doubly-charmed Ξ_{cc} baryon implies a stable ($bb\bar{u}\bar{d}$) tetraquark," Phys. Rev. Lett. **119** (2017) 202001 [arXiv:1707.07666].
- [52] R. Aaij *et al.* [LHCb Collaboration], "Search for the doubly charmed baryon \(\mathbb{Z}_{cc}\)," Sci. China Phys. Mech. Astron. **63** (2020) 221062 [arXiv:1909.12273].
- [53] M. Mattson *et al.* [SELEX Collaboration], "First Observation of the Doubly Charmed Baryon Ξ_{cc}^+ ," Phys. Rev. Lett. **89** (2002) 112001 [arXiv:hep-ex/0208014].
- [54] A. Ali, A. Y. Parkhomenko, Q. Qin and W. Wang, "Prospects of discovering stable double-heavy tetraquarks at a Tera-Z factory," Phys. Lett. B **782** (2018) 412 [arXiv:1805.02535].
- [55] S. L. Glashow, J. Iliopoulos and L. Maiani, "Weak Interactions with Lepton-Hadron Symmetry," Phys. Rev. D 2 (1970) 1285.
- [56] Y. L. Dokshitzer, V. A. Khoze and S. I. Troian, "On specific QCD properties of heavy quark fragmentation ('dead cone')," J. Phys. G 17 (1991) 1602; F. Maltoni, M. Selvaggi and J. Thaler, "Exposing the dead cone effect with jet substructure techniques," Phys. Rev. D 94 (2016) 054015 [arXiv:1606.03449].
- [57] A recent measurement yields $\Gamma(t) = 1.9 \pm 0.5$ GeV, to be compared with the standard-model expectation of 1.32 GeV. See The ATLAS collaboration, "Measurement of the top-quark decay width in top-quark pair events in the dilepton channel at $\sqrt{s} = 13$ TeV with the ATLAS detector," ATLAS-CONF-2019-038.
- [58] For compilations of recent results on the Higgs boson H(125), see The ATLAS Collaboration website, "Public Results: Higgs Physics" and The CMS Collaboration website, "CMS Higgs Physics Results."
- [59] For an instructive narrative, see S. Dawson, C. Englert and T. Plehn, "Higgs Physics: It ain't over till it's over," Phys. Rept. **816** (2019) 1 [arXiv:1808.01324].
- [60] C. Quigg, "Dream Machines," Rev. Accel. Sci. Tech. 10 (2019) 3 [arXiv:1808.06036].
- [61] C. Quigg, "Perspectives and Questions," Yale University Nuclear/Particle/Astrophysics Seminar slides November 5, 2019 [zenodo.3663044].