



FERMILAB-SLIDES-20-023-ND

NEUTRINO INTERACTION MEASUREMENTS ON ARGON

Kirsty Duffy, Fermi National Accelerator Laboratory

on behalf of the MicroBooNE Collaboration

XXIX International Conference on Neutrino Physics

23rd June 2020

This document was prepared by MicroBooNE collaboration using the resources of the Fermi National Accelerator Laboratory (Fermilab), a U.S. Department of Energy, Office of Science, HEP User Facility. Fermilab is managed by Fermi Research Alliance, LLC (FRA), acting under Contract No. DE-AC02-07CH11359.

55 cm

Run 3469 Event 53223, Oct.

Cross-section measurements **on argon** are vital to reduce systematic uncertainties for the **SBN** program and **DUNE**

With **low thresholds** and **4π acceptance**, Liquid Argon Time Projection Chambers (LArTPCs) are powerful detectors to **study detailed final state topologies** and **quantitatively inform theoretical models**

Expanded statistics, better detector understanding since Neutrino 2018

Are models able to describe ν -Ar data?

ArgoNeuT
pathfinder, low statistics

MicroBooNE
high statistics, but initial measurements had large systematic uncertainties

Precision MicroBooNE
improved detector understanding → high statistics, low uncertainties

Future experiments: higher statistics (**SBND, ICARUS, DUNE-ND**) and lower systematics due to measurements by **LArIAT, ProtoDUNEs**

Many measurements of ν -Ar scattering

ν_μ CC inclusive cross section



Single-differential cross section

Phys. Rev. Lett. 108 161802 (2012)



Updated single-differential cross section

Phys. Rev. D 89, 112003 (2014)



Double-differential cross section

Phys. Rev. Lett. 123, 131801 (2019)



Single-differential cross section with updated detector and interaction models

MICROBOONE-NOTE-1069-PUB

ν_μ exclusive channels



Charged-particle multiplicity

Eur. Phys. J. C79, 248 (2019)



ν_μ CCQE-like scattering

Eur. Phys. J. C 79 673 (2019)

arXiv:2006.00108 (submitted to PRL)



ν_μ and $\bar{\nu}_\mu$ CC2p production

Phys. Rev. D 90, 012008 (2014)



ν_μ CC π^0 production

Phys. Rev. D99, 091102(R) (2019)



ν_μ and $\bar{\nu}_\mu$ NC π^0 production

Phys. Rev. D 96, 012006 (2017)



ν_μ and $\bar{\nu}_\mu$ CC π^+ production

Phys. Rev. D 98, 052002 (2018)



ν_μ and $\bar{\nu}_\mu$ Coherent CC π^+ production

Phys. Rev. Lett. 113, 261801 (2014)



ν_μ CC kaon production

MICROBOONE-NOTE-1071-PUB



ν_μ NC 1p production

MICROBOONE-NOTE-1067-PUB

Other measurements



ν_e and $\bar{\nu}_e$ scattering (inclusive)

arXiv:2004.01956[hep-ex]



MeV-scale physics

MICROBOONE-NOTE-1076-PUB



Limits on millicharged particles

Phys. Rev. Lett. 124, 131801 (2020)

Many measurements of ν -Ar scattering

ν_μ CC inclusive cross section



Single-differential cross section

Phys. Rev. Lett. 108 161802 (2012)



Updated single-differential cross section

Phys. Rev. D 89, 112003 (2014)



Double-differential cross section

Phys. Rev. Lett. 123, 131801 (2019)



Single-differential cross section with updated detector and interaction models

MICROBOONE-NOTE-1069-PUB

ν_μ exclusive channels



Charged-particle multiplicity

Eur. Phys. J. C79, 248 (2019)



ν_μ CCQE-like scattering

Eur. Phys. J. C 79 673 (2019)

arXiv:2006.00108 (submitted to PRL)



ν_μ and $\bar{\nu}_\mu$ CC2p production

Phys. Rev. D 90, 012008 (2014)



ν_μ CC π^0 production

Phys. Rev. D99, 091102(R) (2019)



ν_μ and $\bar{\nu}_\mu$ NC π^0 production

Phys. Rev. D 96, 012006 (2017)



ν_μ and $\bar{\nu}_\mu$ CC π^+ production

Phys. Rev. D 98, 052002 (2018)



ν_μ and $\bar{\nu}_\mu$ Coherent CC π^+ production

Phys. Rev. Lett. 113, 261801 (2014)



ν_μ CC kaon production

MICROBOONE-NOTE-1071-PUB



ν_μ NC 1p production

MICROBOONE-NOTE-1067-PUB

Other measurements



ν_e and $\bar{\nu}_e$ scattering (inclusive)

arXiv:2004.01956[hep-ex]

MeV-scale physics

Phys. Rev. D 99, 012002 (2019)



MeV-scale physics

MICROBOONE-NOTE-1076-PUB





Limits on millicharged particles

Phys. Rev. Lett. 124, 131801 (2020)

ArgoNeuT is a 40x47x90cm³ LArTPC

JINST 7 P10019 (2012)



- **2 planes** of wires with 4mm spacing collect charge from drifting electrons following secondary particle tracks
- **1.35x10²⁰ POT** data in NuMI beamline at Fermilab 2009-2010: $\langle E_{\nu e} \rangle = 4.3 \text{ GeV}$, $\langle E_{\bar{\nu} e} \rangle = 10.5 \text{ GeV}$
- Placed in front of MINOS near detector at Fermilab: use as **tracking spectrometer**





Figure from T.Yang, NuINT 2017

See also: "Searches for New Physics with MicroBooNE" by G. Karagiorgi, 2nd July



Thank you to Fermilab Accelerator Division, Cryogenics team, and Operations team!

MicroBooNE: 170 ton LArTPC

JINST 12 P02017 (2017)



- **3 planes** of wires (vertical, $+60^\circ$, -60°) with **3mm spacing**
- **32 PMTs** collect light from flash at time of interaction
- Sits in **two neutrino beams** at Fermilab: BNB (on-axis, $\langle E_{\nu\mu} \rangle = 800$ MeV) and NuMI (off-axis, $\langle E_{\nu e} \rangle = 650$ MeV)
- Stable detector operation since 2015: **longest-running LArTPC to date**
 - >95% DAQ uptime
 - 1.52×10^{21} POT collected in total (analyses shown here use subsets, not full POT)
 - From December 2017: data with **Cosmic Ray Tagger (CRT)**

GETTING THE MOST OUT OF LArTPCs

JINST 13 P07006 (2018)



JINST 13 P07007 (2018)

- MicroBooNE Collaboration has made **huge improvements** in our understanding of the detector since Neutrino 2018
- Detailed understanding of detector is **key to our R&D mission** for future LArTPCs
- **Improved signal processing** (2D deconvolution) accounts for interfering wire signals on all three planes
- Tracking is hard when particles go parallel to wires. Precise calorimetry on all planes → 3D tracking → **4π particle identification**




IMPROVED DETECTOR UNDERSTANDING ENABLES BETTER MEASUREMENTS

MICROBOONE-NOTE-1084-PUB



- Cosmic rejection power (without kinematic requirements) **increased by factor of 8** compared to previous publications
- **High efficiency:** 80.4% for ν_μ CC (87.6% for ν_e CC)
- **Increased statistics:** 11.3k events, compared to 4.3k events in same data set for 2019 CC inclusive measurement



Many measurements of ν -Ar scattering

ν_μ CC inclusive cross section



Single-differential cross section

Phys. Rev. Lett. 108 161802 (2012)



Updated single-differential cross section

Phys. Rev. D 89, 112003 (2014)



Double-differential cross section

Phys. Rev. Lett. 123, 131801 (2019)



Single-differential cross section with updated detector and interaction models

MICROBOONE-NOTE-1069-PUB

ν_μ exclusive channels



Charged-particle multiplicity

Eur. Phys. J. C79, 248 (2019)



ν_μ CCQE-like scattering

Eur. Phys. J. C 79 673 (2019)

arXiv:2006.00108 (submitted to PRL)



ν_μ and $\bar{\nu}_\mu$ CC2p production

Phys. Rev. D 90, 012008 (2014)



ν_μ CC π^0 production

Phys. Rev. D99, 091102(R) (2019)



ν_μ and $\bar{\nu}_\mu$ NC π^0 production

Phys. Rev. D 96, 012006 (2017)



ν_μ and $\bar{\nu}_\mu$ CC π^+ production

Phys. Rev. D 98, 052002 (2018)



ν_μ and $\bar{\nu}_\mu$ Coherent CC π^+ production

Phys. Rev. Lett. 113, 261801 (2014)



ν_μ CC kaon production

MICROBOONE-NOTE-1071-PUB



ν_μ NC 1p production

MICROBOONE-NOTE-1067-PUB

Other measurements



ν_e and $\bar{\nu}_e$ production (inclusive)

arXiv:2004.01956[hep-ex]



MeV-scale physics

MICROBOONE-NOTE-1076-PUB




Limits on millicharged particles



Phys. Rev. Lett. 124, 131801 (2020)

CC INCLUSIVE CROSS SECTION MEASUREMENT


PRL 123, 131801 (2019)



- Selection presented at Neutrino 2018 → new since then: **double-differential** cross section measurement
- **First time** double-differential cross section has been measured on argon: compared to worldwide interaction generators
- All models **overpredict in high-momentum, forward going bins**: interesting physics in this region!




IMPROVED DETECTOR UNDERSTANDING ENABLES BETTER MEASUREMENTS



Previous Measurement


PRL 123, 131801 (2019)

IMPROVED DETECTOR UNDERSTANDING ENABLES BETTER MEASUREMENTS



Previous Measurement



PRL 123, 131801 (2019)



Current Measurement

MICROBOONE-NOTE-1069-PUB

IMPROVED DETECTOR UNDERSTANDING ENABLES BETTER MEASUREMENTS



Previous Measurement

PRL 123,

Improved detector understanding,
reconstruction, CRT → higher purity

ν_μ CC (signal) purity: 50% → 71.9%

Entering backgrounds: 33.3% → 13.2%


IMPROVED INTERACTION MODELING ENABLES BETTER MEASUREMENTS

Better data-simulation

agreement from improved neutrino interaction modeling

- GENIE v2.12.2 → GENIE v3.0.6
- **Tuned** CCQE and CCMEC models to T2K ν_μ CC0 π data
- T2K data is on a carbon target → tuning seems to give **good agreement with MicroBooNE's argon-target data**

PRL 123, 131801 (201)



GENIE v3.0.6 models used:

QE/MEC → **J. Nieves, J.E. Amaro, M. Valverde** Phys. Rev. C 70, 055503 (2004) and

R. Gran, J. Nieves, F. Sanchez, M. Vicente-Vacas Phys. Rev. D 88, 113007 (2013)


RES/COH → **C. Berger, L. Sehgal** Phys. Rev. D 76, 113004 (2007), Phys. Rev. D 79, 053003 (2009)

FSI → work by **L. Salcedo, E. Oset, M. Vicente-Vacas, C. Garcia-Recio**

Nucl. Phys. A 484, 557-592 (1988) and **V. Pandharipande, S.C. Pieper** Phys. Rev. C 45, 791-798 (1992)

DRASTICALLY REDUCED SYSTEMATIC UNCERTAINTIES

MICROBOONE-NOTE-1075-PUB . MICROBOONE-NOTE-1069-PUB



Flux-integrated cross section
consistent with previous measurement


Drastically reduced systematic uncertainties

Source	Uncertainty	
	Previous Analysis	This Analysis
Detector response	16.2%	3.3%
Cross section	3.9%	2.7%
Flux	12.4%	10.5%
Dirt background	10.9%	3.3%
Cosmic ray background	4.2%	N/A
POT counting	2.0%	2.0%
CRT	N/A	1.7%
Total Sys. Error	23.8%	12.1%
Statistics	1.4%	3.8%
Total (Quadratic Sum)	23.8%	12.7%

PRL 123, 131801 (2019)

DRASTICALLY REDUCED SYSTEMATIC UNCERTAINTIES

MICROBOONE-NOTE-1075-PUB . MICROBOONE-NOTE-1069-PUB



Flux-integrated cross section
consistent with previous measurement

Drastically reduced systematic uncertainties


Largest reduction in uncertainties comes from improved detector understanding

Source	Uncertainty	
	Previous Analysis	This Analysis
Detector response	16.2%	3.3%
Cross section	3.9%	2.7%
Flux	12.4%	10.5%
Dirt background	10.9%	3.3%
Cosmic ray background	4.2%	N/A
POT counting	2.0%	2.0%
CRT	N/A	1.7%
Total Sys. Error	23.8%	12.1%
Statistics	1.4%	3.8%
Total (Quadratic Sum)	23.8%	12.7%

PRL 123, 131801 (2019)

DRASTICALLY REDUCED SYSTEMATIC UNCERTAINTIES

MICROBOONE-NOTE-1075-PUB . MICROBOONE-NOTE-1069-PUB



Flux-integrated cross section
consistent with previous measurement

Drastically reduced systematic uncertainties



Instead of cosmic ray simulation, now use overlay: simulated neutrino interactions overlaid on real cosmic data → no uncertainty in cosmic ray model

Source	Uncertainty	
	Previous Analysis	This Analysis
Detector response	16.2%	3.3%
Cross section	3.9%	2.7%
Flux	12.4%	10.5%
Dirt background	10.9%	3.3%
Cosmic ray background	4.2%	N/A
POT counting	2.0%	2.0%
CRT	N/A	1.7%
Total Sys. Error	23.8%	12.1%
Statistics	1.4%	3.8%
Total (Quadratic Sum)	23.8%	12.7%

PRL 123, 131801 (2019)

IMPROVED CROSS SECTION MEASUREMENT

MICROBOONE-NOTE-1074-PUB . MICROBOONE-NOTE-1075-PUB . MICROBOONE-NOTE-1069-PUB



Single-differential cross section as a function of reconstructed muon momentum and angle → **very good agreement with previous measurement**, but **reduced uncertainties**

Future development towards **double-differential** cross-section measurement

Many measurements of ν -Ar scattering

ν_μ CC inclusive cross section



Single-differential cross section

Phys. Rev. Lett. 108 161802 (2012)



Updated single-differential cross section

Phys. Rev. D 89, 112003 (2014)



Double-differential cross section

Phys. Rev. Lett. 123, 131801 (2019)



Single-differential cross section with updated detector and interaction models

MICROBOONE-NOTE-1069-PUB

ν_μ exclusive channels



Charged-particle multiplicity

Eur. Phys. J. C79, 248 (2019)



ν_μ CCQE-like scattering

Eur. Phys. J. C 79 673 (2019)

arXiv:2006.00108 (submitted to PRL)



ν_μ and $\bar{\nu}_\mu$ CC2p production

Phys. Rev. D 90, 012008 (2014)



ν_μ CC π^0 production

Phys. Rev. D99, 091102(R) (2019)



ν_μ and $\bar{\nu}_\mu$ NC π^0 production

Phys. Rev. D 96, 012006 (2017)



ν_μ and $\bar{\nu}_\mu$ CC π^+ production

Phys. Rev. D 98, 052002 (2018)



ν_μ and $\bar{\nu}_\mu$ Coherent CC π^+ production

Phys. Rev. Lett. 113, 261801 (2014)



ν_μ CC kaon production

MICROBOONE-NOTE-1071-PUB



ν_μ NC 1p production

MICROBOONE-NOTE-1067-PUB

Other measurements



ν_e and $\bar{\nu}_e$ scattering (inclusive)

arXiv:2004.01956[hep-ex]



MeV-scale physics

MICROBOONE-NOTE-1076-PUB






Limits on millicharged particles

Phys. Rev. Lett. 124, 131801 (2020)

LArTPC STRENGTH: LOW PROTON THRESHOLDS


MICROBOONE-NOTE-1056-PUB
JINST 15, P03022 (2020)

- **Low thresholds** → probe 2p2h scattering and nuclear processes
- MicroBooNE: **300 MeV/c**
- ArgoNeuT: **200 MeV/c**
Phys. Rev. D 90, 012008 (2014)
- Protons **identified by Bragg peak** in last 30 cm of track



CCQE-LIKE CROSS SECTION


Eur. Phys. J. C 79 673 (2019) arXiv:2006.00108 [hep-ex] (2020)



First extraction of ν_μ - ^{40}Ar CCQE-like cross section using a surface LArTPC





Signal: 1 muon ($p_\mu > 100 \text{ MeV}/c$), 1 proton ($p_p > 300 \text{ MeV}/c$) with extra cuts on coplanarity and transverse imbalance to enhance CCQE contribution
 $\rightarrow \sim 84\%$ CC1p0 π ($\sim 81\%$ CCQE) purity,
 $\sim 20\%$ efficiency

Good agreement with models, except at very **forward muon scattering** angles
 \rightarrow low momentum transfer (similar effects previously seen in carbon scattering)



CCQE-LIKE CROSS SECTION

Eur. Phys. J. C 79 673 (2019) arXiv:2006.00108 [hep-ex] (2020)




**Proton
momentum
threshold 300
MeV/c**


Across all kinematic variables, agreement is improved if forward muon angles are excluded

NCIP CROSS SECTION

MICROBOONE-NOTE-1067-PUB




- Measure cross section for neutral-current single proton production
- Measurement includes events with $Q^2 \sim 2m_p T_p = 0.1 \text{ GeV}^2$, **significantly lower** than previous measurements
- Future development towards a measurement of **NC elastic scattering** cross section → measure strange component of neutral-current axial form factor





CC π^\pm PRODUCTION

Phys. Rev. D 98, 052002 (2018)



ArgoNeuT ν_μ




- **Highly relevant for DUNE:** dominant interaction mode at DUNE energies and less well-understood than CCQE-like scattering
- ArgoNeuT ν_μ and $\bar{\nu}_\mu$ CC π^\pm measurement:
 - Select **two-track events**: one matched to a track in MINOS (muon candidate)
 - Select **CC π^\pm events** using dE/dx of pion candidate, event topology
- MicroBooNE measurement in progress: development work focused on **muon/pion separation** and **pion reinteractions**

LArIAT, ProtoDUNEs

LArTPC STRENGTH: ELECTRONS AND PHOTONS


S. Berkman, Poster
410, Poster Session 2

- **Electrons and photons produce showers in LArTPCs** → important to understand for ν_e appearance searches in SBN and DUNE
- π^0 interactions are a background (although often can be distinguished by energy deposition) — can also be used to **verify shower reconstruction** by reconstructing π^0 mass peak



**MicroBooNE CC π^0 Measurement:
presented at Neutrino 2018**



Phys. Rev. D 99, 091102(R) (2019)



FIRST MEASUREMENT OF ELECTRON NEUTRINO CROSS SECTION

arXiv:2004.01956 [hep-ex]

- Flux-averaged $\nu_e + \bar{\nu}_e$ cross section measured by ArgoNeuT
- Purity 78.9%, efficiency 10.5% → 13 events selected
- First measurement of its kind in an energy regime highly relevant for DUNE, demonstration of fully-automated reconstruction and analysis
- MicroBooNE ν_e measurements in BNB - see “Searches for New Physics with MicroBooNE”, 2nd July
- $\nu_e + \bar{\nu}_e$ cross-section measurements in progress with NuMI beam: purity 40%, efficiency 9% → ~100 events in 5×10^{19} POT



A. Fiorentini, poster
369, poster session 3

I. Lepetic, Poster 88,
Poster Session I

I. Lepetic, Poster 89,
Poster Session 2

A. Bhat, Poster 4,
Poster Session 4


PUSHING THE LIMITS

Phys. Rev. Lett. 124, 131801 (2020)


Phys. Rev. D 99, 012002 (2019)

MICROBOONE-NOTE-1076-PUB

- **CC kaon production:** rare process, few existing measurements, background for **proton decay** $p \rightarrow K^+ \nu$ searches in DUNE
- Selection developed with 68% purity and 7% efficiency → expect 12 candidate interactions in 1.3×10^{21} POT



- **Reconstruct sub-MeV particles:** photons from nucleus de-excitation or neutron re-interactions
- Demonstration of low-threshold LArTPC capabilities
- Used in ArgoNeuT to place constraints on BSM physics (millicharged particles)



Many measurements of ν -Ar scattering

ν_μ CC inclusive cross section



Single-differential cross section

Phys. Rev. Lett. 108 161802 (2012)



Updated single-differential cross section

Phys. Rev. D 89, 112003 (2014)



Double-differential cross section

Phys. Rev. Lett. 123, 131801 (2019)



Single-differential cross section with updated detector and interaction models

MICROBOONE-NOTE-1069-PUB

ν_μ exclusive channels



Charged-particle multiplicity

Eur. Phys. J. C79, 248 (2019)



ν_μ CCQE-like scattering

Eur. Phys. J. C 79 673 (2019)

arXiv:2006.00108 (submitted to PRL)



ν_μ and $\bar{\nu}_\mu$ CC2p production

Phys. Rev. D 90, 012008 (2014)



ν_μ CC π^0 production

Phys. Rev. D99, 091102(R) (2019)



ν_μ and $\bar{\nu}_\mu$ NC π^0 production

Phys. Rev. D 96, 012006 (2017)



ν_μ and $\bar{\nu}_\mu$ CC π^+ production

Phys. Rev. D 98, 052002 (2018)



ν_μ and $\bar{\nu}_\mu$ Coherent CC π^+ production

Phys. Rev. Lett. 113, 261801 (2014)



ν_μ CC kaon production

MICROBOONE-NOTE-1071-PUB



ν_μ NC 1p production

MICROBOONE-NOTE-1067-PUB

Other measurements



ν_e and $\bar{\nu}_e$ scattering (inclusive)

arXiv:2004.01956[hep-ex]



MeV-scale physics

MICROBOONE-NOTE-1076-PUB



Limits on millicharged particles

Phys. Rev. Lett. 124, 131801 (2020)

FUTURE PROSPECTS

- This talk has focused on current results from **MicroBooNE** and recent results from **ArgoNeuT**

FUTURE PROSPECTS

- This talk has focused on current results from **MicroBooNE** and recent results from **ArgoNeuT**

MicroBooNE recent improvements in detector understanding directly results in **reduced systematic uncertainties** on CC inclusive measurement
→ will form the basis of new, **more precise measurements** of neutrino interactions on argon in the near future

Additional measurements in progress: ν_μ CC π^0 , ν_μ CC1 π^+ , ν_μ CC-Coherent π^+ , ν_μ CC0 π Np, ν_μ CC0 π 2p, ν_μ CC0 π STV, ν_μ KDAR CC0 π , ν_μ CC0 π 0p, ν_e CC inclusive, ν_e CC0 π 1p

FUTURE PROSPECTS


- This talk has focused on current results from **MicroBooNE** and recent results from **ArgoNeuT**
- Exclusive measurements will be informed by test-beam measurements of charged particles in LArTPCs (e.g. interactions of pions, protons) by **LArIAT** and **ProtoDUNE**s
- In the future, look out for more measurements from upcoming experiments: **SBND**, **ICARUS**, and eventually **DUNE-ND**

ICARUS will start taking data very soon
SBND will collect **7m v-Ar interactions in 3 years**
See talk: “**ICARUS and the Fermilab Short-Baseline Neutrino Program**” on 2nd July



SUMMARY

- Cross-section measurements on argon are **vital** for the success of the SBN program and eventually DUNE
- **Huge progress** over the past two years since Neutrino 2018
→ measurements with low-energy protons, π^0 s, ν_e s and more are **extremely valuable**
- LArTPC technology has demonstrated 4π acceptance and ability to measure sub-MeV energies — we are already able to make **precise, accurate measurements of exclusive final states**
- First time we can confront **models tuned to carbon** with high-statistics argon data: seem to do well with the data now available
- More (and more precise) measurements expected in the future
→ **stronger tests** of our models



–Cross-section Posters and Supporting Documents

■ MicroBooNE

- ν_μ CCQE-like measurement: **A. Papadopoulou, Poster 145, Poster Session 2**
Cosmic rejection: Eur. Phys. J. C 79 673 (2019) , Cross-section measurement: arXiv:2006.00108 [hep-ex] (2020)
- Updated ν_μ CC-inclusive measurement: MICROBOONE-NOTE-1069-PUB
- ν_μ NC1p measurement: **L. Ren, Poster 292, Poster Session 4** MICROBOONE-NOTE-1067-PUB
- ν_μ CCKaon selection: **A. Fiorentini, Poster 369, Poster Session 3** MICROBOONE-NOTE-1071-PUB
- MeV-scale Physics: **A. Bhat, Poster 4, Poster Session 4** MICROBOONE-NOTE-1076-PUB
- Interaction model and uncertainties: MICROBOONE-NOTE-1074-PUB
- Detector uncertainties: **L. Yates, Poster 176, Poster Session 1** MICROBOONE-NOTE-1075-PUB

■ ArgoNeuT

- $\nu_e + \bar{\nu}_e$ CC inclusive measurement: **R. Fitzpatrick, Poster 139, Poster Session 4**
arXiv:2004.01956 [hep-ex]
- Improved limits on millicharged particles: **I. Lepetic, Poster 89, Poster Session 2**
Phys. Rev. Lett. 124, 131801 (2020)

MICROBOONE PUBLICATIONS

* cross-section specific

- MicroBooNE collaboration, "First Measurement of Differential Charged Current Quasi-Elastic-Like Muon Neutrino Argon Scattering Cross Sections with the MicroBooNE Detector", [arXiv:2006.00108](#), submitted to PRL
- MicroBooNE collaboration, "Vertex-Finding and Reconstruction of Contained Two-track Neutrino Events in the MicroBooNE Detector", [arXiv: 2002.09375](#), submitted to JINST
- MicroBooNE collaboration, "Search for heavy neutral leptons decaying into muon-pion pairs in the MicroBooNE detector", [arXiv:1911.10545](#), *Phys. Rev. D101*, 052001 (2020), [Fermilab News article](#) (02/13/20)
- MicroBooNE collaboration, "Reconstruction and Measurement of O(100) MeV Electromagnetic Activity from $\pi^0 \rightarrow \gamma \gamma$ Decays in the MicroBooNE LAr TPC", [arXiv:1910.02166](#), *JINST 15, P02007* (2020)
- MicroBooNE collaboration, "A Method to Determine the Electric Field of Liquid Argon Time Projection Chambers Using a UV Laser System and its Application in MicroBooNE", [arXiv:1910.01430](#), submitted to JINST
- MicroBooNE collaboration, "Calibration of the Charge and Energy Response of the MicroBooNE Liquid Argon Time Projection Chamber Using Muons and Protons". [arXiv:1907.11736](#). *JINST 15, P03022* (2020)
- MicroBooNE collaboration, "First Measurement of Inclusive Muon Neutrino Charged Current Differential Cross Sections on Argon at Enu ~0.8 GeV with the MicroBooNE Detector", [arXiv:1905.09694](#), *Phys. Rev. Lett.* 123, 131801 (2019), [Fermilab News article](#) (12/13/19)
- MicroBooNE collaboration, "Design and Construction of the MicroBooNE Cosmic Ray Tagger System", [arXiv:1901.02862](#), *JINST 14, P04004* (2019)
- MicroBooNE collaboration, "Rejecting Cosmic Background for Exclusive Neutrino Interaction Studies with Liquid Argon TPCs: A Case Study with the MicroBooNE Detector", [arXiv:1812.05679](#), accepted by *Eur. J. Phys. C*
- MicroBooNE collaboration, "First Measurement of Muon Neutrino Charged Current Neutral Pion Production on Argon with the MicroBooNE LAr TPC", [arXiv:1811.02700](#), *Phys. Rev. D99, 091102(R)* (2019)
- MicroBooNE collaboration, "A Deep Neural Network for Pixel-Level Electromagnetic Particle Identification in the MicroBooNE Liquid Argon Time Projection Chamber", [arXiv:1808.07269](#), *Phys. Rev. D99, 092001* (2019), [Fermilab News article](#) (09/12/18), [DOE HEP Science Highlight](#) (01/30/19)

- MicroBooNE collaboration, "Comparison of Muon-Neutrino-Argon Multiplicity Distributions Observed by MicroBooNE to GENIE Model Predictions", [arXiv:1805.06887](#), *Eur. Phys. J. C79*, 248 (2019), [Fermilab News article](#) (05/31/18)
- MicroBooNE collaboration, "Ionization Electron Signal Processing in Single Phase LAr TPCs II: Data/Simulation Comparison and Performance in MicroBooNE", [arXiv:1804.02583](#), *JINST 13, P07007* (2018), [Fermilab News article](#) (07/09/18), [DOE HEP Science Highlight](#) (05/21/19)
- MicroBooNE collaboration, "Ionization Electron Signal Processing in Single Phase LAr TPCs I: Algorithm Description and Quantitative Evaluation with MicroBooNE Simulation", [arXiv:1802.08709](#), *JINST 13, P07006* (2018), [Fermilab News article](#) (07/09/18), [DOE HEP Science Highlight](#) (05/21/19)
- MicroBooNE collaboration, "The Pandora Multi-Algorithm Approach to Automated Pattern Recognition of Cosmic Ray Muon and Neutrino Events in the MicroBooNE Detector", [arXiv:1708.03135](#), *Eur. Phys. J. C78*, 1, 82 (2018)
- MicroBooNE collaboration, "Measurement of Cosmic Ray Reconstruction Efficiencies in the MicroBooNE LAr TPC Using a Small External Cosmic Ray Counter", [arXiv:1707.09903](#), *JINST 12, P12030* (2017)
- MicroBooNE collaboration, "Noise Characterization and Filtering in the MicroBooNE Liquid Argon TPC", [arXiv:1705.07341](#), *JINST 12, P08003* (2017), [Fermilab News article](#) (07/05/17), [DOE HEP Science Highlight](#) (05/16/18)
- MicroBooNE collaboration, "Michel Electron Reconstruction Using Cosmic Ray Data from the MicroBooNE LAr TPC", [arXiv:1704.02927](#), *JINST 12, P09014* (2017)
- MicroBooNE collaboration, "Determination of Muon Momentum in the MicroBooNE LAr TPC Using an Improved Model of Multiple Coulomb Scattering", [arXiv:1703.06187](#), *JINST 12 P10010* (2017)
- MicroBooNE collaboration, "Convolutional Neural Networks Applied to Neutrino Events in a Liquid Argon Time Projection Chamber", [arXiv:1611.05531](#), *JINST 12, P03011* (2017)
- MicroBooNE collaboration, "Design and Construction of the MicroBooNE Detector", [arXiv:1612.05824](#), *JINST 12, P02017* (2017)


ARGONEUT PUBLICATIONS

* cross-section specific

- ArgoNeuT collaboration, “First Measurement of Electron Neutrino Scattering Cross Section on Argon”, [arXiv:2004.01956\[hep-ex\]](#)
- ArgoNeuT collaboration, “Improved Limits on Millicharged Particles Using the ArgoNeuT Experiment at Fermilab”, [arXiv:1911.07996\[hep-ex\]](#), *Phys. Rev. Lett.* **124**, 131801 (2020)
- ArgoNeuT collaboration, “Demonstration of MeV-Scale Physics in Liquid Argon Time Projection Chambers Using ArgoNeuT”, [arXiv:1810.06502\[hep-ex\]](#), *Phys. Rev. D* **99**, 012002 (2019)
- ArgoNeuT collaboration, “First measurement of the cross section for ν_μ and $\bar{\nu}_\mu$ induced single charged pion production on argon using ArgoNeuT”, [arXiv:1804.10294\[hep-ex\]](#), *Phys. Rev. D* **98**, 052002 (2018)
- ArgoNeuT collaboration, “First Observation of Low Energy Electron Neutrinos in a Liquid Argon Time Projection Chamber”, [arXiv:1610.04102\[hep-ex\]](#), *Phys. Rev. D* **95**, 072005 (2017)
- ArgoNeuT collaboration, “Measurement of ν_μ and $\bar{\nu}_\mu$ neutral current $\pi^0 \rightarrow \gamma\gamma$ production in the ArgoNeuT detector”, [arXiv:1511.00941\[hep-ex\]](#), *Phys. Rev. D* **96**, 012006 (2017)
- ArgoNeuT collaboration, “First Measurement of Neutrino and Antineutrino Coherent Charged Pion Production on Argon”, [arXiv:1408.0598\[hep-ex\]](#), *Phys. Rev. Lett.* **113**, 261801 (2014), *Phys. Rev. Lett.* **114**, 039901 (erratum) (2015)
- ArgoNeuT collaboration, “Detection of Back-to-Back Proton Pairs in Charged-Current Neutrino Interactions with the ArgoNeuT Detector in the NuMI Low Energy Beam Line”, [arXiv:1405.4261\[nucl-ex\]](#), *Phys. Rev. D* **90**, 012008 (2014)
- ArgoNeuT collaboration, “Measurements of Inclusive Muon Neutrino and Antineutrino Charged Current Differential Cross Sections on Argon in the NuMI Antineutrino Beam”, [arXiv:1404.4809\[hep-ex\]](#), *Phys. Rev. D* **89**, 112003 (2014)
- ArgoNeuT collaboration, “A Study of Electron Recombination Using Highly Ionizing Particles in the ArgoNeuT Liquid Argon TPC”, [arXiv:1306.1712\[physics.ins-det\]](#), *JINST* **8** P08005 (2013)
- ArgoNeuT collaboration, “Analysis of a Large Sample of Neutrino-Induced Muons with the ArgoNeuT Detector”, [arXiv:1205.6702\[physics.ins-det\]](#), *JINST* **7** P10020 (2012)
- ArgoNeuT collaboration, “First Measurements of Inclusive Muon Neutrino Charged Current Differential Cross Sections on Argon”, [arXiv:1111.0103\[hep-ex\]](#), *Phys. Rev. Lett.* **108** 161802 (2012)
- ArgoNeuT collaboration, “The ArgoNeuT Detector in the NuMI Low-Energy beam line at Fermilab”, [arxiv:1205.6747\[physics.ins-det\]](#), *JINST* **7** P10019 (2012)

MICROBOONE DATA COLLECTION



Very **stable detector operation**, smooth and **steady data taking**,
efficient data acquisition




COSMIC RAY TAGGER (CRT)

JINST 14 P04004 (2019)

MICROBOONE-NOTE-1069-PUB




ELECTRON-PHOTON DISCRIMINATION



**MicroBooNE ν_e Selection:
presented at Nulnt 2018**

MICROBOONE-NOTE-1054-PUB



Current: with improved detector understanding

MICROBOONE-NOTE-1085-PUB

CC INCLUSIVE CROSS SECTION MEASUREMENT

Phys. Rev. Lett. 123, 131801 (2019)


Selection presented at Neutrino 2018

- Topological and optical information → reject background events from cosmic rays
- Energy deposition profile: select candidate muon

Largest ever sample of neutrino interactions on argon


Signal (CC-inclusive) events: 50.4%

Largest background: **cosmic rays** (29%)
→ directly measured with beam-off data












New since Neutrino 2018:
double-differential cross section measurement

CC Inclusive double-differential measurement












Phys. Rev. Lett. 123, 131801 (2019)

CC Inclusive double-differential measurement




Phys. Rev. Lett. 123, 131801 (2019)

CC Inclusive double-differential measurement





Phys. Rev. Lett. 123, 131801 (2019)

CC Inclusive double-differential measurement




PREVIOUS CC-INCLUSIVE CROSS SECTION: 1D MEASUREMENT

Single-differential cross section measurement presented at
Neutrino 2018: [PRL 123, 131801 \(2019\)](#)



UPDATED CC-INCLUSIVE SELECTION: SYSTEMATIC UNCERTAINTIES

MICROBOONE-NOTE-1075-PUB . MICROBOONE-NOTE-1069-PUB



Flux-integrated cross section
**consistent with previous
measurement**

Drastically reduced systematic
uncertainties

CRT better able to reject interactions
in surrounding material (“dirt”) →
reduced systematic uncertainty

Source	Uncertainty	
	Previous Analysis	This Analysis
Detector response	16.2%	3.3%
Cross section	3.9%	2.7%
Flux	12.4%	10.5%
Dirt background	10.9%	3.3%
Cosmic ray background	4.2%	N/A
POT counting	2.0%	2.0%
CRT	N/A	1.7%
Total Sys. Error	23.8%	12.1%
Statistics	1.4%	3.8%
Total (Quadratic Sum)	23.8%	12.7%

PRL 123, 131801 (2019)

UPDATED CC-INCLUSIVE SELECTION


MICROBOONE-NOTE-1069-PUB

■ **Cosmic rejection:**

- Topological and optical information
- Veto events with CRT hits when all tracks are contained
- Cut on CRT hit-reconstructed vertex z position if tracks are uncontained


■ **Muon selection**

- Longest track > 20 cm is muon candidate
- Topology must be track-like
- Energy deposition must be inconsistent with a proton




UPDATED CC-INCLUSIVE SELECTION EFFICIENCY


MICROBOONE-NOTE-1069-PUB




(a) True neutrino energy



(b) True muon momentum



(c) True muon $\cos(\theta)$



(d) True muon ϕ



- Good efficiency to select QuasiElastic, Meson Exchange Current, RESonant pion production, and Deep Inelastic Scattering interaction channels

- → truly inclusive selection

- Efficiency limited at low neutrino energy/muon momentum due to muon candidate track length > 20 cm requirement

UPDATED CC-INCLUSIVE CROSS SECTION: SYSTEMATIC UNCERTAINTIES

MICROBOONE-NOTE-1069-PUB





Improved detector understanding → **drastically reduced** systematic uncertainties from detector modeling

UPDATED CC-INCLUSIVE CROSS SECTION: REDUCED PHASE SPACE

MICROBOONE-NOTE-1069-PUB



- Efficiency is low for $p_\mu < 150$ MeV due to >20 cm track length requirement
- Main result includes all CC interactions in signal definition (no requirement on muon momentum)
- As a check: re-extract cross section with signal requirement $p_\mu > 150$ MeV/c (note: statistical uncertainties only)



UPDATED CC-INCLUSIVE CROSS SECTION: REDUCED PHASE SPACE

MICROBOONE-NOTE-1069-PUB

- Efficiency is low for $p_\mu < 150$ MeV due to >20 cm track length requirement
- Main result includes all CC interactions in signal definition (no requirement on muon momentum)
- As a check: re-extract cross section with signal requirement $p_\mu > 150$ MeV/c (note: statistical uncertainties only)



CCPIO SELECTIONS

Phys. Rev. D 99, 091102(R) (2019)



Low-energy photons appear more track like
→ low reconstruction efficiency
→ requiring that we reconstruct both π^0 photons limits statistics

Two-shower selection

→ validate π^0 hypothesis by invariant diphoton mass

Single shower selection


→ validate photon hypothesis
→ maximize statistics for cross section measurement



CC π^0 PRODUCTION

Phys. Rev. D 99, 091102(R) (2019)

Select π^0 events by looking for **one** or **two** showers in addition to a candidate muon track




CCQE-LIKE CROSS SECTION

Eur. Phys. J. C 79 673 (2019) arXiv:2006.00108 [hep-ex] (2020)

- **First extraction of ν_μ - ${}^{40}\text{Ar}$ CCQE-like cross section using a surface LArTPC**
- Important channel for low-energy excess search (and other LArTPC oscillation analyses)
- Signal: 1 muon (>100 MeV/c), 1 proton (300 MeV/c)

Selection:

- Two tracks
- Energy deposition consistent with one muon and one proton
- Tracks are not collinear
- Tracks are coplanar
- Low vertex activity
- Low transverse momentum




CCQE CROSS SECTION: MODEL COMPARISONS

A. Papadopoulou, poster 145,
poster session 2





Eur. Phys. J. C 79 673 (2019) arXiv:2006.00108 [hep-ex] (2020)

- **Nominal:** GENIE v2.12.2. Bodek-Ritchie Fermi Gas, Llewellyn-Smith CCQE model, empirical MEC model, Rein-Sehgal resonant and coherent scattering model, “hA” FSI model
- **hA2015:** GENIE v2.12.2 with a more recent “hA2015” FSI model
- **Alternative:** GENIE v2.12.10. Local Fermi Gas, Nieves CCQE model, Nieves MEC model, KLN-BS resonant and BS coherent scattering models, and hA2015 FSI model
- **v3.0.6:** GENIE v3.0.6. Same model configuration as Alternative model, with hA2018 FSI model



NCIP SELECTION

- Single **isolated track**
- Must be contained within fiducial volume
- Length 1.2 - 200 cm
- Must be **forward-going** ($\cos\theta > 0$ w.r.t neutrino beam direction)
- **Deposited energy profile** consistent with a proton
- Multi-class gradient-boosted decision tree used to **further reduce background from cosmic interactions**



NCIP CROSS SECTION MEASUREMENT


L. Ren, poster 292,
poster session 4

MICROBOONE-NOTE-1067-PUB

- Measure cross section for neutral-current single proton production
- Signal: 1 isolated proton

Selection:
42.1% efficiency, 29.8% purity

- Largest backgrounds :
- Proton from charged-current interaction (other particles missed by reconstruction)
 - Proton from non-1p neutral-current interaction (other particles missed by reconstruction)





CC KAON PRODUCTION SELECTION

A. Fiorentini, poster
369, poster session 3

MICROBOONE-NOTE-1071-PUB



- **CC kaon production:** rare process, few existing measurements, background for **proton decay** $p \rightarrow K^+ \nu$ searches in DUNE
- Selection developed on simulation: look for K^+ track from neutrino interaction and μ^+ from K^+ decay
- 67.7% purity and 7% efficiency → expect to select 12 candidate interactions in 1.3×10^{21} POT MicroBooNE data set
- Aim: cross section measurement and study of K^+ in LArTPC



CC KAON SELECTION

MICROBOONE-NOTE-1071-PUB

- Reject cosmic rays based on topology and optical information
- Must have one track with energy deposition consistent with a muon
- K^+ candidate selected based on energy deposition: consistent with a kaon and inconsistent with a proton
- Must have exactly one μ^+ candidate: must start within 5cm of end of kaon track, track length >30cm, energy deposition inconsistent with a proton




ARGONEUT ELECTRON NEUTRINO SELECTION

R. Fitzpatrick,
Poster 139,
Poster Session 4


arXiv:2004.01956 [hep-ex]

- Focus on reconstructing leading shower in neutrino interaction
- Reject events with a muon reconstructed in downstream MINOS detector
- Reject events with through-going muons
- Reconstructed shower must be forward-going: $\cos(\theta) > 0.05$ w.r.t. beam direction
- Shower must start within 2cm of reconstructed vertex
- Electron candidate selected based on topology and charge of entire candidate shower using a BDT: BDT score > 0.9



ARGONEUT CHARGED PION PRODUCTION MEASUREMENT


- ArgoNEUT: CC $\bar{\nu}$ π^\pm production [Phys. Rev. D 98, 052002 \(2018\)](#)
- Select two-track events: one matched to a track in MINOS (muon candidate)
- Select CC $\bar{\nu}$ π^\pm events using dE/dx of pion candidate, event topology
- Overall purity 35.8% (ν), 55.7% ($\bar{\nu}$)
- 337 selected ν events (285 $\bar{\nu}$)



ARGONEUT CHARGED PION PRODUCTION MEASUREMENT

ν_μ CC π^\pm ArgoNeuT measurement


Phys. Rev. D 98, 052002 (2018)



ARGONEUT CHARGED PION PRODUCTION MEASUREMENT

$\bar{\nu}_\mu$ CCI π^\pm ArgoNeuT measurement

Phys. Rev. D 98, 052002 (2018)



Resonant pion production model


- GENIE, NEUT: Rein-Sehgal
- NuWro: $\Delta(1232)$ resonance only

Nonresonant model

- NEUT: Rein-Sehgal
- GENIE, NuWro: Bodek-Yang above resonance region, extrapolate smoothly to converge with resonance model at lower W

FSI


- NEUT, NuWro: Salcedo-Oset cascade
- GENIE: effective cascade model
- GiBUU: quantum-kinetic transport theory



HARGED PION MFASUREMENT

rement

Phys. Rev. D 98, 052002 (2018)



Paper conclusions

GiBUU: good agreement

NuWro, NEUT: similar, higher than measured cross section



GENIE: higher than other generators and measured cross sections (with reanalysis of bubble chamber data in [EPJC \(2016\)](#)
[76: 474](#) points to GENIE's nonresonant background prediction)

All predictions within 2σ of measurement, except GENIE \bar{v} (3.3σ)



MICROBOONE ν_e CC INCLUSIVE SELECTION

MICROBOONE-NOTE-1054-PUB




ν_e selection efficiency: 9%, purity: 40%

MICROBOONE $\nu_e + \bar{\nu}_e$ MEASUREMENTS

MICROBOONE-NOTE-1054-PUB



- Select $\nu_e + \bar{\nu}_e$ CC inclusive interactions by looking for single shower
- Purity 40%, efficiency 9% $\rightarrow \sim 100$ events in 5×10^{19} POT
- Future plans:
 - $\nu_e + \bar{\nu}_e$ CC inclusive flux integrated cross section measurement
 - $\nu_e + \bar{\nu}_e$ CC inclusive differential cross-section measurement
 - Exclusive $\nu_e + \bar{\nu}_e$ CC differential cross-section measurement



CHARGED PARTICLE MULTIPLICITY

Eur. Phys. J. C79, 248 (2019)

Precise resolution of a LArTPC allows us to look in detail at particles produced in a neutrino interaction → measure **number** and **kinematic distributions** of charged tracks produced



MEV-SCALE PHYSICS

Phys. Rev. Lett. 124, 131801 (2020) Phys. Rev. D 99, 012002 (2019) MICROBOONE-NOTE-1076-PUB

- Both ArgoNeuT and MicroBooNE have demonstrated ability to reconstruct **energy depositions from sub-MeV particles** (ArgoNeuT: 300 keV, MicroBooNE: 100 keV)
- Demonstration of **low-threshold LArTPC capabilities**: important for measurements of cross sections, especially solar neutrinos, supernova neutrinos, and neutrinos from μ DAR
- BSM physics **search for millicharged particles** in ArgoNeuT set leading limits (poster by I. Lepetic)

