

Novel constraints on fifth forces and ultralight dark sector with asteroidal data

Yu-Dai Tsai^{a,b,c}, Youjia Wu^d, Sunny Vagnozzi^{e,f,g}
and Luca Visinelli^{h,i}

^aDepartment of Physics and Astronomy, University of California, Irvine, CA 92697-4575, USA

^bFermi National Accelerator Laboratory (Fermilab), Batavia, IL 60510, USA

^cKavli Institute for Cosmological Physics (KICP), University of Chicago, Chicago, IL 60637, USA

^dLeinweber Center for Theoretical Physics, Department of Physics, University of Michigan, Ann Arbor, MI 48109, USA

^eDepartment of Physics, University of Trento, Via Sommarive 14, 38123 Povo (TN), Italy

^fTrento Institute for Fundamental Physics and Applications (TIFPA)-INFN, Via Sommarive 14, 38123 Povo (TN), Italy

^gKavli Institute for Cosmology (KICC), University of Cambridge, Madingley Road, Cambridge CB3 0HA, United Kingdom

^hTsung-Dao Lee Institute (TDLI), 520 Shengrong Road, 201210 Shanghai, P. R. China

ⁱSchool of Physics and Astronomy, Shanghai Jiao Tong University, 800 Dongchuan Road, 200240 Shanghai, P. R. China

E-mail: yt444@cornell.edu, youjiawu@umich.edu, sunny.vagnozzi@unitn.it, luca.visinelli@sjtu.edu.cn

Abstract. We study for the first time the possibility of probing long-range fifth forces utilizing asteroid astrometric data, via the fifth force-induced orbital precession. We examine nine Near-Earth Object (NEO) asteroids whose orbital trajectories are accurately determined via optical and radar astrometry. Focusing on a Yukawa-type potential mediated by a new gauge field (dark photon) or a baryon-coupled scalar, we estimate the sensitivity reach for the fifth force coupling strength and mediator mass in the mass range $m \simeq (10^{-21} - 10^{-15})$ eV, near the “fuzzy” dark matter region. Our estimated sensitivity is comparable to leading limits from equivalence principle tests, potentially exceeding these in a specific mass range. Furthermore, we set new constraints on Yukawa gravity. The fifth force-induced precession increases with the orbital semi-major axis in the limit of a small mass m , motivating the study of objects further away from the Sun. We discuss future prospects for extending our study to more than a million asteroids, including NEOs, main-belt asteroids, Hildas, and Jupiter Trojans, as well as trans-Neptunian objects and exoplanets.

Contents

1	Introduction	1
2	Light particles and orbital precessions	2
3	Methods	3
4	Results and Discussion	5
5	Conclusions and Future Prospects	7
6	Appendix	9

1 Introduction

The study of precessions has revealed some of the deepest secrets of Nature. Most notably, the correct prediction for Mercury’s precession rate from General Relativity (GR) is one of the theory’s major successes [1–3]. The findings of the *Muon $g - 2$* experiment measuring the muon anomalous precession frequency might hint at the existence of physics beyond the Standard Model (SM) [4–6]. New connections between microscopic physics and macroscopic planetary science can be established by studying the precessions of celestial objects, due to long-range forces mediated by (new) ultralight particles.

There are strong motivations to investigate the existence of new light, weakly-coupled degrees of freedom beyond the SM (BSM), which are generic features of string theory [7–10], and are candidates for the dark matter (DM) and dark energy (DE) [11–15]. For example, ultralight (fuzzy) DM may play a significant role in shaping galactic structure [16–18], and DE could be in the form of a quintessential axion [19–21]. Efforts towards detecting the signatures of new light particles and their associated fifth forces range from laboratory and space tests [22–44] to cosmological [45–60] and astrophysical studies [61–79].

In addition, one can also consider modifications to GR (see e.g. [80]). In particular, a Yukawa-type potential, referred to in this context as Yukawa gravity [81–83], has been adopted in order to parametrize the phenomenological effects of theories with massive gravitons [84].

The motion of asteroids is continuously and carefully monitored for various reasons that include planetary defense purposes [85], to the extent that dedicated studies have been recently financed both by the National Aeronautics and Space Administration (NASA) [86] and the European Space Agency (ESA) [87] asteroid missions. These studies benefit from current and future radar and optical data, including from facilities and missions such as Arecibo (decommissioned), Goldstone, Catalina, the Vera Rubin Observatory (VRO), and Gaia [88–91]. Such studies are not free of challenges, as asteroid trajectories are subject to perturbations that range from gravitational effects from other celestial objects to non-gravitational effects due to the thermal and reflective properties of the asteroid’s surface. Recent advances in studying physical parameters and relevant physical processes (including GR parameters, solar quadrupole moment, and Yarkovsky effect) from asteroid data, taking into account these perturbations [92, 93], inspire us to examine the possibility of probing new physics with asteroid astrometry.

In this paper, we provide a proof-of-principle study using asteroid precessions and astrometric data to probe new ultralight particles. Previously, planets, exoplanets, and Kuiper Belt Objects (KBOs) were used to test GR and/or search for dark sector particles [94–114], yet the potential of probing new physics using asteroids remains mostly unharvested. Thanks to advances in radar and optical astrometry, the motion of asteroids, especially those classified as Near-Earth Objects (NEOs), is tracked much more precisely than KBOs and exoplanets. The use of asteroids over planets [115] also carries several advantages, ranging from their sheer number, to their spread in orbital radius allowing to probe a wide range of parameter space. Focusing on light mediators in the mass range $m \simeq (10^{-21} - 10^{-15})$ eV, or equivalently in the Compton wavelength range $\lambda \simeq (10^{-3} - 10^3)$ au, we estimate the sensitivity reach of asteroid precessions to the mediator mass and coupling, which we find to be competitive with some of the most stringent equivalence principle tests [22, 27, 29, 116], and outline further steps to improve the analysis. We find that the strongest bounds are realized in the region where $\lambda \sim a$, where a is the asteroid’s semi-major axis; this motivates the future use of objects orbiting further away from the Sun to probe lighter mediators efficiently. While we focus on gauged SM symmetries and baryon-coupled ultralight scalars as concrete examples, our study is broadly applicable to various well-motivated new physics models.

We expect our study to integrate into ongoing efforts to study gravity in space. We delineate the possibility of conducting similar studies using extended asteroid catalogs, Trans-Neptunian Objects (TNOs), and exoplanets. The growing wealth of available optical and radar data would lead to significant improvements in our results.

The rest of this paper is then organized as follows. In section 2 we lay down the theoretical motivations for our study. In section 3 we discuss the methods used to assess the observables against asteroid data, with the results presented in section 4. Conclusions and future prospects are drawn in section 5.

2 Light particles and orbital precessions

We consider a celestial body of mass M_* orbiting the Sun and subject to an additional fifth-force mediated by a new light particle, with an associated potential $V(r)$ which is modeled as:

$$V(r) = -\frac{GM_\odot M_*}{(1 + \alpha_1)r} \left(1 + \alpha_2 \exp\left(-\frac{r}{\lambda}\right) \right), \quad (2.1)$$

where M_\odot is the solar mass and λ the Yukawa force range. Here, α_1 and α_2 are Yukawa parameters whose values will be specified later in the text according to the different types of Yukawa models. The potential in eq. (2.1) leads to deviations from the body’s Newtonian orbit, introducing an orbital precession alongside GR effects accounted for by $g_{\mu\nu}$.

We consider planar motion and fix $\theta = \pi/2$. Adopting the inverse radius variable $u \equiv 1/r = u(\varphi)$ and defining $\tilde{G} \equiv G/(1 + \alpha_1)$, we obtain the orbit’s fundamental equation (in SI units) [115]:

$$\frac{d^2 u}{d\varphi^2} + u - \frac{\tilde{G}M_\odot}{L^2} = \frac{3\tilde{G}M_\odot}{c^2} u^2 + \alpha_2 \frac{\tilde{G}M_\odot}{L^2} \left(1 + \frac{1}{\lambda u} \right) e^{-\frac{1}{\lambda u}}, \quad (2.2)$$

where L is the orbital angular momentum per unit mass. The first term on the right-hand side of eq. (2.2) leads to well-known GR corrections, while the second term leads to fifth force-induced corrections. Solving eq. (2.2) numerically determines the fifth force-induced

precession, which, upon comparison to observations, can constrain α_1 and α_2 given the force range.

For most of the Yukawa potentials generated from new BSM light scalars or vectors, $\alpha_1 = 0$ and we define $\alpha_2 \equiv \tilde{\alpha}$. Examples to which our study can be applied include gauged $U(1)_B$ [117, 118], $U(1)_{B-L}$ [119–121], $L_\mu - L_{e,\tau}$ [122–124], and baryon-coupled scalar [125–128] models. In these models:

$$\tilde{\alpha} = \pm \frac{g^2}{4\pi G} \frac{Q_\odot Q_*}{M_\odot M_*}, \quad (2.3)$$

where $+$ corresponds to attractive and $-$ corresponds to repulsive force under this notation, and g is the coupling strength. For a BSM boson, the mediator mass $m \equiv \hbar c/\lambda$. For illustrative purposes, we shall focus on the mediator being either a gauged $U(1)_B$ dark photon or an ultralight scalar coupled to the baryon number. The coupling is given by $g = g_{\phi,A'}$ for either a scalar (ϕ) or vector (A') mediator, whose mass is $m = m_{\phi,A'}$. Moreover, $Q_* \equiv M_*/m_p$ and $Q_\odot \equiv M_\odot/m_p$ are the celestial object and Sun total baryon numbers respectively, with m_p the proton mass. The gauged $U(1)_B$ exhibits a chiral anomaly whose cancellation can be achieved, e.g., by introducing additional appropriately constrained particles [129, 130] or invoking extra model building [131–133]. In this phenomenological study, we assume no self-interaction for the scalar. We focus on the asteroid phenomenology of these models and we emphasize again that our method is not limited to the $U(1)_B$ dark photon and scalar mediators case studies. For the scalar coupling, we take the Lagrangian to be $\mathcal{L}_\phi \subset (g_{\phi,p}\bar{p}p + g_{\phi,n}\bar{n}n + g_{\phi,e}\bar{e}e)\phi$, and consider two cases: Case (a) would have the same magnitude as the $U(1)_B$ coupling, modulo a different sign (attractive for scalar and repulsive for vectors), so $g_{\phi,p} = g_{\phi,n}$ and $g_{\phi,e} = 0$.

For the case of Yukawa gravity, $\alpha_1 = \alpha_2$ and we define $\alpha_2 \equiv \tilde{\alpha}_g$ to avoid any confusion. We also define the graviton mass $m_g \equiv \hbar c/\lambda$.

3 Methods

We specialize to asteroids as the celestial objects of interest. Our goal is to estimate the sensitivity reach for the coupling strength and mediator mass of a Yukawa-type fifth force, using the induced orbital precession. To this end, we focus on nine asteroids with precise radar and optical trajectory determinations, studied in detail in Ref. [92]. These are NEOs with semi-major axes $a \in [0.64, 1.08]$ au and eccentricities $e \in [0.48 - 0.90]$. We are only interested in the impact of the fifth force field on the induced orbital precession. A fully-fledged analysis entails *a*) computing the fifth force impact on the asteroid trajectory via an appropriate integrator, accounting for perturbations from all nearby objects, and *b*) using raw astrometric measurements of the asteroid’s trajectory to constrain the fifth force. As this is the beginning attempt to perform this type of analysis, our aim is simply to estimate the fifth force sensitivity reach, while providing a proof-of-principle for the feasibility of such a study and laying the foundations for future detailed analyses.

Various effects contribute to asteroid orbital precession. Perturbations from planetary motions source the largest contributions to the orbital precession of a body, see e.g. Refs. [134, 135]. Two additional contributors are GR effects and solar oblateness [92]. These two effects contribute to the perihelion precession as measured from a fixed reference direction per orbital period, for an orbit whose inclination angle with respect to the solar equator is i_{eq} , as [136]:

$$\Delta\varphi_0 = \frac{6\pi GM_\odot}{a(1-e^2)c^2} \left[\frac{2-\beta+2\gamma}{3} \right] + 3\pi R_\odot^2 \frac{2-3\sin^2 i_{\text{eq}}}{2a^2(1-e^2)^2} J_2, \quad (3.1)$$

where R_\odot is the solar radius and J_2 the solar quadrupole moment [137]. The parameters γ and β describe the deviations from GR in the Parameterized Post-Newtonian (PPN) approach, with GR being recovered when $\beta = \gamma = 1$ [137–139], and with deviations from unity being therefrom tightly constrained by Solar System probes [140–142]. We verified that the precession cross-contribution from both J_2 and $\tilde{\alpha}$ is sub-dominant. The effects on the apsidal precession coming from the presence of planetary perturbations has been assessed e.g. in Ref. [143].

To estimate the fifth force sensitivity reach, we impose that the new physics contribution to the orbital precession in eq. (3.2) does not exceed the uncertainty budget associated to the two major precession contributors β and J_2 (as $|1-\gamma|$ is more tightly constrained than $|1-\beta|$). We lean upon the results of Ref. [92], who estimated the sensitivity to β and J_2 obtainable from a fully-fledged analysis of the 9 asteroids: the analysis was based on the Mission Operations and Navigation Toolkit Environment (MONTE) software [144], which numerically integrates the orbit equations of motion, using a dynamical model including gravitational perturbations from nearby celestial objects and accounting for Yarkovsky drift.¹

We obtain the precession $\Delta\varphi$ by numerically solving eq. (2.2) with initial conditions $u(0) = [a(1-e)]^{-1}$ and $u'(0) = 0$, corresponding to an elliptic orbit with eccentricity e and at its perihelion for $\varphi = 0$. The induced precession is estimated by expressing $u = [a(1-e^2)]^{-1}[1+e\cos\varphi(1-\delta)]$, solving for δ , and deriving the precession as $\Delta\varphi = 2\pi\delta/(1-\delta)$. The new physics contribution is then $\Delta\varphi_{\text{Yukawa}}(\alpha_1, \alpha_2, \lambda) = \Delta\varphi - \Delta\varphi_{\text{GR}}$.

The very light mediator limit $m_\phi \ll \hbar/ac$ (or, equivalently, $\lambda \gg a$) admits an analytical expression for the fifth force-induced precession, obtained by expanding around the exponential term:

$$|\Delta\varphi_{\text{Yukawa}}| \simeq \frac{2\pi\tilde{\alpha}_2}{1+\tilde{\alpha}_2} \left(\frac{am_\phi c}{\hbar}\right)^2 (1-e) = \frac{2\pi\tilde{\alpha}_2}{1+\tilde{\alpha}_2} \left(\frac{a}{\lambda}\right)^2 (1-e). \quad (3.2)$$

We stress that we *do not* use this approximation to estimate our sensitivity reach, but numerically solve eq. (2.2), later verifying that eq. (3.2) holds when $\lambda \gg a$. Note that the precession goes to zero in the limit $\lambda \rightarrow \infty$ where the Newtonian $1/r$ potential is recovered. In this limit $|\Delta\varphi_{\text{Yukawa}}| \propto a^2$, which carries a different functional dependence on a compared to the GR and J_2 contributions, so studying objects within a wide range of a and e can help differentiate the contributions from these terms.

Note that $\left|\frac{\partial\Delta\varphi_0}{\partial\beta}\right|\sigma_\beta \sim \left|\frac{\partial\Delta\varphi_0}{\partial J_2}\right|\sigma_{J_2}$, meaning that both parameters are determined to comparable levels as far as precessions are concerned, and we want to estimate the range of uncertainty for the Yukawa parameters at a given λ . We therefore require that the new physics precession contribution does not exceed the uncertainty budget associated to β and J_2 :

$$\Delta\varphi_{\text{Yukawa}}^2 < \left|\frac{\partial\Delta\varphi_0}{\partial\beta}\right|^2\sigma_\beta^2 + \left|\frac{\partial\Delta\varphi_0}{\partial J_2}\right|^2\sigma_{J_2}^2 + 2\rho\left|\frac{\partial\Delta\varphi_0}{\partial\beta}\right|\left|\frac{\partial\Delta\varphi_0}{\partial J_2}\right|\sigma_{J_2}\sigma_\beta, \quad (3.3)$$

where ρ is the correlation coefficient. The above inequality is a function of the fifth force parameters, and values thereof which saturate the inequality give our estimated sensitivity reach. We repeat these steps for each of the 9 asteroids, obtaining 9 separate (but comparable) limits on the α 's for different values of λ . We implicitly assume that the central values of the measured orbital precessions are consistent with the expectations given GR and all

¹Newton's constant G is measured independently in cold-atom experiments and other techniques to a relative uncertainty of $\sim 2 \times 10^{-5}$ [145–147].

nearby perturbers modelled in Ref. [92], and therefore that there is no detection of fifth force, whose contribution accordingly must not exceed the precession uncertainty budget. In other words, our analysis is akin to a forecast around a fiducial model with no fifth force. For this reason, we also do not account for the perturbations due to the planetary motion in the Solar System, since the presence of a fifth force could already be altering the results from the tinier contributions from GR effects and solar oblateness.

A covariance analysis of the 9 asteroids based on a 2022 sensitivity projection infers $\sigma_\beta = 5.6 \times 10^{-4}$ and $\sigma_{J_2} = 2.7 \times 10^{-8}$, and a correlation coefficient $\rho = -0.72$, with a Monte Carlo forecast recovering similar figures [92]. We base our sensitivity reach estimate on the 2022 values to reflect the current sensitivity. We also present an estimate based on the “optimistic” 2022 values $\sigma_\beta = 2 \times 10^{-4}$ and $\sigma_{J_2} = 10^{-8}$ given in Ref. [92].

4 Results and Discussion

In figure 1 we show the estimated sensitivity to the $U(1)_B$ dark photon and baryon-coupled ultralight scalar couplings, as a function of their masses. In these examples, all the baryons in the Sun and the asteroids are charged and the specific compositions thereof do not affect our results. Three specific asteroids, i.e. TU3, MN, and BD19, deliver the strongest limits (see figure 2 in the Appendix for the sensitivity reaches for each of the asteroids), given by the solid curve in the figure (whereas the dashed curve shows a stronger sensitivity reach based on the optimistic projection described earlier). We have chosen to report the *tightest* sensitivity reach since all 9 curves are comparable. Unsurprisingly, the peak sensitivity is achieved for mediator masses approximately corresponding to the (inverse) orbital radius of each asteroid. On the same figure, we also mark the regions corresponding to typical orbital radii of other (non-NEO) asteroids and TNOs. Finally, we note that the inferred sensitivity to the fifth force coupling strength and mediator mass within the $U(1)_B$ model can easily be converted to other long-range forces, including those associated to gauged $U(1)_{B-L}$ and $L_e - L_{\mu,\tau}$ symmetries, following Refs. [148, 149].

For a fixed coupling strength and mediator mass, the repulsive (attractive) force from a vector dark photon (scalar) mediator would yield a precession of essentially equal magnitude, as we have checked numerically, resulting in eq. (3.3) delivering identical sensitivity reaches. A fully-fledged analysis of the raw asteroid astrometric data should account for the different sign in the precession contribution, and might therefore return different constraints for the two cases.

Also shown in figure 1 are existing leading bounds from equivalence principle tests [22, 27, 29, 116], black hole (BH) superradiance [150], and planetary precession [115]. Lunar Laser Ranging (LLR) [24, 151] provides the leading bound for masses $\gtrsim 10^{-16}$ eV and is included for completeness. A concept for a hypothetical space mission similar to LLR exploiting the Martian moon Phobos is also under development, and is referred to as Phobos Laser Ranging (PLR) [152]. Such a test would be extremely sensitive to the parameter $\tilde{\alpha}$ and could potentially improve bounds by two orders of magnitude on scales of an astronomical unit. Only the vector superradiance bound is present since the scalar superradiance one requires further studies on supermassive BHs, owing to large uncertainties concerning their environments. Asteroids offer an actual probe of fifth forces with range beyond the au scale that is complementary to the scales probed within laboratories; more complex long-range force models may be invoked to bypass torsion balance constraints [15, 153]. We have also set a new limit on Yukawa gravity.




Figure 1: Estimated sensitivity reach for the mass and coupling strength of **(a)** $U(1)_B$ dark photons and baryon-coupled scalars with $g_{\phi,p} = g_{\phi,n}$ and $g_{\phi,e} = 0$ (**left panel**); **(b)** Yukawa gravity with modification parameter $\tilde{\alpha}_g$ as a function of the graviton mass m_g (**right panel**). The asteroid curves are obtained by studying the precessions of nine NEO asteroids with $a \in [0.64, 1.08]$ au and $e \in [0.48 - 0.90]$. The solid black curve shows the tightest 1σ sensitivity reaches from asteroids TU3, MN, and BD19, while the dashed black curve is the 1σ sensitivity reach based on the optimistic 2022 projection of Ref. [92]. Existing constraints include those from planets [115], EP tests [116], and vector superradiance [67], the latter is only applicable to the dark photon A' . LLR [24] provides the leading bound for masses $\gtrsim 10^{-16}$ eV and is included for completeness. As a note of caution, while laboratory bounds are shown along with the results from our analysis, the length scales probed by the two different techniques differ by orders of magnitude. They provide complementary probes of the fifth forces with different screening effects.

Furthermore, we show in figure 1 that precession tests, including asteroid ones, are potentially competitive with EP laboratory tests. This demonstrates that precession tests from asteroids and other planetary objects are especially suitable in probing EP-conserving (or approximately conserving) long-range forces.² Also note that, while laboratory bounds are shown along with the results from our analysis, the length scales probed by the two different techniques differ by orders of magnitude. They provide complementary probes of the fifth forces with different screening effects. We have checked that the results obtained are consistent with the requirement that the precession contribution from the fifth-force is below the limits from GR. This can be seen with an explicit computation at small scales $\lambda \lesssim a$ by comparing the results in Eq. (3.2) with the precession expected in GR, leading to:

$$g_{\phi, A'} \lesssim 6 \times 10^{-23} \frac{1}{(1-e)(1+e)^{1/2}} \left(\frac{a}{\text{au}}\right)^{-3/2} \left(\frac{m_{\phi, A'}}{10^{-18} \text{ eV}}\right)^{-1}, \quad (4.1)$$

a much looser requirement than the bounds derived in Fig. 1 for all asteroids considered.

²Satellite-based EP tests have recently reported no evidence for EP violation [154], or are planned for deployment such as the Satellite Test of the Equivalence Principle (STEP) [155].

Summing up, we note that prospects for advancing our understanding of these nine asteroids, including their hazardous or complex nature, are bright. For example, the binary-asteroid system (66391) 1999 KW4/Moshup is a potential threat to Earth due to its orbital trajectory, and is the subject of intense studies [156].

5 Conclusions and Future Prospects

Our work attempts to connect fundamental new physics and astrometry data for planetary objects. Focusing on nine near-Earth asteroids, our analysis provides a general recipe and sensitivity reach estimate for long-range fifth forces induced by ultralight mediators. Follow-up opportunities are detailed below.

New target objects — There are opportunities to extend our study to $\mathcal{O}(10^6)$ minor planets, classified in table 1. Including asteroids and comets, there are ~ 25000 NEOs (comets are less ideal for our study since they are subject to strong non-gravitational perturbations), a significant number of which have orbits that can be tracked to a similar level of precision as the nine asteroids considered. Among these nine asteroids, the one whose trajectory is determined to lowest accuracy is 2004 KH17, whose semi-major axis is nonetheless measured to $\simeq 1$ km precision. Currently ~ 1800 NEOs have orbits known to comparable or higher accuracy: 247 of these has been analyzed to study Yarkovsky drift [93]. Neglecting systematics, ~ 1800 NEOs can potentially improve our sensitivity reach by more than 1 order of magnitude.

Beyond NEOs, other asteroids including main-belt asteroids (M), Hildas (H), and Jupiter Trojans (JT) can serve similar purposes. Their larger semi-major axes imply that their sensitivity reaches would peak at lower mediator masses, allowing us to probe lighter dark sector particles. Moreover, as the Yarkovsky drift weakens with increasing distance from the Sun as $a^{-1/2}$ [157], the effects over Hildas and Trojans would be negligible compared to NEOs for kilometer-sized bodies and for the same time of the observations. Achieving precision comparable to NEOs might be challenging, but spacecraft ranging may provide data with precision rivalling/surpassing radar observations: for example, the LUCY space mission [158] will provide precision data for Trojans.

TNOs and ETNOs, residing in the outer Solar System, are of extreme interest owing to their trajectories being subject to significantly less gravitational perturbations and solar thermal effects. Their large semi-major axes mean they can be used to probe ultra-light mediators at even lower masses. All these objects are labelled in figure 1 according to their typical semi-major axes.

New observations — Radar studies including Goldstone and the recently decommissioned Arecibo [88] have been collecting high-precision NEOs astrometrical data. VRO will discover a factor of 5 more Solar System minor objects (see table 1 of Ref. [90]), while other optical sky surveys such as Catalina [89], Pan-STARRS [159], ATLAS [160], DECam [161], and ZTF [162] will also be of great use to such studies. High precision astrometry is also achievable with space-based telescopes such as Hubble [163], James Webb [164], Euclid [165], and Roman [166]. LUCY [158] will visit Trojans and the JANUS spacecraft will investigate two binary asteroids [167, 168], providing valuable information to extend our study. New astrometrical techniques such as occultation can substantially improve orbital trajectory determinations [169]. Asteroids also affect gravitational wave detections through gravity gradient noise [170, 171].

Data storage and dedicated software development — Our work motivates the study and inclusion of precession measurements for objects stored in the JPL small objects [172] and

Minor Planets	a [au]	\sim Numbers
Near-Earth Objects (NEOs)	$< 1.3^*$	> 25000
Main-Belt Asteroids (M)	$\sim 2 - 3$	~ 1 million
Hildas (H)	3.7 - 4.2	> 4000
Jupiter Trojans (JT)	5.2	> 9800
Trans-Neptunian Objects (TNOs)	> 30	2700
Extreme TNOs (ETNOs)	> 150	12

Table 1: Targets for our future studies, for which opportunities are provided by sheer numbers and observational programs, classified roughly based on their typical semi-major axes.

*NEOs are defined as having perihelia $a(1 - e) < 1.3$ au.

Minor Planet Center [173] databases. On the analysis side, a fully-fledged study entails re-analyzing the (raw) astrometric asteroid trajectory data. Dedicated computing platforms such as MONTE [144], self-consistently modelling all relevant physical effects, can be used to this end after appropriate modification to include the fifth force effect. We expect this to be an important task for future studies [174].

Theory — Our study can be viewed as an investigation of a specific example of deviations from GR and/or the SM. Of course, the method can be extended to test other theories of gravity (e.g. [175–179]), or other types of dark sector models [125, 180–187], by computing their effects on the dynamics of celestial objects [174]. One can also consider asteroid tracking arrays (ATAs), analogously to pulsar timing arrays, to study gravitational waves and other aspects of fundamental physics.

Final outlook — We expect to broaden up attempts at probing fundamental physics from astrometric data for minor planets in the inner and outer Solar System. More generally, alongside seminal works [94–100, 102–114], we have only just begun exploring the full potential of establishing connections between microscopic new physics and macroscopic planetary observations, from near (NEOs) to far (exoplanets) celestial objects.

Acknowledgments

We thank Alex Drlica-Wagner, Davide Farnocchia, Adam Greenberg, Nick Gnedin, Marco Micheli, Olivier Minazzoli, Matthew Payne, Darryl Seligman, Leo Stein, and Quanzhi Ye for useful discussions regarding asteroid studies, and astronomy and planetary observations in general. We also thank Masha Baryakhtar, Nikita Blinov, Cedric Delaunay, Robert Lasenby, Tanmay Kumar Poddar, Tracy Slatyer, Yotam Soreq, Liantao Wang, Yue Zhang, and Yue Zhao for discussions regarding ultralight dark sector studies. We are grateful to Yuval Grossman, Marco Micheli, and Darryl Seligman for their invaluable comments on our draft. This document was prepared by Y-D.T. using the resources of the Fermi National Accelerator Laboratory (Fermilab), a U.S. Department of Energy, Office of Science, HEP User Facility. Fermilab is managed by the Fermi Research Alliance, LLC (FRA), acting under Contract No. DE-AC02-07CH11359. Y-D.T. also thanks KICP, University of Chicago, for hospitality. Y.W. acknowledges support from an MICDE Catalyst grant at University of Michigan,

DoE grant DE- SC007859, and the LCTP at the University of Michigan.S.V. was partially supported by the Isaac Newton Trust and the Kavli Foundation through a Newton-Kavli Fellowship, and by a grant from the Foundation Blanceflor Boncompagni Ludovisi, née Bildt.

6 Appendix

The method we proposed leads to an estimate for the sensitivity reach for each of the nine individual asteroids. This is shown in more detail in figure 2 in the mass-coupling plane, together with the leading sensitivity reach based on the optimistic projection in Ref. [92] (dashed black curve). Results are comparable across all asteroids.




Figure 2: Estimated sensitivity reach from each of the 9 asteroids (solid colored curves), and leading sensitivity reach based on the optimistic projection in Ref. [92] (dashed black curve). Asteroids are new probes of long-range forces in the \sim au range. The sensitivity can be improved by investigating an additional \sim 25000 NEOs. Long-range forces at larger distances can be studied using main-belt asteroids (M), Jupiter Trojans (JT), Hildas (H), TNOs, and ETNOs, as discussed in the main text.

References

- [1] U.J. Le Verrier, *Theorie du mouvement de Mercure*, *Annales de l'Observatoire de Paris* **5** (1859) 1.
- [2] A. Einstein, *The Foundation of the General Theory of Relativity*, *Annalen Phys.* **49** (1916) 769.

- [3] C. Corda, *The secret of planets' perihelion between Newton and Einstein*, *Phys. Dark Univ.* **32** (2021) 100834.
- [4] MUON G-2 collaboration, *Final Report of the Muon E821 Anomalous Magnetic Moment Measurement at BNL*, *Phys. Rev. D* **73** (2006) 072003 [[hep-ex/0602035](#)].
- [5] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, *A new evaluation of the hadronic vacuum polarisation contributions to the muon anomalous magnetic moment and to $\alpha(m_Z^2)$* , *Eur. Phys. J. C* **80** (2020) 241 [[1908.00921](#)].
- [6] MUON G-2 collaboration, *Measurement of the Positive Muon Anomalous Magnetic Moment to 0.46 ppm*, *Phys. Rev. Lett.* **126** (2021) 141801 [[2104.03281](#)].
- [7] P. Svrcek and E. Witten, *Axions In String Theory*, *JHEP* **06** (2006) 051 [[hep-th/0605206](#)].
- [8] A. Arvanitaki, S. Dimopoulos, S. Dubovsky, N. Kaloper and J. March-Russell, *String Axiverse*, *Phys. Rev. D* **81** (2010) 123530 [[0905.4720](#)].
- [9] M. Cicoli, M. Goodsell and A. Ringwald, *The type IIB string axiverse and its low-energy phenomenology*, *JHEP* **10** (2012) 146 [[1206.0819](#)].
- [10] L. Visinelli and S. Vagnozzi, *Cosmological window onto the string axiverse and the supersymmetry breaking scale*, *Phys. Rev. D* **99** (2019) 063517 [[1809.06382](#)].
- [11] R.D. Peccei, J. Sola and C. Wetterich, *Adjusting the Cosmological Constant Dynamically: Cosmons and a New Force Weaker Than Gravity*, *Phys. Lett. B* **195** (1987) 183.
- [12] C. Wetterich, *Cosmology and the Fate of Dilatation Symmetry*, *Nucl. Phys. B* **302** (1988) 668 [[1711.03844](#)].
- [13] B. Ratra and P.J.E. Peebles, *Cosmological Consequences of a Rolling Homogeneous Scalar Field*, *Phys. Rev. D* **37** (1988) 3406.
- [14] C. Wetterich, *Probing quintessence with time variation of couplings*, *JCAP* **10** (2003) 002 [[hep-ph/0203266](#)].
- [15] J. Khoury and A. Weltman, *Chameleon cosmology*, *Phys. Rev. D* **69** (2004) 044026 [[astro-ph/0309411](#)].
- [16] W. Hu, R. Barkana and A. Gruzinov, *Cold and fuzzy dark matter*, *Phys. Rev. Lett.* **85** (2000) 1158 [[astro-ph/0003365](#)].
- [17] L. Hui, J.P. Ostriker, S. Tremaine and E. Witten, *Ultralight scalars as cosmological dark matter*, *Phys. Rev. D* **95** (2017) 043541 [[1610.08297](#)].
- [18] P. Mocz et al., *First star-forming structures in fuzzy cosmic filaments*, *Phys. Rev. Lett.* **123** (2019) 141301 [[1910.01653](#)].
- [19] J.E. Kim and H.P. Nilles, *A Quintessential axion*, *Phys. Lett. B* **553** (2003) 1 [[hep-ph/0210402](#)].
- [20] M. Ibe, M. Yamazaki and T.T. Yanagida, *Quintessence Axion Revisited in Light of Swampland Conjectures*, *Class. Quant. Grav.* **36** (2019) 235020 [[1811.04664](#)].
- [21] G. Choi, W. Lin, L. Visinelli and T.T. Yanagida, *Cosmic birefringence and electroweak axion dark energy*, *Phys. Rev. D* **104** (2021) L101302 [[2106.12602](#)].
- [22] Y. Su, B.R. Heckel, E.G. Adelberger, J.H. Gundlach, M. Harris, G.L. Smith et al., *New tests of the universality of free fall*, *Phys. Rev. D* **50** (1994) 3614.
- [23] C.D. Hoyle, D.J. Kapner, B.R. Heckel, E.G. Adelberger, J.H. Gundlach, U. Schmidt et al., *Sub-millimeter tests of the gravitational inverse-square law*, *Phys. Rev. D* **70** (2004) 042004 [[hep-ph/0405262](#)].
- [24] J.G. Williams, S.G. Turyshev and D.H. Boggs, *Progress in lunar laser ranging tests of relativistic gravity*, *Phys. Rev. Lett.* **93** (2004) 261101 [[gr-qc/0411113](#)].

- [25] D.F. Mota and D.J. Shaw, *Evading Equivalence Principle Violations, Cosmological and other Experimental Constraints in Scalar Field Theories with a Strong Coupling to Matter*, *Phys. Rev. D* **75** (2007) 063501 [[hep-ph/0608078](#)].
- [26] P. Brax, C. van de Bruck, A.-C. Davis, D.F. Mota and D.J. Shaw, *Detecting chameleons through Casimir force measurements*, *Phys. Rev. D* **76** (2007) 124034 [[0709.2075](#)].
- [27] S. Schlamminger, K.Y. Choi, T.A. Wagner, J.H. Gundlach and E.G. Adelberger, *Test of the equivalence principle using a rotating torsion balance*, *Phys. Rev. Lett.* **100** (2008) 041101 [[0712.0607](#)].
- [28] P. Brax and G. Pignol, *Strongly Coupled Chameleons and the Neutronic Quantum Bouncer*, *Phys. Rev. Lett.* **107** (2011) 111301 [[1105.3420](#)].
- [29] T.A. Wagner, S. Schlamminger, J.H. Gundlach and E.G. Adelberger, *Torsion-balance tests of the weak equivalence principle*, *Class. Quant. Grav.* **29** (2012) 184002 [[1207.2442](#)].
- [30] C. Burrage, E.J. Copeland and E.A. Hinds, *Probing Dark Energy with Atom Interferometry*, *JCAP* **03** (2015) 042 [[1408.1409](#)].
- [31] R. Foot and S. Vagnozzi, *Diurnal modulation signal from dissipative hidden sector dark matter*, *Phys. Lett. B* **748** (2015) 61 [[1412.0762](#)].
- [32] MADMAX WORKING GROUP collaboration, *Dielectric Haloscopes: A New Way to Detect Axion Dark Matter*, *Phys. Rev. Lett.* **118** (2017) 091801 [[1611.05865](#)].
- [33] L. Perivolaropoulos, *Submillimeter spatial oscillations of Newton's constant: Theoretical models and laboratory tests*, *Phys. Rev. D* **95** (2017) 084050 [[1611.07293](#)].
- [34] C. Burrage and J. Sakstein, *Tests of Chameleon Gravity*, *Living Rev. Rel.* **21** (2018) 1 [[1709.09071](#)].
- [35] P. Touboul et al., *MICROSCOPE Mission: First Results of a Space Test of the Equivalence Principle*, *Phys. Rev. Lett.* **119** (2017) 231101 [[1712.01176](#)].
- [36] L. Perivolaropoulos and L. Kazantzidis, *Hints of modified gravity in cosmos and in the lab?*, *Int. J. Mod. Phys. D* **28** (2019) 1942001 [[1904.09462](#)].
- [37] C. Blanco, M. Escudero, D. Hooper and S.J. Witte, *Z' mediated WIMPs: dead, dying, or soon to be detected?*, *JCAP* **11** (2019) 024 [[1907.05893](#)].
- [38] ADMX collaboration, *Extended Search for the Invisible Axion with the Axion Dark Matter Experiment*, *Phys. Rev. Lett.* **124** (2020) 101303 [[1910.08638](#)].
- [39] L. Di Luzio, M. Giannotti, E. Nardi and L. Visinelli, *The landscape of QCD axion models*, *Phys. Rept.* **870** (2020) 1 [[2003.01100](#)].
- [40] I.M. Bloch, A. Caputo, R. Essig, D. Redigolo, M. Sholapurkar and T. Volansky, *Exploring new physics with O(keV) electron recoils in direct detection experiments*, *JHEP* **01** (2021) 178 [[2006.14521](#)].
- [41] S. Vagnozzi, L. Visinelli, P. Brax, A.-C. Davis and J. Sakstein, *Direct detection of dark energy: The XENON1T excess and future prospects*, *Phys. Rev. D* **104** (2021) 063023 [[2103.15834](#)].
- [42] Y.-D. Tsai, J. Eby and M.S. Safronova, *Direct detection of ultralight dark matter bound to the Sun with space quantum sensors*, *Nature Astron.* **7** (2023) 113 [[2112.07674](#)].
- [43] C.B. Adams et al., *Axion Dark Matter*, in *Snowmass 2021*, 3, 2022 [[2203.14923](#)].
- [44] D. Antypas et al., *New Horizons: Scalar and Vector Ultralight Dark Matter*, [2203.14915](#).
- [45] R. Hlozek, D. Grin, D.J.E. Marsh and P.G. Ferreira, *A search for ultralight axions using precision cosmological data*, *Phys. Rev. D* **91** (2015) 103512 [[1410.2896](#)].
- [46] D. Baumann, D. Green, J. Meyers and B. Wallisch, *Phases of New Physics in the CMB*, *JCAP* **01** (2016) 007 [[1508.06342](#)].

- [47] F. D’Eramo, R.Z. Ferreira, A. Notari and J.L. Bernal, *Hot Axions and the H_0 tension*, *JCAP* **11** (2018) 014 [[1808.07430](#)].
- [48] SIMONS OBSERVATORY collaboration, *The Simons Observatory: Science goals and forecasts*, *JCAP* **02** (2019) 056 [[1808.07445](#)].
- [49] V. Poulin, T.L. Smith, D. Grin, T. Karwal and M. Kamionkowski, *Cosmological implications of ultralight axionlike fields*, *Phys. Rev. D* **98** (2018) 083525 [[1806.10608](#)].
- [50] V. Poulin, T.L. Smith, T. Karwal and M. Kamionkowski, *Early Dark Energy Can Resolve The Hubble Tension*, *Phys. Rev. Lett.* **122** (2019) 221301 [[1811.04083](#)].
- [51] S. Vagnozzi, *New physics in light of the H_0 tension: An alternative view*, *Phys. Rev. D* **102** (2020) 023518 [[1907.07569](#)].
- [52] S. Vagnozzi, L. Visinelli, O. Mena and D.F. Mota, *Do we have any hope of detecting scattering between dark energy and baryons through cosmology?*, *Mon. Not. Roy. Astron. Soc.* **493** (2020) 1139 [[1911.12374](#)].
- [53] D. Green et al., *Messengers from the Early Universe: Cosmic Neutrinos and Other Light Relics*, *Bull. Am. Astron. Soc.* **51** (2019) 159 [[1903.04763](#)].
- [54] M. Escudero Abenza, *Precision early universe thermodynamics made simple: N_{eff} and neutrino decoupling in the Standard Model and beyond*, *JCAP* **05** (2020) 048 [[2001.04466](#)].
- [55] K.K. Rogers and H.V. Peiris, *Strong Bound on Canonical Ultralight Axion Dark Matter from the Lyman-Alpha Forest*, *Phys. Rev. Lett.* **126** (2021) 071302 [[2007.12705](#)].
- [56] W. Giarè, E. Di Valentino, A. Melchiorri and O. Mena, *New cosmological bounds on hot relics: Axions & Neutrinos*, *Mon. Not. Roy. Astron. Soc.* **505** (2021) 2703 [[2011.14704](#)].
- [57] I. Esteban and J. Salvado, *Long Range Interactions in Cosmology: Implications for Neutrinos*, *JCAP* **05** (2021) 036 [[2101.05804](#)].
- [58] E. Di Valentino, O. Mena, S. Pan, L. Visinelli, W. Yang, A. Melchiorri et al., *In the realm of the Hubble tension—a review of solutions*, *Class. Quant. Grav.* **38** (2021) 153001 [[2103.01183](#)].
- [59] S. Vagnozzi, *Consistency tests of Λ CDM from the early integrated Sachs-Wolfe effect: Implications for early-time new physics and the Hubble tension*, *Phys. Rev. D* **104** (2021) 063524 [[2105.10425](#)].
- [60] L. Perivolaropoulos and F. Skara, *Challenges for Λ CDM: An update*, *New Astron. Rev.* **95** (2022) 101659 [[2105.05208](#)].
- [61] B. Jain, V. Vikram and J. Sakstein, *Astrophysical Tests of Modified Gravity: Constraints from Distance Indicators in the Nearby Universe*, *Astrophys. J.* **779** (2013) 39 [[1204.6044](#)].
- [62] A. Arvanitaki, M. Baryakhtar and X. Huang, *Discovering the QCD Axion with Black Holes and Gravitational Waves*, *Phys. Rev. D* **91** (2015) 084011 [[1411.2263](#)].
- [63] M. Giannotti, I. Irastorza, J. Redondo and A. Ringwald, *Cool WISPs for stellar cooling excesses*, *JCAP* **05** (2016) 057 [[1512.08108](#)].
- [64] R. Brito, V. Cardoso and P. Pani, *Superradiance: New Frontiers in Black Hole Physics*, *Lect. Notes Phys.* **906** (2015) pp.1 [[1501.06570](#)].
- [65] R. Foot and S. Vagnozzi, *Solving the small-scale structure puzzles with dissipative dark matter*, *JCAP* **07** (2016) 013 [[1602.02467](#)].
- [66] A. Caputo, J. Zavala and D. Blas, *Binary pulsars as probes of a Galactic dark matter disk*, *Phys. Dark Univ.* **19** (2018) 1 [[1709.03991](#)].
- [67] M. Baryakhtar, R. Lasenby and M. Teo, *Black Hole Superradiance Signatures of Ultralight Vectors*, *Phys. Rev. D* **96** (2017) 035019 [[1704.05081](#)].

- [68] M.J. Stott and D.J.E. Marsh, *Black hole spin constraints on the mass spectrum and number of axionlike fields*, *Phys. Rev. D* **98** (2018) 083006 [1805.02016].
- [69] R. Roy and U.A. Yajnik, *Evolution of black hole shadow in the presence of ultralight bosons*, *Phys. Lett. B* **803** (2020) 135284 [1906.03190].
- [70] H. Davoudiasl and P.B. Denton, *Ultralight Boson Dark Matter and Event Horizon Telescope Observations of M87**, *Phys. Rev. Lett.* **123** (2019) 021102 [1904.09242].
- [71] D. Croon, S.D. McDermott and J. Sakstein, *New physics and the black hole mass gap*, *Phys. Rev. D* **102** (2020) 115024 [2007.07889].
- [72] M.J. Stott, *Ultralight Bosonic Field Mass Bounds from Astrophysical Black Hole Spin*, 2009.07206.
- [73] H. Desmond, J. Sakstein and B. Jain, *Five percent measurement of the gravitational constant in the Large Magellanic Cloud*, *Phys. Rev. D* **103** (2021) 024028 [2012.05028].
- [74] R. Roy, S. Vagnozzi and L. Visinelli, *Superradiance evolution of black hole shadows revisited*, *Phys. Rev. D* **105** (2022) 083002 [2112.06932].
- [75] F. Ferlito, S. Vagnozzi, D.F. Mota and M. Baldi, *Cosmological direct detection of dark energy: Non-linear structure formation signatures of dark energy scattering with visible matter*, *Mon. Not. Roy. Astron. Soc.* **512** (2022) 1885 [2201.04528].
- [76] Y. Chen, R. Roy, S. Vagnozzi and L. Visinelli, *Superradiant evolution of the shadow and photon ring of Sgr A**, *Phys. Rev. D* **106** (2022) 043021 [2205.06238].
- [77] S. Vagnozzi et al., *Horizon-scale tests of gravity theories and fundamental physics from the Event Horizon Telescope image of Sagittarius A**, 2205.07787.
- [78] A.K. Saha, P. Parashari, T.N. Maity, A. Dubey, S. Bouri and R. Laha, *Bounds on ultralight bosons from the Event Horizon Telescope observation of Sgr A**, 2208.03530.
- [79] Y.-D. Tsai, J. Eby, J. Arakawa, D. Farnocchia and M.S. Safronova, *New Constraints on Dark Matter and Cosmic Neutrino Profiles through Gravity*, 2210.03749.
- [80] J.W. Moffat, *Modified gravitational theory as an alternative to dark energy and dark matter*, astro-ph/0403266.
- [81] D. Borka, P. Jovanović, V.B. Jovanović and A.F. Zakharov, *Constraining the range of Yukawa gravity interaction from S2 star orbits*, *JCAP* **11** (2013) 050 [1311.1404].
- [82] A.F. Zakharov, P. Jovanovic, D. Borka and V.B. Jovanovic, *Constraining the range of Yukawa gravity interaction from S2 star orbits II: Bounds on graviton mass*, *JCAP* **05** (2016) 045 [1605.00913].
- [83] Y. Dong, L. Shao, Z. Hu, X. Miao and Z. Wang, *Prospects for constraining the Yukawa gravity with pulsars around Sagittarius A**, *JCAP* **11** (2022) 051 [2210.16130].
- [84] C. de Rham, *Massive Gravity*, *Living Rev. Rel.* **17** (2014) 7 [1401.4173].
- [85] S.J. Ostro et al., *Radar Imaging of Binary Near-Earth Asteroid (66391) 1999 KW₄*, *Science* **314** (2006) 1276.
- [86] J.L. Gustetic, V. Friedensen, J.L. Kessler, S. Jackson and J. Parr, *Nasa’s asteroid grand challenge: Strategy, results, and lessons learned*, *Space Policy* **44-45** (2018) 1–13.
- [87] “The european space agency: Science and exploration.” https://www.esa.int/Science_Exploration/Human_and_Robotic_Exploration/Exploration/Asteroids, 2021.
- [88] S.P. Naidu, L.A.M. Benner, J.-L. Margot, M.W. Busch and P.A. Taylor, *Capabilities of Earth-based Radar Facilities for Near-Earth Asteroid Observations*, *Astrophys. J.* **152** (2016) 99 [1604.01080].

- [89] S.G. Djorgovski et al., *The catalina real-time transient survey (crts)*, in *The First Year of MAXI: Monitoring Variable X-ray Sources*, Tokyo: JAXA Special Publ., 2, 2011 [[1102.5004](#)].
- [90] Vera C. Rubin Observatory LSST Solar System Science Collaboration, R.L. Jones, M.T. Bannister, B.T. Bolin, C.O. Chandler, S.R. Chesley et al., *The Scientific Impact of the Vera C. Rubin Observatory's Legacy Survey of Space and Time (LSST) for Solar System Science*, *Bulletin of the AAS* **53** (2020) [arXiv:2009.07653](#) [[2009.07653](#)].
- [91] B. Carry et al., *Potential asteroid discoveries by the esa gaia mission*, *Astron. Astrophys.* **648** (2021) [A96](#).
- [92] A.K. Verma, J.-L. Margot and A.H. Greenberg, *Prospects of Dynamical Determination of General Relativity Parameter β and Solar Quadrupole Moment $J_{2\odot}$ with Asteroid Radar Astronomy*, *Astrophys. J.* **845** (2017) [166](#) [[1707.08675](#)].
- [93] A.H. Greenberg, J.-L. Margot, A.K. Verma, P.A. Taylor and S.E. Hodge, *Yarkovsky drift detections for 247 near-earth asteroids*, *Astron. J.* **159** (2020) [92](#).
- [94] L. Iorio, *On the effects of the Dvali-Gabadadze-Porrati braneworld gravity on the orbital motion of a test particle*, *Class. Quant. Grav.* **22** (2005) [5271](#) [[gr-qc/0504053](#)].
- [95] S.L. Adler, *Planet-bound dark matter and the internal heat of Uranus, Neptune, and hot-Jupiter exoplanets*, *Phys. Lett. B* **671** (2009) [203](#) [[0808.2823](#)].
- [96] A. Jordan and G.A. Bakos, *Observability of the General Relativistic Precession of Periastra in Exoplanets*, *Astrophys. J.* **685** (2008) [543](#) [[0806.0630](#)].
- [97] L. Iorio, *Classical and relativistic long-term time variations of some observables for transiting exoplanets*, *Mon. Not. Roy. Astron. Soc.* **411** (2011) [167](#) [[1007.2780](#)].
- [98] D. Hooper and J.H. Steffen, *Dark Matter And The Habitability of Planets*, *JCAP* **07** (2012) [046](#) [[1103.5086](#)].
- [99] L. Iorio, *Constraints on a MOND effect for isolated aspherical systems in the deep Newtonian regime from orbital motions*, *Class. Quant. Gravit.* **30** (2013) [165018](#) [[1211.3688](#)].
- [100] J. Overduin, J. Mitcham and Z. Warecki, *Expanded solar-system limits on violations of the equivalence principle*, *Class. Quant. Grav.* **31** (2014) [015001](#) [[1307.1202](#)].
- [101] N.P. Pitjeva and E.V. Pitjeva, *Constraints on dark matter in the solar system*, *Astronomy Letters* **39** (2013) [141](#) [[1306.5534](#)].
- [102] L. Iorio, *Post-Keplerian corrections to the orbital periods of a two-body system and their measurability*, *Mon. Not. Roy. Astron. Soc.* **460** (2016) [2445](#) [[1407.5021](#)].
- [103] A. Ain, S. Kastha and S. Mitra, *Stochastic Gravitational Wave Background from Exoplanets*, *Phys. Rev. D* **91** (2015) [124023](#) [[1504.01715](#)].
- [104] K. Masuda and Y. Suto, *Transiting planets as a precision clock to constrain the time variation of the gravitational constant*, *Publ. Astron. Soc. Jap.* **68** (2016) [L5](#) [[1602.02513](#)].
- [105] L. Blanchet, G. Hébrard and F. Larrouturou, *Detecting the General Relativistic Orbital Precession of the Exoplanet HD 80606b*, *Astron. Astrophys.* **628** (2019) [A80](#) [[1905.06630](#)].
- [106] J. Bramante, A. Buchanan, A. Goodman and E. Lodhi, *Terrestrial and Martian Heat Flow Limits on Dark Matter*, *Phys. Rev. D* **101** (2020) [043001](#) [[1909.11683](#)].
- [107] B. Sun, Z. Cao and L. Shao, *Constraints on fifth forces through perihelion precession of planets*, *Phys. Rev. D* **100** (2019) [084030](#) [[1910.05666](#)].
- [108] R. Garani and P. Tinyakov, *Constraints on Dark Matter from the Moon*, *Phys. Lett. B* **804** (2020) [135403](#) [[1912.00443](#)].
- [109] J. Scholtz and J. Unwin, *What if Planet 9 is a Primordial Black Hole?*, *Phys. Rev. Lett.* **125** (2020) [051103](#) [[1909.11090](#)].

- [110] M.L. Ruggiero and L. Iorio, *Probing a r^{-n} modification of the Newtonian potential with exoplanets*, *JCAP* **06** (2020) 042 [[2001.04122](#)].
- [111] M.H. Chan and C.M. Lee, *Constraining the spin-independent elastic scattering cross section of dark matter using the Moon as a detection target and the background neutrino data*, *Phys. Rev. D* **102** (2020) 023024 [[2007.01589](#)].
- [112] R.K. Leane and J. Smirnov, *Exoplanets as Sub-GeV Dark Matter Detectors*, *Phys. Rev. Lett.* **126** (2021) 161101 [[2010.00015](#)].
- [113] H. Wei and Z.-X. Yu, *Inverse Chameleon Mechanism and Mass Limits for Compact Stars*, *JCAP* **08** (2021) 011 [[2103.12696](#)].
- [114] R.K. Leane and T. Linden, *First Analysis of Jupiter in Gamma Rays and a New Search for Dark Matter*, [2104.02068](#).
- [115] T. Kumar Poddar, S. Mohanty and S. Jana, *Constraints on long range force from perihelion precession of planets in a gauged $L_e - L_{\mu,\tau}$ scenario*, *Eur. Phys. J. C* **81** (2021) 286 [[2002.02935](#)].
- [116] J. Bergé, P. Brax, G. Métris, M. Pernot-Borràs, P. Touboul and J.-P. Uzan, *MICROSCOPE Mission: First Constraints on the Violation of the Weak Equivalence Principle by a Light Scalar Dilaton*, *Phys. Rev. Lett.* **120** (2018) 141101 [[1712.00483](#)].
- [117] C.D. Carone and H. Murayama, *Possible light $U(1)$ gauge boson coupled to baryon number*, *Phys. Rev. Lett.* **74** (1995) 3122 [[hep-ph/9411256](#)].
- [118] P. Fileviez Perez and M.B. Wise, *Baryon and lepton number as local gauge symmetries*, *Phys. Rev. D* **82** (2010) 011901 [[1002.1754](#)].
- [119] A. Davidson, *$B - L$ as the fourth color within an $SU(2)_L \times U(1)_R \times U(1)$ model*, *Phys. Rev. D* **20** (1979) 776.
- [120] R.N. Mohapatra and R.E. Marshak, *Local B-L Symmetry of Electroweak Interactions, Majorana Neutrinos and Neutron Oscillations*, *Phys. Rev. Lett.* **44** (1980) 1316.
- [121] A. Davidson and K.C. Wali, *Universal Seesaw Mechanism?*, *Phys. Rev. Lett.* **59** (1987) 393.
- [122] R. Foot, *New Physics From Electric Charge Quantization?*, *Mod. Phys. Lett. A* **6** (1991) 527.
- [123] X.-G. He, G.C. Joshi, H. Lew and R.R. Volkas, *Simplest Z-prime model*, *Phys. Rev. D* **44** (1991) 2118.
- [124] M. Escudero, D. Hooper, G. Krnjaic and M. Pierre, *Cosmology with A Very Light $L_\mu - L_\tau$ Gauge Boson*, *JHEP* **03** (2019) 071 [[1901.02010](#)].
- [125] N. Blinov, S.A.R. Ellis and A. Hook, *Consequences of Fine-Tuning for Fifth Force Searches*, *JHEP* **11** (2018) 029 [[1807.11508](#)].
- [126] S. Sibiryakov, P. Sørensen and T.-T. Yu, *BBN constraints on universally-coupled ultralight scalar dark matter*, *JHEP* **12** (2020) 075 [[2006.04820](#)].
- [127] E. Izaguirre, G. Krnjaic and M. Pospelov, *Probing New Physics with Underground Accelerators and Radioactive Sources*, *Phys. Lett. B* **740** (2015) 61 [[1405.4864](#)].
- [128] M. Pospelov and Y.-D. Tsai, *Light scalars and dark photons in Borexino and LSND experiments*, *Phys. Lett. B* **785** (2018) 288 [[1706.00424](#)].
- [129] J.A. Dror, R. Lasenby and M. Pospelov, *New constraints on light vectors coupled to anomalous currents*, *Phys. Rev. Lett.* **119** (2017) 141803 [[1705.06726](#)].
- [130] J.A. Dror, R. Lasenby and M. Pospelov, *Dark forces coupled to nonconserved currents*, *Phys. Rev. D* **96** (2017) 075036 [[1707.01503](#)].
- [131] M.B. Green and J.H. Schwarz, *Anomaly Cancellation in Supersymmetric $D=10$ Gauge Theory and Superstring Theory*, *Phys. Lett. B* **149** (1984) 117.

- [132] D.B. Kaplan, *Flavor at SSC energies: A New mechanism for dynamically generated fermion masses*, *Nucl. Phys. B* **365** (1991) 259.
- [133] A. Pierce, K. Riles and Y. Zhao, *Searching for Dark Photon Dark Matter with Gravitational Wave Detectors*, *Phys. Rev. Lett.* **121** (2018) 061102 [[1801.10161](#)].
- [134] M.G. Stewart, *Precession of the perihelion of Mercury's orbit*, *American Journal of Physics* **73** (2005) 730.
- [135] S.R. Kane, J. Horner and K. von Braun, *Cyclic Transit Probabilities of Long-period Eccentric Planets due to Periastron Precession*, *Astrophys. J.* **757** (2012) 105 [[1208.4115](#)].
- [136] L. Iorio, *Advances in the measurement of the Lense-Thirring effect with planetary motions in the field of the Sun*, *Schol. Res. Exch.* **2008** (2008) 105235 [[0807.0435](#)].
- [137] C.W. Misner, K.S. Thorne and J.A. Wheeler, *Gravitation*, W. H. Freeman, San Francisco (1973).
- [138] C.M. Will and K. Nordtvedt, Jr., *Conservation Laws and Preferred Frames in Relativistic Gravity. I. Preferred-Frame Theories and an Extended PPN Formalism*, *Astrophys. J.* **177** (1972) 757.
- [139] C.M. Will, *Theory and Experiment in Gravitational Physics*, Cambridge University Press (9, 2018).
- [140] S.C. Solomon et al., *The MESSENGER mission to Mercury: scientific objectives and implementation*, *Planet. Space Sci.* **49** (2001) 1445.
- [141] B. Bertotti, L. Iess and P. Tortora, *A test of general relativity using radio links with the Cassini spacecraft*, *Nature* **425** (2003) 374.
- [142] D.E. Smith et al., *Gravity Field and Internal Structure of Mercury from MESSENGER*, *Science* **336** (2012) 214.
- [143] L. Barbieri and F. Talamucci, *Calculation of apsidal precession via perturbation theory*, *Advanc. Astrophys.* **4** (2019) 96 [[1802.07115](#)].
- [144] S. Evans et al., *MONTE: the next generation of mission design and navigation software*, *CEAS Space Journal* **10** (2018) 79.
- [145] G. Rosi, F. Sorrentino, L. Cacciapuoti, M. Prevedelli and G.M. Tino, *Precision Measurement of the Newtonian Gravitational Constant Using Cold Atoms*, *Nature* **510** (2014) 518 [[1412.7954](#)].
- [146] P. Hamilton, M. Jaffe, P. Haslinger, Q. Simmons, H. Müller and J. Khoury, *Atom-interferometry constraints on dark energy*, *Science* **349** (2015) 849 [[1502.03888](#)].
- [147] C. Xue, J.-P. Liu, Q. Li, J.-F. Wu, S.-Q. Yang, Q. Liu et al., *Precision measurement of the Newtonian gravitational constant*, *Natl. Sci. Rev.* **7** (2020) 1803.
- [148] E.G. Adelberger, B.R. Heckel, S.A. Hoedl, C.D. Hoyle, D.J. Kapner and A. Upadhye, *Particle Physics Implications of a Recent Test of the Gravitational Inverse Square Law*, *Phys. Rev. Lett.* **98** (2007) 131104 [[hep-ph/0611223](#)].
- [149] P. Fayet, *MICROSCOPE limits on the strength of a new force, with comparisons to gravity and electromagnetism*, *Phys. Rev. D* **99** (2019) 055043 [[1809.04991](#)].
- [150] M. Baryakhtar, M. Galanis, R. Lasenby and O. Simon, *Black hole superradiance of self-interacting scalar fields*, *Phys. Rev. D* **103** (2021) 095019 [[2011.11646](#)].
- [151] T.W. Murphy, *Lunar laser ranging: the millimeter challenge*, *Rept. Prog. Phys.* **76** (2013) 076901 [[1309.6294](#)].
- [152] S.G. Turyshev, W. Farr, W.M. Folkner, A.R. Giererd, H. Hemmati, T.W. Murphy, Jr. et al.,

- Advancing Tests of Relativistic Gravity via Laser Ranging to Phobos*, *Exper. Astron.* **28** (2010) 209 [1003.4961].
- [153] C. Burrage and J. Sakstein, *A Compendium of Chameleon Constraints*, *JCAP* **11** (2016) 045 [1609.01192].
- [154] MICROSCOPE collaboration, *MICROSCOPE Mission: Final Results of the Test of the Equivalence Principle*, *Phys. Rev. Lett.* **129** (2022) 121102 [2209.15487].
- [155] J. Overduin, F. Everitt, P. Worden and J. Mester, *STEP and fundamental physics*, *Class. Quant. Grav.* **29** (2012) 184012 [1401.4784].
- [156] “Jpl small-body database browser - 66391 moshup (1999 kw4).” <https://ssd.jpl.nasa.gov/sbdb.cgi?sstr=1999+KW4&orb=1>, 2021.
- [157] A.H. Greenberg, J.-L. Margot, A.K. Verma, P.A. Taylor, S.P. Naidu, M. Brozovic et al., *Asteroid 1566 Icarus’s Size, Shape, Orbit, and Yarkovsky Drift from Radar Observations*, *Astrophys. J.* **153** (2017) 108 [1612.07434].
- [158] C.B. Olkin et al., *Lucy Mission to the Trojan Asteroids: Instrumentation and Encounter Concept of Operations*, *Planet. Sci.* **2** (2021) 172 [2104.04575].
- [159] L. Denneau et al., *The Pan-STARRS Moving Object Processing System*, *Publ.Astron.Soc.Pac.* **125** (2013) 357 [1302.7281].
- [160] J.L. Tonry et al., *ATLAS: A High-cadence All-sky Survey System*, *Publ.Astron.Soc.Pac.* **130** (2018) 064505 [1802.00879].
- [161] N. Golovich, N. Lifset, R. Armstrong, E. Green, M.D. Schneider and R. Pearce, *A New Blind Asteroid Detection Scheme*, **2104.03411**.
- [162] M.J. Graham et al., *The zwicky transient facility: Science objectives*, *Publ.Astron.Soc.Pac.* **131** (2019) 078001.
- [163] A.A. Parfeni, L.I. Caramete, A.M. Dobre and N. Tran Bach, *Detection of asteroid trails in Hubble Space Telescope images using Deep Learning*, *arXiv e-prints* (2020) arXiv:2010.15425 [2010.15425].
- [164] A.S. Rivkin, F. Marchis, J.A. Stansberry, D. Takir and C. Thomas, *Asteroids and the james webb space telescope*, *Publ. Astron. Soc. Pac.* **128** (2016) 018003.
- [165] B. Carry, *Solar system science with esa euclid*, *Astron. Astrophys.* **609** (2018) A113.
- [166] R. Akeson, L. Armus, E. Bachelet, V. Bailey, L. Bartusek, A. Bellini et al., *The Wide Field Infrared Survey Telescope: 100 Hubbles for the 2020s*, *arXiv e-prints* (2019) [1902.05569].
- [167] C. Lewicki, A. Graps, M. Elvis, P. Metzger and A. Rivkin, *Furthering Asteroid Resource Utilization in the Next Decade through Technology Leadership*, *arXiv e-prints* (2021) [2103.02435].
- [168] D. Seligman and K. Batygin, *The onset of chaos in permanently deformed binaries from spin-orbit and spin-spin coupling*, *The Astrophysical Journal* **913** (2021) 31.
- [169] J.N. Winn, *Transits and Occultations*, *arXiv e-prints* (2010) arXiv:1001.2010 [1001.2010].
- [170] M.A. Fedderke, P.W. Graham and S. Rajendran, *Gravity Gradient Noise from Asteroids*, *Phys. Rev. D* **103** (2021) 103017 [2011.13833].
- [171] M.A. Fedderke, P.W. Graham and S. Rajendran, *Asteroids for μ Hz gravitational-wave detection*, *Phys. Rev. D* **105** (2022) 103018 [2112.11431].
- [172] “Jpl small-body database search engine.” https://ssd.jpl.nasa.gov/sbdb_query.cgi, 2021.
- [173] “The international astronomical union minor planet center.” <https://www.minorplanetcenter.net/>, 2021.

- [174] Y.-D. Tsai *et al.*, *Leading constraints on fifth forces and ultralight dark matter from the OSIRIS-REx mission and ultraprecise asteroid astrometry*, in preparation (2023) .
- [175] P.G. Bergmann, *Comments on the scalar tensor theory*, *Int. J. Theor. Phys.* **1** (1968) 25.
- [176] M. Milgrom, *A modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis.*, *Astrophys. J.* **270** (1983) 365.
- [177] T.P. Sotiriou and V. Faraoni, *f(R) Theories Of Gravity*, *Rev. Mod. Phys.* **82** (2010) 451 [[0805.1726](#)].
- [178] T. Clifton, P.G. Ferreira, A. Padilla and C. Skordis, *Modified Gravity and Cosmology*, *Phys. Rept.* **513** (2012) 1 [[1106.2476](#)].
- [179] S. Nojiri, S.D. Odintsov and V.K. Oikonomou, *Modified Gravity Theories on a Nutshell: Inflation, Bounce and Late-time Evolution*, *Phys. Rept.* **692** (2017) 1 [[1705.11098](#)].
- [180] D.E. Kaplan, G.Z. Krnjaic, K.R. Rehermann and C.M. Wells, *Atomic Dark Matter*, *JCAP* **05** (2010) 021 [[0909.0753](#)].
- [181] Y. Farzan and A.R. Akbarieh, *VDM: A model for Vector Dark Matter*, *JCAP* **10** (2012) 026 [[1207.4272](#)].
- [182] F.-Y. Cyr-Racine and K. Sigurdson, *Cosmology of atomic dark matter*, *Phys. Rev. D* **87** (2013) 103515 [[1209.5752](#)].
- [183] K. Petraki, L. Pearce and A. Kusenko, *Self-interacting asymmetric dark matter coupled to a light massive dark photon*, *JCAP* **07** (2014) 039 [[1403.1077](#)].
- [184] L. Randall and M. Reece, *Dark Matter as a Trigger for Periodic Comet Impacts*, *Phys. Rev. Lett.* **112** (2014) 161301 [[1403.0576](#)].
- [185] R. Foot and S. Vagnozzi, *Dissipative hidden sector dark matter*, *Phys. Rev. D* **91** (2015) 023512 [[1409.7174](#)].
- [186] Y.-D. Tsai, R. McGehee and H. Murayama, *Resonant Self-Interacting Dark Matter from Dark QCD*, *Phys. Rev. Lett.* **128** (2022) 172001 [[2008.08608](#)].
- [187] S. Knapen, T. Lin and K.M. Zurek, *Light Dark Matter: Models and Constraints*, *Phys. Rev. D* **96** (2017) 115021 [[1709.07882](#)].