

Snowmass2021 - Letter of Interest

Cosmology Intertwined IV: The Age of the Universe and its Curvature

Thematic Areas: (check all that apply /■)

- (CF1) Dark Matter: Particle Like
- (CF2) Dark Matter: Wavelike
- (CF3) Dark Matter: Cosmic Probes
- (CF4) Dark Energy and Cosmic Acceleration: The Modern Universe
- (CF5) Dark Energy and Cosmic Acceleration: Cosmic Dawn and Before
- (CF6) Dark Energy and Cosmic Acceleration: Complementarity of Probes and New Facilities
- (CF7) Cosmic Probes of Fundamental Physics
- (Other) [*Please specify frontier/topical group*]

Contact Information:

Eleonora Di Valentino (JBCA, University of Manchester, UK) [eleonora.divalentino@manchester.ac.uk]

Authors:

Eleonora Di Valentino (JBCA, University of Manchester, UK)
Luis A. Anchordoqui (City University of New York, USA)
Özgür Akarsu (Istanbul Technical University, Istanbul, Turkey)
Yacine Ali-Haïmoud (New York University, USA)
Luca Amendola (University of Heidelberg, Germany)
Nikki Arendse (DARK, Niels Bohr Institute, Denmark)
Marika Asgari (University of Edinburgh, UK)
Mario Ballardini (Alma Mater Studiorum Università di Bologna, Italy)
Spyros Basilakos (Academy of Athens and Nat. Observatory of Athens, Greece)
Elia Battistelli (Sapienza Università di Roma and INFN sezione di Roma, Italy)
Micol Benetti (Università degli Studi di Napoli Federico II and INFN sezione di Napoli, Italy)
Simon Birrer (Stanford University, USA)
François R. Bouchet (Institut d'Astrophysique de Paris, CNRS & Sorbonne University, France)
Marco Bruni (Institute of Cosmology and Gravitation, Portsmouth, UK, and INFN Sezione di Trieste, Italy)
Erminia Calabrese (Cardiff University, UK)
David Camarena (Federal University of Espirito Santo, Brazil)
Salvatore Capozziello (Università degli Studi di Napoli Federico II, Napoli, Italy)
Angela Chen (University of Michigan, Ann Arbor, USA)
Jens Chluba (JBCA, University of Manchester, UK)
Anton Chudaykin (Institute for Nuclear Research, Russia)
Eoin Ó Colgáin (Asia Pacific Center for Theoretical Physics, Korea)
Francis-Yan Cyr-Racine (University of New Mexico, USA)
Paolo de Bernardis (Sapienza Università di Roma and INFN sezione di Roma, Italy)
Javier de Cruz Pérez (Departament FQA and ICCUB, Universitat de Barcelona, Spain)

Jacques Delabrouille (CNRS/IN2P3, Laboratoire APC, France & CEA/IRFU, France & USTC, China)
Celia Escamilla-Rivera (ICN, Universidad Nacional Autónoma de México, Mexico)
Agnès Ferté (JPL, Caltech, Pasadena, USA)
Fabio Finelli (INAF OAS Bologna and INFN Sezione di Bologna, Italy)
Wendy Freedman (University of Chicago, Chicago IL, USA)
Noemi Frusciante (Instituto de Astrofísica e Ciências do Espaço, Lisboa, Portugal)
Elena Giusarma (Michigan Technological University, USA)
Adrià Gómez-Valent (University of Heidelberg, Germany)
Will Handley (University of Cambridge, UK)
Ian Harrison (JBCA, University of Manchester, UK)
Luke Hart (JBCA, University of Manchester, UK)
Alan Heavens (ICIC, Imperial College London, UK)
Hendrik Hildebrandt (Ruhr-University Bochum, Germany)
Daniel Holz (University of Chicago, Chicago IL, USA)
Dragan Huterer (University of Michigan, Ann Arbor, USA)
Mikhail M. Ivanov (New York University, USA)
Shahab Joudaki (University of Oxford, UK and University of Waterloo, Canada)
Marc Kamionkowski (Johns Hopkins University, Baltimore, MD, USA)
Tanvi Karwal (University of Pennsylvania, Philadelphia, USA)
Lloyd Knox (UC Davis, Davis CA, USA)
Suresh Kumar (BITS Pilani, Pilani Campus, India)
Luca Lamagna (Sapienza Università di Roma and INFN sezione di Roma, Italy)
Julien Lesgourgues (RWTH Aachen University)
Matteo Lucca (Université Libre de Bruxelles, Belgium)
Valerio Marra (Federal University of Espirito Santo, Brazil)
Silvia Masi (Sapienza Università di Roma and INFN sezione di Roma, Italy)
Sabino Matarrese (University of Padova and INFN Sezione di Padova, Italy)
Arindam Mazumdar (Centre for Theoretical Studies, IIT Kharagpur, India)
Alessandro Melchiorri (Sapienza Università di Roma and INFN sezione di Roma, Italy)
Olga Mena (IFIC, CSIC-UV, Spain)
Laura Mersini-Houghton (University of North Carolina at Chapel Hill, USA)
Vivian Miranda (University of Arizona, USA)
Cristian Moreno-Pulido (Departament FQA and ICCUB, Universitat de Barcelona, Spain)
David F. Mota (University of Oslo, Norway)
Jessica Muir (KIPAC, Stanford University, USA)
Ankan Mukherjee (Jamia Millia Islamia Central University, India)
Florian Niedermann (CP3-Origins, University of Southern Denmark)
Alessio Notari (ICCUB, Universitat de Barcelona, Spain)
Rafael C. Nunes (National Institute for Space Research, Brazil)
Francesco Pace (JBCA, University of Manchester, UK)
Andronikos Paliathanasis (DUT, South Africa and UACH, Chile)
Antonella Palmese (Fermi National Accelerator Laboratory, USA)
Supriya Pan (Presidency University, Kolkata, India)
Daniela Paoletti (INAF OAS Bologna and INFN Sezione di Bologna, Italy)
Valeria Pettorino (AIM, CEA, CNRS, Université Paris-Saclay, Université de Paris, France)
Francesco Piacentini (Sapienza Università di Roma and INFN sezione di Roma, Italy)
Vivian Poulin (LUPM, CNRS & University of Montpellier, France)
Marco Raveri (University of Pennsylvania, Philadelphia, USA)

Adam G. Riess (Johns Hopkins University, Baltimore, USA)
Vincenzo Salzano (University of Szczecin, Poland)
Emmanuel N. Saridakis (National Observatory of Athens, Greece)
Anjan A. Sen (Jamia Millia Islamia Central University New Delhi, India)
Arman Shafieloo (Korea Astronomy and Space Science Institute (KASI), Korea)
Anowar J. Shajib (University of California, Los Angeles, USA)
Joseph Silk (IAP Sorbonne University & CNRS, France, and Johns Hopkins University, USA)
Alessandra Silvestri (Leiden University, NL)
Martin S. Sloth (CP3-Origins, University of Southern Denmark)
Tristan L. Smith (Swarthmore College, Swarthmore, USA)
Joan Solà Peracaula (Departament FQA and ICCUB, Universitat de Barcelona, Spain)
Carsten van de Bruck (University of Sheffield, UK)
Licia Verde (ICREA, Universidad de Barcelona, Spain)
Luca Visinelli (GRAPPA, University of Amsterdam, NL)
Benjamin D. Wandelt (IAP Sorbonne University & CNRS, France, and CCA, USA)
Deng Wang (National Astronomical Observatories, CAS, China)
Jian-Min Wang (Key Laboratory for Particle Astrophysics, IHEP of the CAS, Beijing, China)
Anil K. Yadav (United College of Engg. & Research, GN, India)
Weiqiang Yang (Liaoning Normal University, Dalian, China)

Abstract: A precise measurement of the curvature of the Universe is of primeval importance for cosmology since it could not only confirm the paradigm of primordial inflation but also help in discriminating between different early Universe scenarios. The recent observations, while broadly consistent with a spatially flat standard Λ Cold Dark Matter (Λ CDM) model, are showing tensions that still allow (and, in some cases, even suggest) a few percent deviations from a flat universe. In particular, the Planck Cosmic Microwave Background power spectra, assuming the nominal likelihood, prefer a closed universe at more than 99% confidence level. While new physics could be in action, this anomaly may be the result of an unresolved systematic error or just a statistical fluctuation. However, since a positive curvature allows a larger age of the Universe, an accurate determination of the age of the oldest objects provides a smoking gun in confirming or falsifying the current flat Λ CDM model.

The curvature of the Universe – The flat Λ Cold Dark Matter (Λ CDM) cosmological model describes incredibly well the current cosmological observations. However, together with the long standing *Hubble constant* H_0 disagreement¹, and $\sigma_8 - S_8$ tension², there are some anomalies in the Planck 2018 cosmological results that deserve further investigations. Between them the most significant from the statistical point of view, is the preference at 3.4σ for a closed Universe^{3–5}. Moreover, the Planck dataset also suggest an indication at more than 2σ for Modified Gravity^{4;6;7}. This disagreement with the predictions for a flat universe of the standard model is connected with the higher, anomalous, lensing contribution in the Cosmic Microwave Background (CMB) power spectra, characterized by the A_L parameter^{4;8}, that is strongly degenerate with Ω_k (see Fig. 1). A closed universe also solves a well-know tension above 2σ between the low and high multipoles regions of the angular power spectra^{3;9;10}. This indication for curvature can be due to unresolved systematics in the Planck 2018 data, or can be simply due to a statistical fluctuation.

Indeed, while Planck 2018⁴ finds $\Omega_k = -0.044^{+0.0181}_{-0.015}$, i.e. $\Omega_k < 0$ at about 3.4σ ($\Delta\chi^2 \sim -11$), using the official baseline Plik likelihood¹¹, the evidence is reduced when considering the alternative CamSpec¹² likelihood (see discussion in¹³), albeit with the marginalized constraint still above the 99% CL ($\Omega_k = -0.035^{+0.018}_{-0.013}$). Moreover, the recent results from the ground-based experiment ACT, in combination with data from the WMAP experiment, is fully compatible with a flat universe with $\Omega_k = -0.001^{+0.014}_{-0.010}$ while slightly preferring a closed universe when combined with a portion of the Planck dataset with $\Omega_k = -0.018^{+0.013}_{-0.010}$ ¹⁴ (see Fig. 2). A closed universe is also preferred by a combination of non-CMB data made by Baryon Acoustic Oscillation (BAO) measurements^{15–17}, supernovae (SNe) distances from the recent Pantheon catalog¹⁸, and a prior on the baryon density derived from measurements of primordial deuterium¹⁹ assuming Big Bang Nucleosynthesis (BBN), but with a much larger H_0 ³ completely in agreement with the SH0ES collaboration value R19²⁰. However, letting the curvature free to vary means to increase both the H_0 and the S_8 tensions³. Therefore, at the moment there are not theoretical models that can explain at the same time all the tensions and anomalies we see in the data. On the other hand, a flat universe is preferred also by Planck + BAO, or + CMB lensing²¹ or + Pantheon data. However these dataset combinations are in disagreement at more than 3σ when the curvature is free to vary^{3;5}. In addition, though the error bars are so large that cannot discriminate between the models, a flat Universe is also in agreement with the analysis made by²² using the $H(z)$ sample from the cosmic chronometers (CC) and the luminosity distance $D_L(z)$ from the 1598 quasars ($\Omega_k = 0.08 \pm 0.31$) or the Pantheon sample ($\Omega_k = -0.02 \pm 0.14$), in agreement with the previous²³. Finally, in²⁴ a combination of BAO+BBN+H0LiCOW provides $\Omega_k = -0.07^{+0.14}_{-0.26}$ with H_0 in agreement with R19, while BAO+BBN+CC gives a positive $\Omega_k = 0.28^{+0.17}_{-0.28}$. In¹³ it has been pointed out that is difficult to believe to a possible cosmological data conspiracy towards $\Omega_k = 0$. However, a full agreement of the luminosity distance measurements, like Pantheon or R19, with Planck can be reached also ruling out both, a flat universe and a cosmological constant²⁵.

The Age of the Universe – The age of the universe is an important piece of the puzzle because it connects H_0 and Ω_m , both of which can be measured in the early and the late universe. The age is not just a




Figure 1: 68% CL and 95% CL contour plots for Ω_k and A_L (from Ref.³).

¹All the bounds are reported at 68% confidence level in the text.

prediction of the Λ CDM model, that for Planck 2018 is $t_U = 13.800 \pm 0.024$ Gyr, but can also be measured using very old objects. For example, in²⁶ it is obtained $t_U = 13.35 \pm 0.16(\text{stat.}) \pm 0.5(\text{sys.})$ Gyr using populations of stars in globular clusters. Nevertheless, while robustness and accuracy tests have been done very extensively for CMB and quite extensively for BAO and SNe, for the age of the oldest objects we have to be more careful. For example, one finds the ages of the oldest stars 2MASS J18082002–5104378 B equal to $t_* = 13.535 \pm 0.002$ Gyr²⁷, but if the scatter among different models to fit for the age is taken into account the age becomes $t_* = 13.0 \pm 0.6$ Gyr²⁸, and the age of HD 140283 equal to $t_* = 14.46 \pm 0.8$ Gyr²⁹, but becomes $t_* = 13.5 \pm 0.7$ Gyr²⁸ using the new Gaia parallaxes instead of original HST parallaxes. Therefore, even if at present there is not real tension between the different t_u determinations, most of the error-bars in the age determination comes from the fact that different stellar models do not really agree with each other at the required level of precision to be really able to help with the tensions in cosmology. Nevertheless, stellar models can/are expected to improve reducing this error significantly, and this could potentially unveil a tension on the age of the Universe. Trying to alleviate it by changing the Planck determination, would interestingly have an effect on the cosmological tensions. A possibility to increase the age of the Universe, to be larger than the age of oldest stars, is by lowering the Hubble constant value, because of the anti-correlation between these parameters³⁰. For example, a positive curvature for the Universe, as suggested by Planck 2018, preferring a lower H_0 and worsening significantly the H_0 tension, predicts an older Universe $t_U = 15.31 \pm 0.47$ Gyr. Therefore, it seems that the only way to address the H_0 crisis if R19 is correct is to introduce an extremely recent (after $z \sim 0.1$) departure from Λ CDM, requiring a great deal of fine-tuning.

Future – Detecting a curvature Ω_k different from zero could be due to a local inhomogeneity biasing our bounds³¹, and in this case CMB spectral distortions such as the KSZ effect and Compton- γ distortions, present a viable method to constrain the curvature at a level potentially detectable by a next-generation experiment. If a curvature Ω_k different from zero is the evidence for a truly superhorizon departure from flatness, this will have profound implication for a broad class of inflationary scenarios. While open universe are easier to obtain in inflationary models^{32–36}, with a fine-tuning at the level of about one percent one can obtain also a semi-realistic model of a closed inflationary universe^{37;38}. In³⁹ it has been shown that forthcoming surveys, even combined together, are likely to place constraints on the spatial curvature of $\sim 10^{-3}$ at 95% CL at best, but enough for solving the current anomaly in the Planck data. Experiments like Euclid and SKA, instead, may further produce tighter measurements of Ω_k by helping to break parameter degeneracies^{40;41}.

Summary – In these four LoIs^{1;2;42} we presented a snapshot, at the beginning of the SNOWMASS process, of the concordance Λ CDM model and its connections with the experiment. This is a cutting-edge field in the area of cosmology, with unrestrained growth over the last decade. On the experimental side, we have learned that it is really important to have multiple precise and robust measurements of the same observable, with experiments conducted blind in regard to the expected outcome. This provides a unique opportunity to study similar physics from various points of view. While on the theory side, it is really important having robust and testable predictions for the proposed physical models that can be probed with the data. With the synergy between these two sides, significant progress can be made to answer fundamental physics questions. During the SNOWMASS process we plan to monitor the new advances on the field to come out with a clear roadmap for the coming decades.




Figure 2: 1D posterior distributions on Ω_k (from Ref. 14).

References

- [1] E. Di Valentino *et al.*, “Cosmology Intertwined II: The Hubble Constant Tension,” [arXiv:2008.11284](#) [[astro-ph.CO](#)].
- [2] E. Di Valentino *et al.*, “Cosmology Intertwined III: $f\sigma_8$ and S_8 ,” [arXiv:2008.11285](#) [[astro-ph.CO](#)].
- [3] E. Di Valentino, A. Melchiorri, and J. Silk, “Planck evidence for a closed Universe and a possible crisis for cosmology,” *Nature Astron.* **4** no. 2, (2019) 196–203, [arXiv:1911.02087](#) [[astro-ph.CO](#)].
- [4] **Planck** Collaboration, N. Aghanim *et al.*, “Planck 2018 results. VI. Cosmological parameters,” [arXiv:1807.06209](#) [[astro-ph.CO](#)].
- [5] W. Handley, “Curvature tension: evidence for a closed universe,” [arXiv:1908.09139](#) [[astro-ph.CO](#)].
- [6] **Planck** Collaboration, P. A. R. Ade *et al.*, “Planck 2015 results. XIV. Dark energy and modified gravity,” *Astron. Astrophys.* **594** (2016) A14, [arXiv:1502.01590](#) [[astro-ph.CO](#)].
- [7] S. Capozziello, “Curvature quintessence,” *Int. J. Mod. Phys. D* **11** (2002) 483–492, [arXiv:gr-qc/0201033](#).
- [8] E. Calabrese, A. Slosar, A. Melchiorri, G. F. Smoot, and O. Zahn, “Cosmic Microwave Weak lensing data as a test for the dark universe,” *Phys. Rev.* **D77** (2008) 123531, [arXiv:0803.2309](#) [[astro-ph](#)].
- [9] G. Addison, Y. Huang, D. Watts, C. Bennett, M. Halpern, G. Hinshaw, and J. Weiland, “Quantifying discordance in the 2015 Planck CMB spectrum,” *Astrophys. J.* **818** no. 2, (2016) 132, [arXiv:1511.00055](#) [[astro-ph.CO](#)].
- [10] **Planck** Collaboration, N. Aghanim *et al.*, “Planck intermediate results. LI. Features in the cosmic microwave background temperature power spectrum and shifts in cosmological parameters,” *Astron. Astrophys.* **607** (2017) A95, [arXiv:1608.02487](#) [[astro-ph.CO](#)].
- [11] **Planck** Collaboration, N. Aghanim *et al.*, “Planck 2018 results. V. CMB power spectra and likelihoods,” [arXiv:1907.12875](#) [[astro-ph.CO](#)].
- [12] G. Efstathiou and S. Gratton, “A Detailed Description of the CamSpec Likelihood Pipeline and a Reanalysis of the Planck High Frequency Maps,” [arXiv:1910.00483](#) [[astro-ph.CO](#)].
- [13] G. Efstathiou and S. Gratton, “The evidence for a spatially flat Universe,” [arXiv:2002.06892](#) [[astro-ph.CO](#)].
- [14] **ACT** Collaboration, S. Aiola *et al.*, “The Atacama Cosmology Telescope: DR4 Maps and Cosmological Parameters,” [arXiv:2007.07288](#) [[astro-ph.CO](#)].
- [15] F. Beutler, C. Blake, M. Colless, D. H. Jones, L. Staveley-Smith, L. Campbell, Q. Parker, W. Saunders, and F. Watson, “The 6dF Galaxy Survey: Baryon Acoustic Oscillations and the Local Hubble Constant,” *Mon. Not. Roy. Astron. Soc.* **416** (2011) 3017–3032, [arXiv:1106.3366](#) [[astro-ph.CO](#)].

- [16] A. J. Ross, L. Samushia, C. Howlett, W. J. Percival, A. Burden, and M. Manera, “The clustering of the SDSS DR7 main Galaxy sample – I. A 4 per cent distance measure at $z = 0.15$,” *Mon. Not. Roy. Astron. Soc.* **449** no. 1, (2015) 835–847, [arXiv:1409.3242 \[astro-ph.CO\]](#).
- [17] BOSS Collaboration, S. Alam *et al.*, “The clustering of galaxies in the completed SDSS-III Baryon Oscillation Spectroscopic Survey: cosmological analysis of the DR12 galaxy sample,” *Mon. Not. Roy. Astron. Soc.* **470** no. 3, (2017) 2617–2652, [arXiv:1607.03155 \[astro-ph.CO\]](#).
- [18] D. M. Scolnic *et al.*, “The Complete Light-curve Sample of Spectroscopically Confirmed SNe Ia from Pan-STARRS1 and Cosmological Constraints from the Combined Pantheon Sample,” *Astrophys. J.* **859** no. 2, (2018) 101, [arXiv:1710.00845 \[astro-ph.CO\]](#).
- [19] R. J. Cooke, M. Pettini, and C. C. Steidel, “One Percent Determination of the Primordial Deuterium Abundance,” *Astrophys. J.* **855** no. 2, (2018) 102, [arXiv:1710.11129 \[astro-ph.CO\]](#).
- [20] A. G. Riess, S. Casertano, W. Yuan, L. M. Macri, and D. Scolnic, “Large Magellanic Cloud Cepheid Standards Provide a 1% Foundation for the Determination of the Hubble Constant and Stronger Evidence for Physics beyond Λ CDM,” *Astrophys. J.* **876** no. 1, (2019) 85, [arXiv:1903.07603 \[astro-ph.CO\]](#).
- [21] Planck Collaboration, N. Aghanim *et al.*, “Planck 2018 results. VIII. Gravitational lensing,” [arXiv:1807.06210 \[astro-ph.CO\]](#).
- [22] Y. Liu, S. Cao, T. Liu, X. Li, S. Geng, Y. Lian, and W. Guo, “Model-independent constraints on cosmic curvature: implication from updated Hubble diagram of high-redshift standard candles,” [arXiv:2008.08378 \[astro-ph.CO\]](#).
- [23] R.-G. Cai, Z.-K. Guo, and T. Yang, “Null test of the cosmic curvature using $H(z)$ and supernovae data,” *Phys. Rev. D* **93** no. 4, (2016) 043517, [arXiv:1509.06283 \[astro-ph.CO\]](#).
- [24] R. C. Nunes and A. Bernui, “ θ_{BAO} estimates and the H_0 tension,” [arXiv:2008.03259 \[astro-ph.CO\]](#).
- [25] E. Di Valentino, A. Melchiorri, and J. Silk, “Cosmic Discordance: Planck and luminosity distance data exclude LCDM,” [arXiv:2003.04935 \[astro-ph.CO\]](#).
- [26] D. Valcin, J. L. Bernal, R. Jimenez, L. Verde, and B. D. Wandelt, “Inferring the Age of the Universe with Globular Clusters,” [arXiv:2007.06594 \[astro-ph.CO\]](#).
- [27] K. C. Schlafman, I. B. Thompson, and A. R. Casey, “An ultra metal-poor star near the hydrogen-burning limit,” *The Astrophysical Journal* **867** no. 2, (Nov, 2018) 98. <http://dx.doi.org/10.3847/1538-4357/aadd97>.
- [28] R. Jimenez, A. Cimatti, L. Verde, M. Moresco, and B. Wandelt, “The local and distant Universe: stellar ages and H_0 ,” *JCAP* **03** (2019) 043, [arXiv:1902.07081 \[astro-ph.CO\]](#).
- [29] H. E. Bond, E. P. Nelan, D. A. VandenBerg, G. H. Schaefer, and D. Harmer, “Hd 140283: A star in the solar neighborhood that formed shortly after the big bang,” *The Astrophysical Journal* **765** no. 1, (Feb, 2013) L12. <http://dx.doi.org/10.1088/2041-8205/765/1/L12>.
- [30] F. de Bernardis, A. Melchiorri, L. Verde, and R. Jimenez, “The Cosmic Neutrino Background and the Age of the Universe,” *JCAP* **03** (2008) 020, [arXiv:0707.4170 \[astro-ph\]](#).

- [31] P. Bull and M. Kamionkowski, “What if Planck’s Universe isn’t flat?,” *Phys. Rev. D* **87** no. 8, (2013) 081301, [arXiv:1302.1617 \[astro-ph.CO\]](#). [Erratum: *Phys.Rev.D* 87, 129901 (2013)].
- [32] J. Gott, “Creation of open universes from de Sitter space.,” *Nature* **295** (1982) 304–307.
- [33] A. D. Linde, “Quantum creation of an open inflationary universe,” *Phys. Rev. D* **58** (1998) 083514, [arXiv:gr-qc/9802038](#).
- [34] M. Bucher, A. S. Goldhaber, and N. Turok, “An open universe from inflation,” *Phys. Rev. D* **52** (1995) 3314–3337, [arXiv:hep-ph/9411206](#).
- [35] M. Kamionkowski, B. Ratra, D. N. Spergel, and N. Sugiyama, “CBR anisotropy in an open inflation, CDM cosmogony,” *Astrophys. J. Lett.* **434** (1994) L1–L4, [arXiv:astro-ph/9406069](#).
- [36] M. Kamionkowski and D. N. Spergel, “Large angle cosmic microwave background anisotropies in an open universe,” *Astrophys. J.* **432** (1994) 7, [arXiv:astro-ph/9312017](#).
- [37] A. D. Linde, “Can we have inflation with $\Omega_k \neq 0$?,” *JCAP* **05** (2003) 002, [arXiv:astro-ph/0303245](#).
- [38] A. Lasenby and C. Doran, “Closed universes, de sitter space, and inflation,” *Phys. Rev. D* **71** (Mar, 2005) 063502. <https://link.aps.org/doi/10.1103/PhysRevD.71.063502>.
- [39] C. D. Leonard, P. Bull, and R. Allison, “Spatial curvature endgame: Reaching the limit of curvature determination,” *Phys. Rev. D* **94** no. 2, (2016) 023502, [arXiv:1604.01410 \[astro-ph.CO\]](#).
- [40] E. Di Dio, F. Montanari, A. Raccanelli, R. Durrer, M. Kamionkowski, and J. Lesgourgues, “Curvature constraints from Large Scale Structure,” *JCAP* **06** (2016) 013, [arXiv:1603.09073 \[astro-ph.CO\]](#).
- [41] M. Vardanyan, R. Trotta, and J. Silk, “How flat can you get? a model comparison perspective on the curvature of the universe,” *Monthly Notices of the Royal Astronomical Society* **397** no. 1, (Jul, 2009) 431–444. <http://dx.doi.org/10.1111/j.1365-2966.2009.14938.x>.
- [42] E. Di Valentino *et al.*, “Cosmology Intertwined I: Perspectives for the Next Decade,” [arXiv:2008.11283 \[astro-ph.CO\]](#).