

The Lick AGN Monitoring Project 2016: Velocity-Resolved H Lags in Luminous Seyfert Galaxies

VIVIAN U,^{1,2} AARON J. BARTH,¹ H. ALEXANDER VOGLER,^{1,3} HENGXIAO GUO,¹ TOMMASO TREU,⁴
VARDHA N. BENNETT,⁵ GABRIELA CANALIZO,² ALEXEI V. FILIPPENKO,^{6,7} ELINOR GATES,⁸ FREDERICK HAMANN,²
MICHAEL D. JONER,⁹ MATTHEW A. MALKAN,⁴ ANNA PANCOAST,¹⁰ PETER R. WILLIAMS,⁴ JONG-HAK WOO,^{11,12}
BELA ABOLFATHI,¹ L. E. ABRAMSON,¹³ STEPHEN F. ARMEN,¹⁴ HYUN-JIN BAE,¹¹ THOMAS BOHN,²
BENJAMIN D. BOIZELLE,^{9,1} AZALEE BOSTROEM,^{3,15} ANDREW BRANDEL,¹ THOMAS G. BRINK,⁶ SANYUM CHANNA,^{16,17}
M. C. COOPER,¹ MAREN COSENS,^{5,18,19} EDWARD DONOHUE,^{5,20} SEAN P. FILLINGHAM,¹
DIEGO GONZÁLEZ-BUITRAGO,^{1,21} GONI HALEVI,^{6,22} ANDREW HALLE,¹⁶ CAROL E. HOOD,²³ KEITH HORNE,²⁴
J. CHUCK HORST,¹⁴ MAXIME DE KOUCHKOVSKY,⁶ BENJAMIN KUHN,^{25,14} SAHANA KUMAR,^{6,26} DOUGLAS C. LEONARD,¹⁴
DONALD LOVELAND,^{5,27} CHRISTINA MANZANO-KING,² IAN MCHARDY,²⁸ RAÚL MICHEL,²⁹ MELANIE KAE B. OLAES,¹⁴
DAESEONG PARK,^{30,31} SONGYOUN PARK,¹¹ LIUYI PEI,¹ TIMOTHY W. ROSS,⁶ JORDAN N. RUNCO,⁴ JENNA SAMUEL,^{32,3}
JAVIER SÁNCHEZ,^{1,33,34} BRYAN SCOTT,² REMINGTON O. SEXTON,^{2,35,36} JAEJIN SHIN,¹¹ ISAAC SHIVVERS,⁶
CHANCE L. SPENCER,^{5,37} BENJAMIN E. STAHL,^{6,16} SAMANTHA STEGMAN,^{6,38} ISAK STOMBERG,^{5,39} STEFANO VALENTI,³
L. VILLAFANA,⁴ JONELLE L. WALSH,⁴⁰ HEECHAN YUK,^{6,41} AND WEIKANG ZHENG⁶

¹Department of Physics and Astronomy, 4129 Frederick Reines Hall, University of California, Irvine, CA 92697, USA

²Department of Physics and Astronomy, University of California, Riverside, CA 92521, USA

³Department of Physics and Astronomy, University of California, 1 Shields Avenue, Davis, CA 95616, USA

⁴Department of Physics and Astronomy, University of California, Los Angeles, CA 90095-1547, USA

⁵Physics Department, California Polytechnic State University, San Luis Obispo CA 93407, USA

⁶Department of Astronomy, University of California, 501 Campbell Hall, Berkeley, CA 94720-3411, USA

⁷Miller Institute for Basic Research in Science, University of California, Berkeley, CA 94720, USA

⁸Lick Observatory, P.O. Box 85, Mt. Hamilton, CA 95140, USA

⁹Department of Physics and Astronomy, N283 ESC, Brigham Young University, Provo, UT 84602, USA

¹⁰Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

¹¹Astronomy Program, Department of Physics and Astronomy, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Korea

¹²SNU Astronomy Research Center, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Republic of Korea

¹³Carnegie Observatories, 813 Santa Barbara Street, Pasadena, CA 91101, USA

¹⁴Department of Astronomy, San Diego State University, San Diego, CA 92182-1221, USA

¹⁵DiRAC Institute, Department of Astronomy, University of Washington, 3910 15th Avenue, NE, Seattle, WA 98195, USA

¹⁶Department of Physics, University of California, Berkeley, CA 94720, USA

¹⁷Department of Physics, Stanford University, Stanford, CA 94305, USA

¹⁸Physics Department, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093 USA

¹⁹Center for Astrophysics and Space Sciences, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093 USA

²⁰Booz Allen, 1615 Murray Canyon Road, Suite 8000, San Diego, CA 92108, USA

²¹Universidad Nacional Autónoma de México, Instituto de Astronomía, AP 106, Ensenada 22860, BC, México

²²Department of Astrophysical Sciences, Princeton University, 4 Ivy Lane, Princeton, NJ 08544, USA

²³Department of Physics, California State University, San Bernardino, 5500 University Parkway, San Bernardino, CA 92407, USA

²⁴SUPA Physics and Astronomy, University of St Andrews, North Haugh, St Andrews, KY16 9SS, Scotland, UK

²⁵Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

²⁶Department of Physics, Florida State University, 77 Chieftan Way, Tallahassee, FL 32306, USA

²⁷Lawrence Livermore National Laboratory, 7000 East Avenue, Livermore, CA 94550

²⁸University of Southampton, Highfield, Southampton, SO17 1BJ, UK

²⁹Instituto de Astronomía, Universidad Nacional Autónoma de México, AP 877, Ensenada, Baja California, C.P. 22830 México

³⁰Department of Astronomy and Atmospheric Sciences, Kyungpook National University, Daegu, 41566, Republic of Korea

³¹Korea Astronomy and Space Science Institute, Daejeon, 34055, Republic of Korea

³²Department of Astronomy, The University of Texas at Austin, 2515 Speedway, Stop C1400, Austin, TX 78712, USA

³³Fermi National Accelerator Laboratory, Kirk Rd. & Pine St, Batavia, IL 60510, USA

³⁴Kauli Institute for Cosmological Physics, 5640 South Ellis Avenue, Chicago, IL 60637, USA

Corresponding author: Vivian U

vivianu@uci.edu

³⁵ *U.S. Naval Observatory, 3450 Massachusetts Ave NW, Washington, DC 20392-5420, USA*

³⁶ *Department of Physics and Astronomy, George Mason University, 4400 University Dr, Fairfax, VA 22030-4444, USA*

³⁷ *Department of Physics, California State University Fresno, Fresno, CA 93740-8031, USA*

³⁸ *Department of Chemistry, University of Wisconsin, Madison, WI 53706, USA*

³⁹ *Deutsches Elektronen-Synchrotron DESY, 22607 Hamburg, Germany*

⁴⁰ *George P. and Cynthia W. Mitchell Institute for Fundamental Physics and Astronomy, Department of Physics & Astronomy, Texas A&M University, 4242 TAMU, College Station, TX 77843, USA*

⁴¹ *Department of Physics and Astronomy, University of Oklahoma, 440 W. Brooks St., Norman, OK 73019, USA*

ABSTRACT

We carried out spectroscopic monitoring of 21 low-redshift Seyfert 1 galaxies using the Kast double spectrograph on the 3 m Shane telescope at Lick Observatory from April 2016 to May 2017. Targeting active galactic nuclei (AGN) with luminosities of $L \sim 10^{44}$ erg s⁻¹ and predicted H β lags of 20–30 days or black hole masses of 10^7 – $10^8 M_\odot$, our campaign probes luminosity-dependent trends in broad-line region (BLR) structure and dynamics as well as to improve calibrations for single-epoch estimates of quasar black hole masses. Here we present the first results from the campaign, including H β emission-line light curves, integrated H β lag times (8–30 days) measured against V-band continuum light curves, velocity-resolved reverberation lags, line widths of the broad H β components, and virial black hole mass estimates (10^7 – $10^8 M_\odot$). Our results add significantly to the number of existing velocity-resolved lag measurements and reveal a diversity of BLR gas kinematics at moderately high AGN luminosities. AGN continuum luminosity appears not to be correlated with the type of kinematics that its BLR gas may exhibit. Follow-up direct modeling of this dataset will elucidate the detailed kinematics and provide robust dynamical black hole masses for several objects in this sample.

Keywords: Seyfert galaxies, supermassive black holes, active galactic nuclei, reverberation mapping

1. INTRODUCTION

It has been known for about two decades that black hole (BH) mass exhibits a tight correlation with the stellar velocity dispersion of the galactic bulge within which it resides, and this relation holds over several orders of magnitude in black hole mass (e.g., Ferrarese & Merritt 2000; Gebhardt et al. 2000). This relation suggests that supermassive black holes and their host galaxies are tightly linked throughout their lifetimes. A fundamental understanding of the growth of supermassive black holes and their interaction with their immediate vicinity will provide key constraints on cosmological models of galaxy evolution (Springel et al. 2005; Croton et al. 2006). A crucial component in these numerical recipes is therefore the accurate determination of black hole masses as a function of cosmic history.

Robust black hole mass measurements have been limited mostly to nearby galaxies owing to the exquisite angular resolution needed to spatially resolve a black hole’s sphere of influence. The technique of temporally resolving the structure of the broad-line region (BLR) around an actively accreting black hole, called reverberation mapping (RM; Blandford & McKee 1982; Peterson 1993), provides a viable alternative—or in the case of the distant universe, the only currently available tool for directly measuring black hole masses.

Black hole scaling relations, such as that between the BLR size and the active galactic nuclei (AGN) rest-frame luminosity at 5100 Å (the radius luminosity relationship; Wandel et al. 1999; Kaspi et al. 2000, 2005; Bentz et al. 2006, 2009a, 2013; Shen 2013), enable single-epoch mass-determination methods for estimating black hole masses in broad-lined AGN out to high redshifts. The calibrations for such mass estimates, however, rest upon the presumptions that distant quasars are similar to the low-redshift reverberation-mapped AGN, while in fact they exhibit higher luminosities and larger Eddington ratios (Richards et al. 2011; Shen 2013). Whether virial assumptions for BLR gas dynamics based on local Seyferts can be extended to distant quasars must be tested against a broad range of black hole masses and luminosities. RM, as a tool that probes BLR gas structure, provides the means for such a test.

The RM method is successful for black hole mass determination because the emission lines in the BLR gas are found to respond to the stochastic continuum variations in an AGN via the transfer function (Peterson 1993)

$$L(v_z, t) = \int C(v_z, t - \tau) d\tau \quad (1)$$

where $L(v_z, t)$ is the emission-line luminosity at line-of-sight velocity v_z at observed time t , $C(t)$ is the con-

tinuum light curve, and (v_z) is the transfer function that maps continuum variability to the emission-line response at v_z after some time delay t . This simple model assumes a linear response and no background light; fitting AGN light curves requires the linearized echo model that subtracts off the reference levels, e.g. $L(v_z, t) = L_0(v_z)$ (see discussion in Horne et al. 2021). Monitoring both the continuum and the emission-line light curves provides data that can allow the determination of (v_z) , also known as the velocity-delay map, whose shape depends on the structure and kinematics of the BLR (Horne et al. 2004). In practice, precise velocity-resolved information in the form of the transfer function demands intense monitoring with frequent sampling, high signal-to-noise ratio (S/N) data, and long duration. As a result, the number of AGN with such data available to-date has been limited.

With the goal of refining black hole scaling relations through expanding the local reverberation-mapped AGN database by which they are anchored, we embarked on a campaign aimed to monitor AGN at higher luminosity than samples targeted in the 2008 and 2011 campaigns of the Lick AGN Monitoring Project (LAMP, e.g., Bentz et al. 2009b; Barth et al. 2015). Higher-luminosity AGN tend to have longer lags. Coupled with the fact that they have weaker variability amplitudes owing to the ample fuel supply in their accretion disks (vanden Berk et al. 2004; Wilhite et al. 2008; MacLeod et al. 2010), lag recovery is more challenging for these objects. The advent of large, multi-object programs such as SDSS-RM (Shen et al. 2015) and OzDES (Yuan et al. 2015) provides a new landscape where large numbers of reverberation lags for H β , Mg II, and C IV in quasars are determined over a broad redshift range (Shen et al. 2016; Grier et al. 2017, 2019). Grier et al. (2017) measured H β time lags for 44 AGN with luminosities $\log[L(5100 \text{ \AA})/L_{43-45.5}]$ at redshifts $z = 0.12-1$. Early results from the SDSS-RM campaign determined C IV lags in 52 quasars (Grier et al. 2019), whereas those from OzDES found two C IV-based black hole masses to be among the highest redshift ($z = 1.9-2.6$) and highest mass black holes [$M_{\text{BH}} = (3.3-4.4) \times 10^9 M_{\odot}$] measured thus far with RM studies (Hoormann et al. 2019). Complementing these large multi-fiber spectroscopic studies, our campaign targeted AGN at low redshifts and aimed to yield high-fidelity data for velocity-resolved lag measurements and dynamical modeling.

During the past several years, different groups have determined resolved velocity-delay reverberation signatures for dozens of unique objects among local Seyferts, changing-look AGN, and those with

H β asymmetry (e.g., Bentz et al. 2010a; Grier et al. 2013; Pancoast et al. 2014b; De Rosa et al. 2018; Du et al. 2018a; Williams et al. 2018, 2020; Lu et al. 2019; Zhang et al. 2019; Sergeev 2020; Lu et al. 2021; Bentz et al. 2021). Our primary intent is to use a large, relatively broad sample to investigate statistical luminosity-dependent trends in BLR structure and gas dynamics using velocity-resolved reverberation, which could directly impact the accuracy of virial mass estimates. Here we present our first results that focus on new H β velocity-resolved measurements along with integrated H β lags and virial black hole masses. Our dataset will allow for forward-modeling work using, for instance, the CAMEL code (Pancoast et al. 2011, 2014a, Villafañana et al., in prep.) to directly determine black hole masses.

This paper is organized as follows. The sample selection is described in Section 2. In Section 3, we present our observational program at Lick Observatory and details of the data reduction and processing work. Section 4 reports on our photometric campaign to provide the continuum light curves for our AGN sources. In Section 5, we illustrate our emission-line light curves and subsequent integrated and velocity-resolved H β lag detections, including an assessment of the lag significance and the observed variety of BLR kinematics. Section 6 depicts our line-width measurements and derived virial black hole masses, within the larger context of how our results compare with the existing AGN radius luminosity relation.

Throughout the paper, we have adopted $H_0 = 67.8 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.308$, and $\Omega_{\text{vac}} = 0.692$ (Planck Collaboration et al. 2016).

2. SAMPLE SELECTION

The past Lick AGN Monitoring Project (LAMP) campaigns targeted AGN with H β lags of $\geq 3-15$ days (Bentz et al. 2009b, 2010b; Barth et al. 2015). The chief objective of this campaign is to investigate BLR kinematics of moderate-luminosity AGN, so we targeted primarily Seyfert 1s with extinction-corrected $\log[L(5100 \text{ \AA})/L_{43-543.9}]$. Based on the radius luminosity relation (e.g., Bentz et al. 2013), the targeted AGN continuum luminosity range corresponds to an H β lag range of 20-30 days. We note, however, that recent results from the SEAMBH (Du et al. 2016a, 2018b; Du & Wang 2019) and SDSS-RM (Grier et al. 2017) programs have shown that some AGN have lags shorter than would be expected from earlier versions of the radius luminosity relation (Bentz et al. 2013).

We applied this 20-30-day criterion in H β lag or equivalently $L(5100 \text{ \AA})$ to various catalogs of Type 1

AGN (e.g., Boroson & Green 1992; Marziani et al. 2003; Peterson et al. 2004; Vestergaard & Peterson 2006; Bachev et al. 2008; Winter et al. 2010; Shen et al. 2011; Zu et al. 2011; Joshi et al. 2012; Bennert et al. 2015; Sun & Shen 2015). Our selection was further narrowed with a redshift limit at $z < 0.08$ to ensure that the H β and [O III] lines fall blueward of the cutoff wavelength of the dichroic used in the Kast spectrograph (5500 Å). Sources were selected with declination

$\delta > 5^\circ$ and magnitude < 17 in the optical V or r band. We also assessed the short-term variability in previous light curves from the Catalina Real-time Transient Survey (Drake et al. 2009) for all our candidate sources and selected those with at least 0.1 magnitude of variations. Targets having high-quality velocity-resolved lag measurement from previous RM campaigns were excluded (e.g., Denney et al. 2010; Grier et al. 2012; Du et al. 2014; Wang et al. 2014), as were those no longer featuring broad hydrogen recombination lines in archival Lick optical spectra.

In order to ensure that lags can be measured accurately, we placed a further constraint requiring that the monitoring duration of an AGN, defined as the longest continuous period when the AGN can be observed at airmass < 2 during our campaign, be at least 3 times larger than the expected H β lag based on simulations. Given that our initial campaign duration was designed to be 9 months long (which extended to 1 year later on), this constraint resulted in a sample of 29 objects with duration-to-lag ratio > 3 . Two sources with slightly shorter H β lag estimates (Mrk 315: 13 days; Mrk 704: 16 days) were included to better fill the right-ascension range of the sample.

After an initial probationary period of 1 month, we discarded 8 sources showing low continuum variability amplitude, retaining a sample of 21 objects for ongoing monitoring. This final sample of Seyfert 1s and their properties are listed in Table 1.

3. THE SPECTROSCOPIC CAMPAIGN AT LICK

3.1. Overview of the Observational Program

We conducted a spectroscopic monitoring program of 1 yr duration at Lick Observatory on Mount Hamilton, California. Prior to the start of the campaign, we ran simulations to determine the minimum threshold for successful lag recovery among monitoring periods of varying lengths and cadences. Our estimated H β lag range required a sampling cadence of two nights per week over the course of a year to sufficiently resolve time-dependent emission-line variations and to extend the temporal baseline for detecting robust reverberation sig-

nal across the entire extent of the BLR, accounting for anticipated weather losses and seasonal gaps.

This project was allocated 100 nights at the Lick 3 m Shane telescope across three observing semesters between 2016 April 28 and 2017 May 6 (UT). The observing runs were distributed mostly with a frequency of 2 nights per week as requested, with the exception of bright time and occasional scheduling constraints owing to other time-sensitive observing programs. Partial nights were occasionally exchanged with or obtained from other programs to facilitate the sampling cadence of certain targets during this monitoring period.

Spectroscopic data were taken using the Kast Double Spectrograph (Miller & Stone 1994). The red CCD detector was upgraded halfway into our campaign (September 17 20, 2016). The upgrade from the Reticon 400 1200 pixel CCD to the Hamamatsu 4096 1024 pixel CCD significantly improved the quantum efficiency, up to a factor of 2 at the red end, and removed severe fringing effects. Images taken with the new red CCD suffered from a high incidence rate of cosmic-ray hits and alpha-particle hits due to radioactive material in the new dewar window. These defects were removed at the reduction stage where frames were combined before the spectrum was extracted.

We employed the following setup for our AGN observations: the D55 dichroic with the 600/4310 grism (nominal coverage of 3300 5520 Å at 1.02 Å/pixel) on the blue side and the 600/7500 grating (nominal coverage of 4000 11,000 Å at 2.35 Å/pixel pre-upgrade, and 3800 10,000 Å at 1.31 Å/pixel post-upgrade) on the red side, a compromise between broad spectral coverage and moderate spectral resolution. The plate scale was 0.43/pixel for both the blue and the new red detectors, and 0.78/pixel for the old red detector. A slit width of 4 μ was employed to minimize slit losses due to seeing variations between different nights (Filippenko 1982). A fixed position angle, optimized for each individual target, was chosen so that the spectroscopic aperture sampled the same portion of the host galaxy across the duration of the monitoring campaign.

Exposure times were typically up to 30 min per object each night, split into 2 or 3 separate exposures (pre- and post-upgrade of the red CCD, respectively) to facilitate cosmic-ray cleaning. Standard calibrations including bias exposures, arc frames (using the Ar, He, Hg, Cd, and Ne lamps), and dome flats were taken during the afternoon, and well-calibrated flux standard stars (G191B2B, Feige 34, BD+284211, HZ44, BD+262606, and/or BD+174708) were observed during the nights. Before the red CCD upgrade, red-side observations suffered from severe fringing at wavelengths longward of

Table 1. Sample Properties

Object	Other Name(s)	Right Ascension (J2000)	Declination (J2000)	Redshift	Reference
Zw 535-012		00:36:20.983	+45:39:54.08	0.04764	1
I Zw 1	UGC 00545, Mrk 1502, PG 0050+124	00:53:34.940	+12:41:36.20	0.05890	2
Mrk 1048	NGC 985, VV 285	02:34:37.769	08:47:15.44	0.04314	2
Ark 120	UGC 03271, Mrk 1095	05:16:11.421	00:08:59.38	0.03271	3
Mrk 376	KUG 0710+457, IRAS 07105+4547	07:14:15.070	+45:41:55.78	0.05598	3
Mrk 9		07:36:56.979	+58:46:13.43	0.03987	1,3
Mrk 704	CGCG 091-065, MCG +03-24-043	09:18:26.005	+16:18:19.22	0.02923	4
MCG +04 22 042		09:23:43.003	+22:54:32.64	0.03235	5
Mrk 110	PG 0921+525	09:25:12.870	+52:17:10.52	0.03529	6,7
RBS 1303	CGS R14.01	13:41:12.904	14:38:40.58	0.04179	4
Mrk 684		14:31:04.783	+28:17:14.11	0.04608	1
Mrk 841	J15040+1026	15:04:01.201	+10:26:16.15	0.03642	5
Mrk 1392	1505+0342	15:05:56.553	+03:42:26.32	0.03614	8
SBS 1518+593		15:19:21.650	+59:08:23.70	0.07810	9
3C 382	CGCG 173-014	18:35:03.390	+32:41:46.80	0.05787	2
NPM1G+27.0587	2MASX J18530389+2750275	18:53:03.874	+27:50:27.72	0.06200	10
RXJ 2044.0+2833		20:44:04.500	+28:33:12.10	0.05000	10
PG 2209+184	II Zw 171	22:11:53.889	+18:41:49.86	0.07000	2
PG 2214+139	Mrk 304	22:17:12.262	+14:14:20.89	0.06576	2
RBS 1917	2MASX J22563642+0525167	22:56:36.500	+05:25:17.20	0.06600	2
Mrk 315		23:04:02.622	+22:37:27.53	0.03887	1

NOTE Objects in this and subsequent tables and figures are listed in RA order. Redshifts are from NED. References for L (5100 Å) used to estimate the H lag in our sample selection: (1) Previous Lick spectra; (2) Marziani et al. (2003); (3) Joshi et al. (2012); (4) Barth et al. (2015); (5) Winter et al. (2010); (6) Peterson et al. (2004); (7) Zu et al. (2011); (8) Bennert et al. (2015); (9) Sun & Shen (2015); (10) Bachev et al. (2008)

7000 Å. In order to remove the fringe pattern, dome flats at the position of every object were taken before or after each standard star and AGN observation on the red side. This time-consuming procedure was no longer needed after the upgrade that eliminated the fringing effect. The observing parameters for our sample are listed in Table 2.

The wet 2016–2017 winter season at Lick hampered our observing effort, particularly during December through February. Overall, our spectroscopic campaign achieved an observational success rate of $\lesssim 70\%$, where 30 nights were lost entirely to weather or dome issues. For another 12 nights, only 6 or fewer AGN were observed when typically 12–15 objects would have been observed on a night with good conditions. We were confident that at least fourteen nights of observations were carried out under photometric conditions. Of the 70 nights when data were obtained, half were done with the old red CCD and half with the upgraded detector.

The number of total epochs observed for each object ranges from 22 to 50, with a median of 38 (see Table 2).

3.2. Spectroscopic Data Reduction

The Kast spectroscopic data were reduced using a combination of standard routines in IRAF and IDL, following the procedure outlined by Barth et al. (2015). Here we provide a brief description of the reduction process of the blue-side data and pre-/post-upgrade red-side data.

In general, the data were processed via bias subtraction, flat fielding, cosmic-ray cleaning, one-dimensional (1D) extraction using an unweighted boxcar extraction region, wavelength calibration, and flux calibration. Error spectra were extracted and propagated through all subsequent calibration steps. Multiple spectra taken on the same night were combined. The width adopted for the spectral extraction was 10 \AA , though it was widened to $15 \text{--} 20 \text{ \AA}$ for some sources (Mrk 315, Mrk 9, MCG

Table 2. Observing Parameters

Object	Slit PA (deg)	t_{exp} (s)	Monitoring Period (UT)		N_{obs}	S/N	N_{phot}
Zw 535 012	100	1800	2016/06/27	2017/03/04	41	57	10
I Zw 1	45	1600	2016/07/03	2017/02/01	34	103	6
Mrk 1048	0	1800	2016/08/08	2017/02/16	27	88	5
Ark 120	0	1800	2016/08/15	2017/03/26	34	124	5
Mrk 376	70	1800	2016/05/01	2017/05/01	31	84	6
Mrk 9	90	1800	2016/05/01	2017/05/01	33	78	5
Mrk 704	130	1800	2016/10/20	2017/05/01	23	106	3
MCG +04 22 042	145	1800	2016/05/01	2017/05/01	34	54	7
Mrk 110	44	1200	2016/05/01	2017/05/01	41	74	6
RBS 1303	20	1200	2016/05/01	2017/05/01	22	67	5
Mrk 684	60	1800	2016/05/01	2017/05/01	43	98	12
Mrk 841	50	800	2016/05/01	2017/05/01	45	77	11
Mrk 1392	50	1800	2016/05/01	2017/05/01	39	55	10
SBS 1518+593	90	1800	2016/05/01	2017/05/01	48	44	8
3C 382	70	1800	2016/05/01	2016/12/03	50	81	12
NPM1G+27.0587	60	1800	2016/05/01	2016/12/03	38	55	7
RXJ 2044.0+2833	60	1800	2016/05/01	2016/12/31	46	58	9
PG 2209+184	60	1800	2016/05/01	2016/12/31	40	32	9
PG 2214+139	60	1800	2016/05/18	2016/12/31	43	81	10
RBS 1917	30	1800	2016/06/01	2016/12/31	32	39	9
Mrk 315	60	1800	2016/06/07	2016/12/31	35	59	9

NOTE t_{exp} represents the typical total on-source exposure time for each AGN every night it was observed, usually split among 2–4 frames. N_{obs} represents the total number of spectroscopic observations for each source. S/N represents the median S/N per pixel in the continuum at $(5100–5200) (1+z) \text{ \AA}$. N_{phot} represents the number of photometric nights for each source.

+04 22 042, and NPM1G+27.0587) and during particular nights where the seeing was exceptionally poor.

Most of the AGN were ux calibrated using the standard-star exposure that was closest in airmass. However, the ux -calibrated spectra for a few epochs exhibited some abnormal slopes and features close to the dichroic cutoff. We were unable to conclusively determine the cause of these anomalies, but they were most likely related to the dichroic. Since the [O III] 5007 line, situated close to the dichroic cutoff, was assumed to be constant throughout the observing campaign and was used to normalize the nightly spectra, this peculiarity identified in several spectra added extra scatter to the H β light curves. In these cases, we attempted to correct the problem by calibrating the AGN with standard stars that were taken close in time for six nights. As a result, this ux -calibration issue was partially rectified but remained responsible for some residual scatter in the emission-line light curves. For eight sources (Ark 120, RBS 1303, Mrk 9, RXJ 2044.0+2833, Zw 535 012,

SBS 1518+593, Mrk 684, and Mrk 110), we further divided the [O III] light curve by the median [O III] ux , and subsequently normalized the H β light curve with the residual [O III] light curve. This normalization helped to mitigate the residual scatter remaining in the H β light curves. We suspect that this ux -calibration anomaly due to the dichroic was occurring at a lower level throughout much of the campaign, which caused scatter larger than what might be typically achievable in other RM datasets.

For red-side data taken prior to 2016 Sep. 20, each observation was attenuated using a dome-flat exposure taken at the same telescope position. For red-side data taken between 2016 Sep. 20 and 2017 Jan. 31, the Kast red dewar was slightly tilted and thus the two-dimensional spectra had to be adjusted with a rotation of 0.9° before spectral extraction.

3.3. Photometric and Spectral Scaling

To ensure that all the spectra for each AGN are on consistent flux and wavelength scales across the monitoring period, a modified version of the procedure described by van Groningen & Wanders (1992, hereafter VGW92) was applied to the blue-side data for internal calibration (see Barth et al. 2015; Fausnaugh 2017, for more details). Having the spectra exhibit consistent spectral resolution throughout the temporal series is important particularly for extracting velocity-resolved measurements. First, we computed the shifts in wavelength among the time-series spectra from cross correlation. We applied these shifts to align the spectra and calculated an initial mean spectrum that was then designated as the reference. The [O III] $\lambda 5007$ fluxes, presumed to be intrinsically constant throughout the course of the observing campaign, were measured from spectra taken during photometric nights and were averaged to estimate the true absolute flux of [O III]. The number of photometric nights designated per object ranged from 3 to 12, with a median of 8 nights. The resulting reference spectrum was then scaled to match this measurement of the [O III] flux, with a median uncertainty of 8%.

Each night's spectrum was then aligned and scaled to the reference by matching the [O III] $\lambda 5007$ emission-line profile following our modified VGW92 scaling method. In order to minimize variations outside of intrinsic AGN variability, we aimed to achieve a uniform spectral resolution across the full time series of spectra. Where VGW92 ignores resolution corrections for epochs observed at lower resolution than the reference spectrum, we adopted instead a modified approach similar to that described by Fausnaugh (2017): we first applied a Gaussian broadening kernel to the reference spectrum to ensure that all the rescaled spectra match a single resolution, typically the worst among the spectra. Excluding the occasional extreme outliers, this broadening kernel had a typical σ of $1.5 \pm 3 \text{ \AA}$ with a median σ of 2 \AA for each AGN. Each of the spectra was then aligned by wavelength, scaled in flux, and broadened in spectral resolution to match the [O III] line profile of the broadened reference spectrum.

We performed a comparison of the modified VGW92 spectral scaling method to that using the Python code `mapspec` (Fausnaugh 2017) with the same wavelength windows and broadened reference spectrum. Differences between the two approaches include the following: (i) VGW92 calls for using a Gaussian kernel for smoothing, while `mapspec` offers an additional option of using Gauss-Hermite polynomials; and (ii) our modified VGW92 approach fits for free parameters via χ^2 minimization, while `mapspec` uses a Bayesian framework to optimize rescaling parameters and estimate model

uncertainties. The runtime for the modified VGW92 method was typically ~ 100 times shorter than for `mapspec`. We ran several tests comparing the level of scatter in the light curves of the [O III] line that resulted from both methods and concluded that modified VGW92 performed better for most of the AGN in our sample. We subsequently adopted the spectra scaled with the modified VGW92 approach for the ensuing analysis.

The scaling of the red-side data took a different approach. To normalize the red-side flux scale, red-side spectra were stitched to the corresponding blue-side spectra via aligning their overlapping regions (~ 5300 – 5500 \AA). All spectra were first aligned to the reference spectrum in wavelength. Because the ends of the spectra tended to be noisy, we used a weighted average to determine the overall multiplicative scaling factor. This technique worked well in most cases where the spectra were smooth or well behaved at the ends. The average scaled spectra for the sample are presented in Figure 1.

3.4. Noise-Corrected RMS Spectra

Relative variability across the spectrum can be visualized using the root-mean-square (rms) spectrum. The Peterson et al. (2004) procedure of taking the standard deviation of flux values at each wavelength element over all the epochs may inadvertently bias the rms spectrum given the inclusion of various noise factors in addition to genuine AGN variability (Barth et al. 2015). To remove the contribution due to photon-counting noise from the rms spectrum, Pei et al. (2017) suggested an approach to produce the excess rms (e-rms hereafter) spectrum defined per wavelength element and epoch i as

$$\text{e-rms} = \frac{1}{N-1} \sqrt{\sum_{i=1}^N [(F_i - \bar{F})^2 - \sigma_i^2]} \quad (2)$$

where N is the number of epochs in the time series, \bar{F} is the flux averaged over the time series, and F_i and σ_i refer to the flux and rms uncertainty in the flux at each band i , respectively. This was applied to the set of scaled spectra for each AGN.

In the case of an AGN with strongly varying broad-line profiles, broad features are expected to dominate the e-rms spectra in the relevant spectral regions. The narrow [O III] lines, on the other hand, generally vanish in the rms given that they are assumed to be constant within the time-series data. Presented in Figure 2, the e-rms spectra generally feature a strong blue continuum (except for Mrk 315), broad-line emission in the Balmer series and (usually) in He II. The absence of




Figure 1. Mean spectra for the LAMP2016 sample including both blue- and red-side data. Here, as in Table 1, the AGN are shown in order of increasing right ascensions.

residual narrow [O III] features confirms that our internal flux calibration worked well near this wavelength, but other residual narrow-line features are likely artifacts of imperfect flux calibration errors that increase at bluer wavelengths. This is particularly the case for Mrk 315, which has a red e-rms continuum slope and prominent [O II] $\lambda 3727$.

3.5. Spectral Decomposition

The traditional approach of measuring broad emission-line fluxes using a simple linear continuum subtraction is subject to several inadequacies. A linear continuum model is unable to separate blended emission-line features, such as He II or Fe II lines that often overlap the broad $H\beta$ profile. Furthermore, a linear fit is a poor approximation to the actual continuum, which includes both AGN and host-galaxy components. Residual errors from continuum subtraction can be particu-

larly problematic for velocity-resolved RM in the faint high-velocity wings of broad emission lines, where over-subtraction or under-subtraction of the continuum could lead to biased inferences on BLR structure and kinematics. In recent years, spectral decomposition approaches have been applied to the data from some RM campaigns (e.g., Barth et al. 2013, 2015; Hu et al. 2015). By fitting a multicomponent model to each night’s spectrum, the contributions of individual line and continuum components can be better isolated. We find that in most of our objects, the $H\beta$ line profiles isolated using the two methods do not differ dramatically except in the red wings, but spectral decomposition is useful for separating blended line components and removes the starlight component for objects with high starlight fraction much more effectively.

Here we have adopted and applied the spectral fitting method described by Barth et al. (2015) for the




Figure 2. The blue-side e-rms spectra for the VGW92-scaled time-series data featuring strong blue continuum and broad-line emission in the Balmer series, overlaid with dashed lines indicating the wavelengths of the prominent emission features (labeled in the top panels). The absence of residual narrow [O III] features demonstrates the effectiveness of our internal flux calibration at the red end, but other residual narrow-line features (such as the [O II] λ 3727 feature in Mrk 315) is most likely due to a combination of underlying continuum variability and imperfect spectrophotometric calibration errors at the blue end.

H β spectral region. We refer the reader to Barth et al. (2015) for a detailed description, and provide a brief overview of the method here. The model components include a power-law AGN continuum, starlight from a single-burst old stellar population at solar metallicity (Bruzual & Charlot 2003) convolved with a Gaussian velocity broadening, and emission lines including [O III] (narrow), H β (broad and narrow), He II (broad and narrow), He I (broad), and an Fe II emission template convolved with a Gaussian velocity broadening. Emission-line profiles were modeled using a fourth-order Gauss-Hermite function, except for the He I and He II lines for which a Gaussian was used. Each line’s velocity centroid was allowed to vary independently, except for the [O III] λ 4959, 5007 lines which were required to have the same velocity profile and a 1:3

flux ratio. For the Fe II blends, we tested different template spectra as described by Boroson & Green (1992), Véron-Cetty et al. (2004), and Kovačević et al. (2010). In the near future, we will employ a promising, new Fe II template spectrum based on Mrk 493 (Park et al., in prep.). Similar to Barth et al. (2015), the multicomponent Kovačević et al. (2010) template provided the best fit to the Fe II lines in all of our AGN and was used for the final fits, except for the case of I Zw 1. For I Zw 1 we found that the Boroson & Green (1992) template, which is based on I Zw 1 itself, provided the best fit with minimized residuals. As part of the fitting process, a Cardelli et al. (1989) reddening model is applied to the model spectrum, allowing $E(B - V)$ to be a free parameter. Fitting was carried out over a rest-wavelength range of approximately 4200–

5200 Å, with the exact range tailored to the data for each AGN depending on its redshift. The H γ + [O III] 4363 blend was masked out from the fit, because decomposing this blend would add several additional free parameters to the model. The full model, including the multicomponent Kovačević et al. (2010) iron template, includes 33 free parameters. Model fits were optimized using a Levenberg-Marquardt algorithm as implemented by Markwardt (2009). For each AGN, the model was first fitted to the mean spectrum, and the fit parameters from the mean spectrum fit were used as the starting parameter estimates for the fit to each individual night's spectrum. The overall mean spectrum as fitted with the different model components for each galaxy is illustrated in Figure 3.

4. THE PHOTOMETRIC CAMPAIGN

AGN continuum light curves were measured from imaging data. We chose the V band since it provides a fairly clean continuum measurement with relatively little contamination from broad emission lines for low-redshift sources. To maximize the chances of detecting the delayed response of the emission-line variations to those of the continuum, we began the photometric monitoring campaign two months before the start of the spectroscopic program. From February 2016 to May 2017, our team used a network of eight telescopes across the world and obtained high-precision V -band images with up to nightly cadence within each object's monitoring season.

4.1. Telescopes and Cameras

Our photometric campaign included observations with the following telescopes: the 0.76 m Katzman Automatic Imaging Telescope (KAIT) (Filippenko et al. 2001) and the Anna Nickel telescope at Lick Observatory on Mount Hamilton, California; the Las Cumbres Observatories Global Telescope (LCOGT) network (Brown et al. 2013; Boroson et al. 2014); the 2 m Liverpool Telescope at the Observatorio del Roque de Los Muchachos on the Canary island of La Palma, Spain (Steele et al. 2004); the 1 m Illinois Telescope at Mount Laguna Observatory (MLO; Smith & Nelson 1969) in the Laguna Mountains, California; the San Pedro Mártir Observatory (SPM) 1.5 m Johnson telescope (Butler et al. 2012; Watson et al. 2012) at the Observatorio Astronómico Nacional located in Baja California, México; the Fred Lawrence Whipple Observatory 1.2 m telescope on Mount Hopkins, Arizona; and the 0.9 m West Mountain Observatory (WMO) telescope at Utah Lake in Utah. KAIT, LCOGT, SPM-1.5 m, and

Liverpool are fully robotic telescopes. Table 3 summarizes the telescope and detector properties.

4.2. Photometry Measurements and Continuum Light Curves

All images were processed with reduction procedures applied as part of the standard pipeline for each facility, including overscan subtraction and flat-fielding. For telescopes that did not include world coordinate system (WCS) solutions as part of their standard processing, we used `astrometry.net` (Lang et al. 2010) to add WCS information to the FITS headers.

Measurement of the AGN light curves was carried out using the automated aperture photometry pipeline described by Pei et al. (2014), which can be used to measure AGN light curves using data from any number of telescopes with diverse camera properties. This procedure, written in IDL and based on the photometry routines in the IDL Astronomy User's Library (Landsman 1993), is designed to automate the process of identifying the AGN and a set of comparison stars in each image by their coordinates, measuring their instrumental magnitudes, and using the comparison-star measurements to obtain a consistent magnitude scale across the full time series for data from multiple telescopes.

The aperture-photometry routine returns photometric uncertainties (in magnitudes) based on the photon-counting statistics, background uncertainty, and CCD readout noise. However, other error sources contribute to the actual error budget, such as imperfect flat-fielding and point-spread function (PSF) variations across the field of view. To account for these additional error sources, we measured the excess variance $\frac{\sigma^2}{x}$ in the light curves of the comparison stars for each AGN, for each telescope's data. Assuming that the excess variance in the comparison-star light curves is a good estimate of the additional error over and above the statistical uncertainties, we inflated the fractional flux uncertainties on the AGN photometry by addition in quadrature using $\frac{\sigma_{\text{tot}}^2}{x} = \frac{\sigma_{\text{phot}}^2}{x} + \frac{\sigma^2}{x}$, where $\frac{\sigma_{\text{phot}}^2}{x}$ is the original photometric uncertainty on a given data point, and $\frac{\sigma^2}{x}$ is the average (over all comparison stars in the field) of the excess variance in the comparison-star light curves. This error adjustment was carried out separately for each telescope's data.

AGN light curves measured from different telescopes tend to have small offsets in magnitude relative to one another as a result of differences in filter transmission curves, even after normalizing them to the same set of comparison stars. To correct for these offsets, we applied an additive shift (in magnitudes) to bring each telescope's data into best average agreement with the




Figure 3. Spectral decomposition results. The observed mean spectrum (black) for each galaxy is plotted alongside the decomposed model components: starlight (purple), AGN power-law continuum (green), H β (magenta), He II (cyan), He I (orange), Fe II (grey), and [O III] (blue). The sum of the fits is represented by the red line. The He I component is negligible in most cases. The blend of H γ and [O III] λ 4363 is excluded from the fit. The Kovačević et al. (2010) Fe II template was applied to all sources except for I Zw 1, which was fitted with the Boroson & Green (1992) template.

Table 3. Telescope and CCD Properties for Imaging Observations

Telescope	D_{mirror} (m)	Detector	Field of View		Pixel Scale (\prime /pix)	Gain (e^- /ADU)	Read Noise (e^-)	Binning	
FLWO-1.2 m	1.2	Fairchild CCD 486	23 1	23 1	0.336	4.45	7.18	2	2
KAIT	0.76	Apogee AP7	6 8	6 8	0.80	4.5	12.0	1	1
LCOGT-2 m	2.0	Merope	4 7	4 7	0.467	1.0	9.0	2	2
LCOGT-1 m	1.0	Fairchild Imaging	26 5	26 5	0.389	1.0	13.5	1	1
Liverpool	2.0	e2V CCD 231	10	10	0.304	1.62	8.0	2	2
MLO-1 m	1.0	Fairchild 446	13 3	13 3	0.718	2.13	3.88	1	1
Nickel	1.0	Loral	6 3	6 3	0.368	1.7	8.3	2	2
SPM-1.5 m	1.5	Fairchild 3041	5 4	5 4	0.32	4.2	14.0	2	2
WMO-0.9 m	0.9	Finger Lakes PL-09000	25 2	25 2	0.61	1.37	12.0	1	1

data from the telescope having the most data points for each AGN. Finally, the V -band magnitude scale for each AGN field was calibrated using observations of standard stars taken on photometric nights at WMO and cross-checked against the AAVSO Photometric All-Sky Survey (Henden et al. 2016).

Magnitudes were converted to f for measurement of broad emission-line lags. The typical uncertainties of the photometric data points f are at the 0.4% level, not exceeding 0.7% for any AGN. The final V -band light curves are presented in Table 4 and shown alongside the emission-line light curves in 5.

Table 4. V -band Light Curve Data

Object	HJD	f	f	Telescope
IZw1	7555.8493	7.124	0.020	LCOGT-
IZw1	7557.8446	7.197	0.022	LCOGT-
IZw1	7558.8839	7.210	0.023	LCOGT-
IZw1	7559.8607	7.207	0.022	LCOGT-
IZw1	7561.9280	7.195	0.021	LCOGT-
IZw1	7563.9231	7.312	0.021	LCOGT-
IZw1	7564.9379	7.237	0.018	LCOGT-McDonald
IZw1	7566.9208	7.226	0.021	LCOGT-
IZw1	7567.9343	7.194	0.022	LCOGT-
IZw1	7570.9122	7.242	0.017	LCOGT-McDonald

NOTE Dates are listed as HJD - 2450000. Units for continuum flux density f are 10^{-15} erg s $^{-1}$ cm $^{-2}$ Å $^{-1}$. This table is published in its entirety in machine-readable format on the online version of the journal. Sample entries are shown here for guidance regarding the table's form and content.

5. EMISSION-LINE LIGHT CURVES AND LAG DETERMINATION

We used results from spectral decomposition to derive H emission-line light curves. The flexibility of the multicomponent modeling afforded us a number of ways to extract the H flux. In the spectral region near the H line, sources with strong Fe II emission were subjected to substantial degeneracy of spectral decomposition that could result in, from night to night, instability in model fitting in terms of how much flux gets assigned to the different model components. This instability results in additional noise in the H light curve. To minimize the noise due to degeneracy from spectral fits, we used a version of the decomposed model spectra where only the AGN and stellar continuum components have been removed. This version consisted of the residuals of all present emission lines after continuum sub-

traction. We then selected a corresponding extraction window free of other emission lines for the calculation of H flux (see Table 5). A different way to extract H flux is to use a version of the decomposed model spectra that only contained H broad and narrow components, with other emission-line components as well as the AGN and stellar continua removed. The resulting H light curves computed from the two methods were very similar for objects that have relatively clean H spectral regions, but those for the emission-line residual versions were less noisy for objects with more complex line profiles within the H spectral region. The final H light-curve measurements for the full sample are presented in Table 6. The light curves for the V band and H are shown in Figures 4–10. The rest of the emission-line features will be presented in a forthcoming paper.

Table 5. Emission-Line Extraction Windows in Å (observed frame), F_{var} , and [O III] Excess Scatter

Object	H		F_{var}	[O III]		[O III]
Zw 535-012	5018	5181	0.050	5220	5270	0.016
I Zw 1	5099	5199	0.044	5263	5337	0.054
Mrk 1048	5002	5153	0.071	5200	5252	0.004
Ark 120	4931	5102	0.195	5143	5200	0.043
Mrk 376	5069	5201	0.064	5259	5322	0.016
Mrk 9	5007	5127	0.069	5179	5241	0.016
Mrk 704	4945	5059	0.046	5126	5192	0.015
MCG +04-22-042	4986	5079	0.268	5149	5203	0.012
Mrk 110	4985	5120	0.083	5161	5208	0.015
RBS 1303	5001	5136	0.081	5185	5251	0.019
Mrk 684	5032	5131	0.058	5209	5262	0.048
Mrk 841	4949	5120	0.073	5167	5218	0.006
Mrk 1392	4958	5108	0.173	5165	5212	0.007
SBS 1518+593	5191	5310	0.051	5369	5428	0.024
3C 382	5030	5226	0.142	5268	5342	0.067
NPM1G+27.0587	5092	5246	0.051	5289	5352	0.024
RXJ 2044.0+2833	5040	5187	0.055	5240	5276	0.019
PG 2209+184	5147	5280	0.122	5329	5387	0.052
PG 2214+139	5110	5260	0.028	5307	5355	0.024
RBS 1917	5111	5261	0.028	5231	5444	0.016
Mrk 315	5002	5111	0.109	5174	5236	0.016

To account for the residual uncertainties from spectral scaling, we computed the excess variance in the [O III] light curve for each AGN as the additional uncertainty term. We then combined this excess variance term with the statistical error in quadrature such that

$$\sigma_f^2(\text{modified}) = \sigma_f^2(\text{stat}) + (f_{[\text{O III}]})^2 \quad (3)$$

Table 6. H Light Curve Data

Object	HJD	f	f (stat)	f (modified)
Mrk 110	7509.74	690.6	0.76	10.47
Mrk 110	7520.75	683.9	0.94	10.38
Mrk 110	7526.73	686.3	0.96	10.42
Mrk 110	7527.73	678.7	0.93	10.30
Mrk 110	7536.71	701.7	1.10	10.67
Mrk 110	7540.71	695.0	1.05	10.56
Mrk 110	7541.71	717.2	1.39	10.93
Mrk 110	7543.72	665.7	1.34	10.15
Mrk 110	7546.72	685.8	1.13	10.43
Mrk 110	7549.69	683.1	1.35	10.41

NOTE Dates are listed as HJD 2450000. Units for emission-line flux f are in 10^{15} erg s $^{-1}$ cm $^{-2}$. The original f represents the photon-counting errors on the flux measurement, while the modified f incorporates the spectral scaling uncertainties from [O III] scatter. This table is published in its entirety in machine-readable format on the online version of the journal. Sample entries are shown here for guidance regarding the table's form and content.

Both the statistical and the modified errors are listed in Table 6. Following Bentz et al. (2009a), we compute the variability statistics F_{var} for each light curve and found that it ranges from 0.028 to 0.268, with a median of 0.069 (see Table 5). Almost a third of our sample (3C 382, Ark 120, MCG +04 22 042, Mrk 1392, Mrk 315, and PG 2209+184) exhibit strong variability in the H light curves with $F_{\text{var}} > 0.1$. Among the rest of the sample, a few galaxies (such as Zw 535 012, Mrk 110, Mrk 841, RBS 1303, and RXJ2044.0+2833) still display discernible variability for plausible lag determination even if the signal is diluted by large error bars or outliers.

5.1. Integrated H Lags

Following White & Peterson (1994), Peterson et al. (2004) and others, cross-correlation was employed to determine the delay in the integrated H line signal relative to that from the V-band continuum from the photometric campaign. We chose to adopt the interpolated cross-correlation function (ICCF; e.g., Peterson et al. 1998) method. Given a lag range that represents reasonable bounds of the expected lag, the cross-correlation function (CCF) between the two light curves is com-

puted at every time lag at 0.5 day intervals via

$$F_{\text{CCF}}(\tau) = \frac{1}{N-1} \sum_{i=1}^N \frac{[L(t_i) - L][C(t_i - \tau) - C]}{L_{\text{std}} C_{\text{std}}} \quad (4)$$

where L (C) and L_{std} (C_{std}) are the mean and standard deviation of the emission-line (continuum) time series, respectively.

Two lag measures are computed: τ_{peak} , the lag at the peak of the CCF, and τ_{cen} , the centroid of the CCF for all points in the CCF above a predetermined threshold value (80% of the peak CCF value). We adopt the latter for the subsequent analysis. For error analysis, we employed the Monte Carlo flux randomization method (Peterson et al. 1998) and repeated the CCF calculation for 10^3 realizations, building up distributions of correlation measurements. The median and 1- σ widths of the cross-correlation centroid distributions (CCCD) were then adopted as the final lag measurements, and their associated uncertainties as shown in Figures 4-10.

5.2. Assessment of Cross-Correlation Reliability

In any RM campaign, it is expected that some AGN will yield highly robust measurements of reverberation lag, thanks to strong variability and high-quality data, while other objects may exhibit little or no evidence for correlated variability between the continuum and emission-line light curves. This can occur as a result of observational factors including low S/N or poor temporal sampling, or factors intrinsic to the AGN including low variability amplitude or low responsivity of the emission line to variations in the ionizing continuum. There is not necessarily a clear demarcation between reliable and unreliable lag measurements, and a variety of methods have been used to assess the significance or quality of lag detections (e.g., Grier et al. 2017). In some cases, a simple threshold value of the correlation strength r_{max} is used, but r_{max} alone is not necessarily a good indicator of correlation significance for red-noise light curves. An alternative method is to employ null-hypothesis testing to determine the probability that two uncorrelated red-noise light curves having the same S/N and cadence as the data would yield an r_{max} at least as strong as the observed value. Such methods have been employed for analysis of multiwavelength continuum correlations in AGN (e.g., Uttley et al. 2003; Arévalo et al. 2008; Chatterjee et al. 2008), but until recently have rarely been employed to assess broad-line reverberation lags (Penton et al. 2021; Li et al. 2021). Our method will be presented in detail by Guo et al. (in prep.), and we provide a brief description here.




Figure 4. Light curves for the LAMP2016 sample (Zw 535–012, I Zw 1, Mrk 1048): V-band continuum flux density f_λ in 10^{-15} erg cm $^{-2}$ s $^{-1}$ Å $^{-1}$, color-coded by telescopes (top left); the H β emission line flux f_{line} in 10^{-15} erg cm $^{-2}$ s $^{-1}$ (bottom left). The error bars plotted here for the H β light curves incorporate the uncertainty term from the normalized excess scatter of [O III]. The cross-correlation function is shown in the right panel for each AGN, alongside the cross-correlation centroid distribution in yellow. The dotted vertical line indicates the median value of the CCF.




Figure 5. Same as Figure 4 (but for Ark 120, Mrk 376, Mrk 9)




Figure 6. Same as Figure 4 (but for Mrk 704, MCG +04-22-042, Mrk 110)




Figure 7. Same as Figure 4 (but for RBS 1303, Mrk 684, Mrk 841)




Figure 8. Same as Figure 4 (but for Mrk 1392, SBS 1518+593, 3C 382)




Figure 9. Same as Figure 4 (but for NPM1G+27.0587, RXJ 2044.0+2833, PG 2209+184)




Figure 10. Same as Figure 4 (but for PG 2214+139, RBS 1917, Mrk 315)

We first generated light curves according to the damped random walk (DRW) model with the Python software `CARMA`¹ (Kelly et al. 2009, 2014). The DRW model provides an adequate description of ultraviolet/optical variabilities in AGN, though with plausible deviations on short (McHardy et al. 2006; Mushotzky et al. 2011; Kasliwal et al. 2015; Smith et al. 2018) or long (MacLeod et al. 2010; Guo et al. 2017) timescales. Each simulated light curve represents a segment randomly selected from a 100-times longer light curve predicted from the same DRW model fitted to the observed light curve. We resampled each mock light curve to have the same cadence as the real observations and added Gaussian random noise based on the S/N of the data, creating 10^3 realizations of each light curve. Cross-correlation measurements were then carried out using the simulated data to determine the distribution of r_{\max} values that would be obtained for uncorrelated light curves. A two-way simulation was performed – that is, we calculated the CCF between the real continuum and each simulated emission-line light curve for the first 500 simulations, and then the opposite way for the rest of the realizations.

We measured the CCF for all the simulations, using the same lag search range employed for each AGN, and counted, out of 10^3 , the number of positive lags ($\tau > 0$) with peak values r_{\max} higher than our observed r_{\max} . The resulting fraction represents the probability that a correlation signal found between two uncorrelated light curves would exceed the correlation signal of the data. This derived p -value, denoted $p(r_{\max})$, thus provides an indicator of the robustness of our lag detections given the observed r_{\max} and the observed properties of the light curves including S/N, cadence, and duration.

We emphasize that the p -values derived from this method are not false-positive probabilities, and they do not give a measure of the absolute significance of a lag detection. Strictly speaking, our p -values simply give the probability that uncorrelated light curves having the same statistical properties as the data would give a cross-correlation signal as strong as that seen in the data. Consequently, a smaller p -value denotes a more robust and reliable detection of the correlation signal between two light curves. In general, for broad-line RM we have a strong prior that a reverberation signal is very likely to be present in the data, thus our null hypothesis of intrinsically uncorrelated light curves is an extreme and fairly unlikely scenario. We further emphasize that there is no strict cutoff between significant and insignif-

icant lag detections by this method; the p -value merely gives an indication of the relative degree of reliability between different measurements.

We assess the quality of our resulting lags based on these quantitative assessment indicators in Figure 11. As expected, there is an anticorrelation between r_{\max} and $p(r_{\max})$, but it is not a tight one-to-one relationship owing to the differences in S/N, sampling cadence, and variability amplitude among our sample. The results are consistent with the general expectation that the highest-quality light curves that exhibit clear correlated variability have high r_{\max} and small $p(r_{\max})$ values. Considering these lag-assessment results, we conclude that 16 of the 21 AGN in our sample falling in the region $p(r_{\max}) < 0.2$ and $r_{\max} > 0.6$ have correlations between continuum and H β light curves that are sufficiently robust for the ensuing analyses (noting that they display a broad range of p -values and thus range from highly robust to relatively weak detections of correlated variability), while we discard the remaining five objects from further analysis based on their low r_{\max} and high p -values. Based partly on visual inspection of the light-curve quality, our chosen r_{\max} threshold is relatively conservative compared to those selected by other large surveys (e.g., minimum $r_{\max} = 0.45$ by SDSS-RM; Grier et al. 2017). These r_{\max} and $p(r_{\max})$ values, along with AGN luminosities and lag measurements in both observed and rest frames, are reported in Table 7. [Only the r_{\max} and $p(r_{\max})$ are reported for the subsample of sources with unreliable lag measurements.]




Figure 11. $p(r_{\max})$ versus r_{\max} as two objective lag-assessment parameters for our sample. Sources with $p(r_{\max})$ above 0.2 or r_{\max} less than 0.6 are deemed to have unreliable lags as represented by open circles.

¹ https://github.com/brandonckelly/carma_pack

Table 7. AGN Luminosity and H β Cross-Correlation Lag Results

Object	L (5100 Å) (10^{43} erg s $^{-1}$)		Observed Frame		Rest Frame		r_{\max}	$p(r_{\max})$	BLR Kinematics
			cen	peak	cen	peak			
			(days)	(days)	(days)	(days)			
Zw 535 012	4.9	0.7	21.3 $^{+8.5}_{-4.8}$	22.0 $^{+6.5}_{-8.0}$	20.3 $^{+8.1}_{-4.6}$	21.0 $^{+6.2}_{-7.6}$	0.63	0.12	Infalling
Mrk 1048	9.5	1.8	7.8 $^{+10.1}_{-9.8}$	10.0 $^{+9.0}_{-13.0}$	7.4 $^{+9.7}_{-9.4}$	9.6 $^{+8.6}_{-12.5}$	0.62	0.11	Infalling
Ark 120	9.2	3.4	19.3 $^{+6.1}_{-4.6}$	20.5 $^{+5.5}_{-4.5}$	18.7 $^{+5.9}_{-4.5}$	19.9 $^{+5.3}_{-4.4}$	0.95	0.03	Ambiguous
Mrk 9	5.6	0.4	20.3 $^{+4.2}_{-7.6}$	15.0 $^{+12.0}_{-7.0}$	19.5 $^{+4.1}_{-7.3}$	14.4 $^{+11.5}_{-6.7}$	0.79	0.10	Ambiguous
Mrk 704	6.2	0.6	29.8 $^{+10.4}_{-10.3}$	34.5 $^{+5.5}_{-19.5}$	28.9 $^{+10.2}_{-10.0}$	33.5 $^{+5.3}_{-18.9}$	0.82	0.19	Infalling
MCG +04 22 042	1.6	0.4	13.7 $^{+2.5}_{-1.9}$	10.5 $^{+3.0}_{-1.0}$	13.3 $^{+2.4}_{-1.8}$	10.2 $^{+2.9}_{-1.0}$	0.95	0.00	Symmetric
Mrk 110	7.2	1.7	28.8 $^{+4.4}_{-5.2}$	22.5 $^{+7.5}_{-4.0}$	27.8 $^{+4.3}_{-5.1}$	21.7 $^{+7.2}_{-3.9}$	0.92	0.02	Symmetric
RBS 1303	2.4	0.4	19.4 $^{+3.6}_{-4.5}$	13.5 $^{+7.5}_{-4.5}$	18.7 $^{+3.4}_{-4.3}$	13.0 $^{+7.2}_{-4.3}$	0.90	0.02	Out owing
Mrk 841	6.7	0.9	11.7 $^{+5.0}_{-3.7}$	9.0 $^{+5.0}_{-2.0}$	11.2 $^{+4.8}_{-3.5}$	8.7 $^{+4.8}_{-1.9}$	0.74	0.02	Infalling
Mrk 1392	1.6	0.6	27.6 $^{+3.6}_{-4.0}$	17.5 $^{+3.0}_{-2.0}$	26.7 $^{+3.5}_{-3.9}$	16.9 $^{+2.9}_{-1.9}$	0.95	0.01	Symmetric
SBS 1518+593	10.6	1.0	21.7 $^{+9.0}_{-9.5}$	22.0 $^{+9.5}_{-11.0}$	20.1 $^{+8.4}_{-8.9}$	20.4 $^{+8.8}_{-10.2}$	0.60	0.03	Infalling
3C 382	15.0	3.1	10.0 $^{+7.2}_{-7.1}$	6.5 $^{+13.5}_{-7.0}$	9.5 $^{+6.8}_{-6.7}$	6.1 $^{+12.8}_{-6.6}$	0.86	0.12	Symmetric
NPM1G +27.0587	10.9	1.0	8.5 $^{+5.0}_{-4.7}$	7.5 $^{+6.5}_{-4.0}$	8.0 $^{+4.7}_{-4.5}$	7.1 $^{+6.1}_{-3.8}$	0.74	0.16	Infalling
RXJ 2044.0+2833	5.3	0.5	15.1 $^{+1.7}_{-2.0}$	15.0 $^{+4.0}_{-3.5}$	14.4 $^{+1.6}_{-1.9}$	14.3 $^{+3.8}_{-3.3}$	0.78	0.04	Infalling
PG 2209+184	2.3	0.6	14.6 $^{+3.0}_{-3.1}$	16.0 $^{+3.0}_{-5.0}$	13.7 $^{+2.9}_{-2.9}$	15.0 $^{+2.8}_{-4.7}$	0.89	0.01	Ambiguous
RBS 1917	4.9	0.5	12.7 $^{+4.6}_{-4.1}$	12.5 $^{+4.5}_{-3.5}$	11.9 $^{+4.3}_{-3.9}$	11.7 $^{+4.2}_{-3.3}$	0.76	0.02	Out owing
I Zw 1	28.5	4.0					0.18	0.89	
Mrk 376	17.6	1.8					0.26	0.77	
Mrk 684	8.3	0.9					0.46	0.12	
PG 2214+139	21.2	1.3					0.62	0.45	
Mrk 315	3.2	0.4					0.53	0.23	

5.3. Velocity-Resolved H β Lags

To obtain velocity-resolved reverberation results as a means to probe the kinematics of the BLR gas, we used CCF to compute lag measurements for individual velocity segments of the emission line. Our procedure follows that described by Bentz et al. (2009b), Denney et al. (2009), and Grier et al. (2013), and is detailed below.

To verify the effect of binning on the apparent BLR kinematics, we divided the broad H β component into eight bins via two different schemes: (i) bins of equal rms flux, and (ii) bins of equal velocity width. In the first approach, we initially determined the total rms flux from integrating the H β line in the e-rms spectrum. Then we established velocity bins such that the rms flux within each bin would equal one-eighth of the total flux. There are some exceptions in the cases where the red wing of the rms profile suffered from significant noise and the last bin was thus truncated at the red line wing. In the second approach, bins are divided evenly across the width of the line in velocity space. In either binning scheme, light curves were computed from each bin of

the continuum-subtracted H β profile and lags were measured against the V-band continuum light curve. Resulting lags from the two binning schemes generally agree in the line core where the S/N is high; minor disagreement is found in the noisy line wings of some sources. For clarity, we adopt the first scheme (equal-rms-flux binning) and show the resulting velocity-resolved lag spectra for each AGN with robust lag in Figure 12 and in Table 8.

5.4. Implications for BLR Kinematics

The velocity-resolved lags allow us to make simple qualitative inferences about the BLR geometry and kinematics from the transfer-function models (e.g., Welsh & Horne 1991; Horne et al. 2004; Bentz et al. 2009b). Qualitatively, symmetric velocity-resolved structure around zero velocity is consistent with either Keplerian, disk-like rotation or random motion without net radial inflow or outflow over an extended BLR. In such a case, the high-velocity wings exhibit the shortest lags because the highest rotation speeds correspond to material orbiting closest to the central black




Figure 12. Velocity-resolved reverberation lags for our sample. In each of the top panels, the H β lags in days (y-axis) shown are for bins of equal rms flux, plotted against relative velocity in km s^{-1} (x-axis). The blue dotted line indicates the overall integrated H β lag with $\pm 1\sigma$ uncertainty spanned by the blue bar. The middle and bottom panels show the H β profile of the mean and rms spectra, respectively.




Figure 12. Continued

Table 8. Velocity-Resolved Lags

Object	Velocity Bin		τ_{cen} (days)
	(km s ⁻¹)		
Mrk110	-1910	2324	30.34 ^{+3.90} _{3.86}
Mrk110	-539	417	31.66 ^{+2.60} _{2.79}
Mrk110	-151	357	34.66 ^{+2.57} _{2.26}
Mrk110	205	357	35.83 ^{+3.04} _{2.85}
Mrk110	563	357	34.65 ^{+2.79} _{3.48}
Mrk110	1040	596	29.80 ^{+4.45} _{4.04}
Mrk110	1874	1073	14.12 ^{+8.08} _{5.93}

NOTE This table is published in its entirety in machine-readable format on the online version of the journal. Sample entries are shown here for guidance regarding the table's form and content.

hole. Radially infalling gas produces an asymmetric pattern displaying longer lags at the high-velocity blue wing, while the opposite is true in the case of outflow-dominated kinematics. We note that the presence of absorption, noise, or other geometric complexities may muddle these interpretations, which could be elucidated with direct modeling using forward-modeling codes such as CAMEL in follow-up work (Villaña et al., in prep.).

We find a diversity of velocity-resolved lag spectra among our sample representing the qualitative signature of symmetric (3C 382, MCG +04 22 042, Mrk 110, Mrk 1392), infalling (Mrk 1048, Mrk 704, Zw 535 012, Mrk 841, RXJ 2044.0+2833, NPM1G+27.0587, SBS 1518+593), and outflowing (RBS 1303 and RBS 1917) scenarios. In the cases of Ark 120, Mrk 9, and PG 2209+184, the velocity-resolved structure appears flat across the H β profile within the uncertainties, which may be difficult to describe with simple models. Whether these kinematics exhibit luminosity-dependent trends and how a subset of these results relate to other RM samples will be further examined in §6 and the Appendix, respectively.

6. BLACK HOLE SCALING RELATIONS

6.1. AGN Radius Luminosity Relationship

The applicability of the radius luminosity relationship to single-epoch mass determination (Shen et al. 2008; Shen & Liu 2012) relies on the small scatter initially quantified (~ 0.2 dex; Kaspi et al. 2005; Bentz et al. 2009a, 2013; Kilerci Eser et al. 2015; Martínez-Aldama et al. 2020), but recent studies of in-

creasingly diverse AGN samples led to an increase in this scatter (e.g., Grier et al. 2017; Du et al. 2018b). The main reason for this increased scatter seems to depend on the Eddington ratio (Bian et al. 2012; Du et al. 2015, 2016a; Martínez-Aldama et al. 2019; Dalla Bontà et al. 2020), an important third parameter suggesting that there is more diversity in the AGN population than can be explained by the simple two-parameter radius luminosity relationship. Here we compare our sample on the AGN radius luminosity relationship with other RM samples from the literature (Bentz et al. 2013; Grier et al. 2017; Du et al. 2018b) in Figure 13. The best-fit line from SDSS-RM for this relation is

$$\log(R_{\text{BLR}}/\text{ltday}) = K + \log(L(5100)/10^{44} \text{ erg s}^{-1}) \quad (5)$$

with a median slope $\alpha = 0.45$ and a median normalization $K = 1.46$ (Fonseca Alvarez et al. 2019). We computed the AGN luminosity $L(5100 \text{ \AA})$ based on the AGN continuum flux at 5100 \AA as extracted from the decomposition of the mean spectrum. Luminosity distances were derived using the cosmology calculator ($H_0 = 67.8 \text{ km s}^{-1} \text{ Mpc}^{-1}$) provided by Wright (2006). All of our AGN fall within 0.5 dex of the radius luminosity relation as defined by the literature points within the range of $L(5100 \text{ \AA}) = 10^{43.5-44.4} \text{ erg s}^{-1}$. The best-fit line in the form of Equation 5 for all the data points combining our current work with those from the literature has a normalization $K = 1.35$ and $\alpha = 0.38$.

We further explore whether BLR kinematics might depend on AGN luminosity. Past work has shown that while inflow or outflow are indicated in a subset of AGN, the majority of H β velocity-delay maps appear roughly symmetric with rotationally dominated kinematics (e.g., Bentz et al. 2009b, 2010b; Denney et al. 2010; Barth et al. 2011; Horne et al. 2021). The AGN with velocity-resolved reverberation detected to-date are mostly within the low-luminosity range [$L(5100 \text{ \AA}) = 10^{42.43-5} \text{ erg s}^{-1}$]. Our results add significantly to the existing number of sources having velocity-resolved measurements with $L(5100 \text{ \AA}) > 10^{44} \text{ erg s}^{-1}$. While we caution that velocity-resolved lag spectra provide only a qualitative impression of the BLR kinematics, we illustrate the BLR kinematics via different colored symbols shown alongside those from the literature in Figure 14. No trend is seen between BLR kinematics and AGN luminosity either within our LAMP2016 sample or when combined with past results from the literature. It is plausible that the BLR shows time-dependent structural changes, from both the kinematic and the geometric points of view, and there may be a lag in its correlation with the luminosity of the AGN. The diversity in BLR kinematics found in our mod-




Figure 13. Radius–luminosity relationship of our sample (blue, filled) superimposed on top of other work from the literature (open symbols; Bentz et al. 2013; Grier et al. 2017; Du et al. 2018b). The best-fit relation from SDSS with a median slope α of 0.45 and a median normalization K of 1.46 (Fonseca Alvarez et al. 2019) is plotted as the dashed line. The best-fit line in the form of Equation 5 for all the data points combining our current work with those from the literature (blue) has a normalization $K = 1.35$ and $\alpha = 0.38$.

erately high-luminosity sample appears consistent with that seen among AGN with super-Eddington accretion rates (e.g., Du et al. 2016b).

6.2. H β Line Widths

Following the procedure outlined by Barth et al. (2015), we measured the line width of the broad H β component from both the spectrally-decomposed mean spectrum and the rms spectrum. The mean spectrum constructed from the time-series data has a continuum level of zero with just residual noise from spectral fitting, but the continuum of the rms spectrum incorporates photon-counting errors in addition to residual noise. The noise continuum is removed after a simple linear fit before making line-width measurements.

We computed the line width using two separate parameters according to Peterson et al. (2004): the full width at half-maximum intensity (FWHM) of the line, and the line dispersion σ_{line} . The FWHM of a single-peaked line is defined as the difference between the wavelengths blueward and redward of the peak $P(\lambda)_{\text{max}}$ that correspond to half of its height. A double-peaked line requires additional attention to defining $P(\lambda_{\text{blue,red}})_{\text{max}}$ for the blue and red peaks, respectively, and thus their corresponding wavelengths at half the maximum height, but otherwise the procedure in determining the differ-




Figure 14. Radius–luminosity relationship of our sample (filled and color-coded by BLR kinematics) along with other velocity-resolved results from Bentz et al. (2010a), Pancoast et al. (2014b), Grier et al. (2013), De Rosa et al. (2018), and Du et al. (2018b) (open symbols with the same color coding). No trend is seen for the different BLR kinematics within our LAMP2016 sample or when combined with the literature.

ence between the resulting wavelengths is similar. As for σ_{line} (square root of the second moment of the H β line profile), we adopted from Peterson et al. (2004) the following:

$$\sigma_{\text{line}}^2(\lambda) = \langle \lambda^2 \rangle - \lambda_0^2, \quad (6)$$

where

$$\langle \lambda^2 \rangle = \frac{\int \lambda^2 P(\lambda) d\lambda}{\int P(\lambda) d\lambda} \quad (7)$$

and

$$\lambda_0 = \frac{\int \lambda P(\lambda) d\lambda}{\int P(\lambda) d\lambda}. \quad (8)$$

In each of these calculations, we also varied the endpoints of the H β extraction window randomly within $\pm 5 \text{ \AA}$ ten times in order to account for uncertainties brought about by the precise choice of the H β spectral limits. The resulting line-width measurement is the mean from these ten trials.

For error analysis associated with the measured width parameters, we undertook a Monte Carlo bootstrapping procedure and simulated 100 realizations of the mean and rms spectra, respectively, based on random subset sampling of the existing nightly spectra. For a source with n epochs of observations, we simulate new mean

and rms spectra of the broad H β component from combining n randomly-drawn time-series spectra with repetition allowed. We repeated our measurement of the FWHM and σ_{line} on these new spectra, where the means and the standard deviations of the resulting distributions were taken as the line widths and their corresponding uncertainties, respectively.

The final line-width measurements are corrected for instrumental broadening subtracted in quadrature from the observed line width σ_{obs} :

$$\sigma_{\text{intrinsic}}^2 = \sigma_{\text{obs}}^2 - \sigma_{\text{inst}}^2 \quad (9)$$

Given that we used the same blue-side spectral setup as the LAMP2011 campaign, we adopt from Barth et al. (2015) the instrumental FWHM of 380 km s^{-1} and dispersion $\sigma_{\text{inst}} = 162 \text{ km s}^{-1}$ for our small corrections. The resulting FWHM and σ_{line} measured from both the mean and the e-rms spectra are presented in Table 9.

6.3. Virial Black Hole Masses

The mass of a black hole may be determined most robustly using dynamical modeling codes such as CARMEL (e.g., Pancoast et al. 2011). Nonetheless, virial estimates are useful within uncertainties that depend on the BLR geometry. The black hole mass and its associated uncertainty are computed and propagated from the virial equation

$$M_{\text{virial}} = f \frac{c}{G} v^2 \quad (10)$$

$$M = \frac{\frac{M_{\text{cen}}^2}{\sigma_{\text{line}}^2} + \frac{M_{\text{cen}}^2}{v_{\text{line}}^2}}{\frac{1}{\sigma_{\text{line}}^2} + 4 \frac{1}{v_{\text{line}}^2}} \quad (11)$$

where f is the scaling factor that depends on the geometry and kinematics of the BLR, c is the speed of light,

τ_{cen} is the time delay with uncertainties, v_{line} is the velocity of the BLR gas with uncertainties as measured by the line width, and G is the gravitational constant. The empirically fitted virial factor from Woo et al. (2015) is consistent with that from CARMEL dynamical modeling (Pancoast et al. 2014b), so we have adopted for our black hole mass calculations $f = 10^{0.65} = 4.47$ from Woo et al. (2015), $M_{\text{cen}} = M_{\text{cen}}$, and $v_{\text{line}} = \sigma_{\text{line}}(\text{rms})$ from Dalla Bontà et al. (2020) to facilitate comparisons with other studies. We present both the virial products $c_{\text{cen}}^2 \tau_{\text{cen}}^2 G$ and the derived f -dependent black hole masses in Table 10. Comparisons of our results to existing prior black hole mass measurements for several

sources can be found in the Appendix. Overall, we find agreements between our measurements and those from the literature within uncertainties.

7. CONCLUSIONS

We present the first results from our 100-night LAMP2016 campaign where we monitored a sample of 21 luminous Seyfert 1 nuclei with AGN luminosity $L_{5100\text{\AA}} \approx 10^{44} \text{ erg s}^{-1}$ during April 2016–May 2017 at Lick Observatory. Our analysis here provided the H β emission-line light curves and integrated H β lag detections measured against V -band continuum photometric light curves for the full sample. We further assessed the significance of the lag determinations, and computed, for a subset of sources with good-quality lags, their velocity-resolved reverberations, inferred BLR kinematics, broad H β line widths, and virial black hole mass estimates.

The unusually rainy weather at Mount Hamilton during the winter months of 2016–2017 hampered our ability to monitor the variability of our AGN as closely as would have been ideal, leaving large gaps in several of our emission-line light curves. Consequently, we were not able to measure reliable lags for six of our AGN owing to inadequate light curves. Our overall results nearly double the number of existing velocity-resolved lag measurements, particularly at the moderately high-luminosity regime among previous reverberation-mapped samples, and revealed a diversity of signatures potentially indicative of different BLR gas kinematics. Given the complexity of the multiple factors involved, the lack of any clear correlation between AGN luminosity and velocity-resolved lag structure suggests that luminosity itself is not the sole factor responsible for setting the kinematic state of the BLR. It may also depend on other properties of the accreting black holes. Follow-up direct dynamical modeling work will shed light on the detailed kinematics and provide robust dynamical black hole masses to help further calibrate widely-adopted black hole scaling relations.

ACKNOWLEDGMENTS

We thank the anonymous referee for thoughtful comments and great suggestions that significantly improved this manuscript. We are grateful to M. Fausnaugh for offering help with their `mapspec` code used to test different spectral scaling of our Shane/Kast data, and M. Bentz for providing her table of values for the radius–luminosity relationship. We acknowledge the following individuals for their time and effort contributed to

Table 9. Rest-Frame Line Widths of the Broad H α Component

Object	Mean				rms			
	FWHM		line		FWHM		line	
Zw 535 012	2705	33	1474	28	2005	80	1259	112
I Zw 1	2187	920	1644	102	2665	700	916	165
Mrk 1048	4830	80	1840	58	4042	406	1726	76
Ark 120	5656	44	2125	41	4765	84	1882	42
Mrk 376	10055	341	2662	38	14438	1439	3633	210
Mrk 9	3751	71	1555	68	3159	144	1326	69
Mrk 704	8597	448	2002	63	8597	448	2009	102
MCG +04 22 042	2658	57	1141	39	2120	39	977	29
Mrk 110	2048	13	1203	32	2052	100	1314	69
RBS 1303	2286	21	1243	26	1738	113	1292	156
Mrk 684	2174	33	1216	74	2391	632	2010	773
Mrk 841	7073	311	2139	55	7452	660	2278	96
Mrk 1392	4267	25	1635	13	3690	138	1501	38
SBS 1518+593	3312	24	1782	21	4656	1692	2249	255
3C 382	7772	54	4218	67	13926	2337	5616	97
NPM1G +27.0587	3501	28	1683	42	2893	177	1735	136
RXJ 2044.0+2833	2196	31	989	32	2047	72	870	50
PG 2209+184	4045	34	1573	40	3247	88	1353	64
PG 2214+139	5133	17	1947	55	4867	473	1984	90
RBS 1917	2399	11	1180	50	1653	287	851	154
Mrk 315	3097	252	1824	67	3411	1438	2031	260

NOTE All line-width measurements are in km s⁻¹.

the Lick observing campaign: (Shane) Zachary Parsons, Estefania Padilla Gonzalez, Noah Rivera, Cristilyn Gardner, Jake Haslemann, Sean Lewis, and Ellen Glad; (Nickel) Nick Choksi, Sameen Yunus, Je Molloy, Andrew Rikhter, and Haynes Stephens. Photometric data collection at MLO was supported by NSF grant AST-1210311; we thank Robert Quimby, Emma Lee, Joseph Tinglof, Eric McLaughlin, Amy Igarashi, and Tariq Johnson for assistance with these observations.

We are deeply grateful to the UCO/Lick staff for help with scheduling and supporting the observations. Research at Lick Observatory is partially supported by a generous gift from Google. The Kast red CCD detector upgrade, led by B. Holden, was made possible by the Heising-Simons Foundation, William and Marina Kast, and the University of California Observatories. KAIT and its ongoing operation were made possible by donations from Sun Microsystems, Inc., the Hewlett-Packard Company, AutoScope Corporation, Lick Observatory, the NSF, the University of California, the Sylvia & Jim Katzman Foundation, and the TABASGO Foundation. Data presented herein were obtained using the UCI Re-

mote Observing Facility, made possible by a generous gift from John and Ruth Ann Evans.

Research at UC Irvine has been supported by NSF grants AST-1412693 and AST-1907208. V.U. acknowledges funding support from the University of California Riverside's Chancellor's Postdoctoral Fellowship and NASA Astrophysics Data Analysis Program Grant #80NSSC20K0450. Her work was conducted in part at the Aspen Center for Physics, which is supported by NSF grant PHY-1607611; she thanks the Center for its hospitality during the "Astrophysics of Massive Black Holes Merger" workshop in June and July 2018. T.T. acknowledges support by the Packard Foundation through a Packard research fellowship. V.N.B. and I.S. gratefully acknowledge assistance from NSF Research at Undergraduate Institutions (RUI) grants AST-1312296 and AST-1909297. Note that findings and conclusions do not necessarily represent views of the NSF. G.C. acknowledges NSF support under grant AST-1817233. J.H.W. acknowledges the funding from the Basic Science Research Program through the National Research Foundation of Korean Government (NRF-2021R1A2C3008486).

Table 10. Virial Products and Derived BH Masses

Object	$c_{\text{cen}} \frac{2}{\text{line}} G$	M_{BH}
	($10^7 M_{\odot}$)	($10^8 M_{\odot}$)
Zw 535 012	$0.85^{+0.36}_{-0.18}$	$0.38^{+0.16}_{-0.08}$
Mrk 1048	$0.50^{+0.60}_{-0.64}$	$0.22^{+0.27}_{-0.29}$
Ark 120	$1.62^{+0.55}_{-0.38}$	$0.73^{+0.25}_{-0.17}$
Mrk 9	$0.91^{+0.21}_{-0.41}$	$0.41^{+0.09}_{-0.18}$
Mrk 704	$2.30^{+0.91}_{-0.79}$	$1.03^{+0.41}_{-0.36}$
MCG +04 22 042	$0.34^{+0.07}_{-0.05}$	$0.15^{+0.03}_{-0.02}$
Mrk 110	$0.77^{+0.14}_{-0.15}$	$0.35^{+0.06}_{-0.07}$
RBS 1303	$0.57^{+0.09}_{-0.12}$	$0.25^{+0.04}_{-0.06}$
Mrk 841	$1.04^{+0.58}_{-0.35}$	$0.47^{+0.26}_{-0.16}$
Mrk 1392	$1.40^{+0.17}_{-0.20}$	$0.63^{+0.08}_{-0.09}$
SBS 1518+593	$1.23^{+0.49}_{-0.56}$	$0.55^{+0.22}_{-0.25}$
3C 382	$3.12^{+2.51}_{-2.19}$	$1.39^{+1.12}_{-0.98}$
NPM1G +27.0587	$0.51^{+0.37}_{-0.30}$	$0.23^{+0.16}_{-0.13}$
RXJ 2044.0+2833	$0.27^{+0.04}_{-0.04}$	$0.12^{+0.02}_{-0.02}$
PG 2209+184	$0.66^{+0.14}_{-0.14}$	$0.29^{+0.06}_{-0.06}$
RBS 1917	$0.32^{+0.12}_{-0.12}$	$0.14^{+0.06}_{-0.05}$

NOTE Assuming $f = 4.47$ (Woo et al. 2015).

A.V.F.'s group at U.C. Berkeley is grateful for support from the TABASGO Foundation, the Christopher R. Redlich Fund, the Miller Institute for Basic Research in Science (in which he is a Miller Senior Fellow), and many individual donors. K.H. acknowledges support from STFC grant ST/R000824/1. We acknowledge the generous support of Marc J. Staley, whose fellowship partly funded B.E.S. whilst contributing to the work presented herein as a graduate student. I.S. acknowledges support from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy EXC 2121 "Quantum Universe" 390833306. Research by S.V. is supported by NSF grants AST 1813176 and AST-2008108.

This work makes use of observations from the LCOGT network. The Liverpool Telescope is operated on the island of La Palma by Liverpool John Moores University

in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias with financial support from the UK Science and Technology Facilities Council. Based on observations acquired at the Observatorio Astronómico Nacional in the Sierra San Pedro Mártir (OAN-SPM), Baja California, México, we thank the daytime and night support staff at the OAN-SPM for facilitating and helping obtain our observations. Some of the data used in this paper were acquired with the RATIR instrument, funded by the University of California and NASA Goddard Space Flight Center, and the 1.5-meter Harold L. Johnson telescope at the Observatorio Astronómico Nacional on the Sierra de San Pedro Mártir, operated and maintained by the Observatorio Astronómico Nacional and the Instituto de Astronomía of the Universidad Nacional Autónoma de México. Operations are partially funded by the Universidad Nacional Autónoma de México (DGAPA/PAPIIT IG100414, IT102715, AG100317, IN109418, IG100820, and IN105921). We acknowledge the contribution of Leonid Georgiev and Neil Gehrels to the development of RATIR. This research was made possible through the use of the AAVSO Photometric All-Sky Survey (APASS), funded by the Robert Martin Ayers Sciences Fund and NSF grant AST-1412587 and contributed by observers worldwide. We acknowledge the use of The AGN Black Hole Mass Database as a compilation of some of the reverberation mapped black hole masses prior to 2015 (Bentz & Katz 2015). This research has made use of the NASA/IPAC Extragalactic Database (NED), which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

Facility: Shane (Kast double spectrograph), KAIT, Nickel, LCOGT, MLO:1m, OANSPM:1.5m, FLWO:1.2m, BYU:0.9m, Liverpool:2m, AAVSO

Software: mp t (Markwardt 2009), CARMA (Kelly et al. 2009, 2014), IDL Astronomy User's Library (Landsman 1993), IRAF (Tody 1986), CAMEL (Pancoast et al. 2011, 2014a, Villafañe et al., in prep.), mapspec (Fausnaugh 2017)

APPENDIX

A. COMPARISON WITH PRIOR M_{BH} MEASUREMENTS

RM studies over the years have shown the intrinsic scatter among BH mass measurements from the RM technique to be inherently small, demonstrating its robustness for measuring BH masses. In the well-studied case of NGC 5548 with many epochs of RM-based BH masses (see Dalla Bontà et al. 2020, for a recent compilation), the distribution of

these masses exhibits an intrinsic scatter of about 0.2 dex, suggesting that reverberation BH mass measurements for individual sources may typically fall within this range. Here we examine the level of agreement between our results and those from the literature for the AGN where BH mass measurements from previous RM campaigns are available. Given the expected ~ 0.2 dex scatter in mass estimates obtained from light curves at different epochs, our measurements for the majority of these sources are generally consistent with past M_{BH} estimates, with a factor of 2.7 difference (0.44 dex) in the most discrepant case.

A.1. *Ark 120*

Ark 120 was monitored by Peterson et al. (1998) and Peterson et al. (2004) who had measured a rest-frame H β lag of $47.1^{+8.3}_{-12.4}$ and $37.1^{+4.8}_{-5.4}$ days and v_{line} of 1959 \pm 109 and 1884 \pm 48 km s $^{-1}$ from two different epochs, respectively. Adopting the same f of 4.47 from Woo et al. (2015) as in this study, the black hole masses calculated are $1.57^{+0.45}_{-0.59} \times 10^8 M_{\odot}$ and $1.15^{+0.21}_{-0.23} \times 10^8 M_{\odot}$, respectively.

In comparison, we measured a shorter lag of $18.7^{+5.9}_{-4.5}$ days in H β , resulting in a smaller black hole mass of $0.73^{+0.25}_{-0.17} \times 10^8 M_{\odot}$. In a concurrent campaign, the MAHA group determined an H β lag of $16.2^{+3.2}_{-3.1}$ days and consequently a black hole mass of $0.68^{+0.14}_{-0.13} \times 10^8 M_{\odot}$ (using the same virial factor) in Ark 120 (Du et al. 2018a). Our measurement ($\log(M_{\text{BH}}/M_{\odot}) = 7.86$) is very similar to the MAHA result ($\log(M_{\text{BH}}/M_{\odot}) = 7.83$), and roughly agree with the previous masses ($\log(M_{\text{BH}}/M_{\odot}) = 8.20$) from over a decade ago within ~ 0.2 dex, indicating that the state of the AGN and the intrinsic BLR properties stay generally consistent in the past two decades.

A.2. *Mrk 704*

Mrk 704 was among the bright Seyfert 1 galaxies that were targeted by De Rosa et al. (2018) in an RM campaign in early 2012. De Rosa et al. (2018) measured an H β lag of $12.65^{+1.49}_{-2.14}$ days and v_{line} of 1860^{+108}_{-130} km s $^{-1}$, resulting in a black hole mass of $0.43^{+0.16}_{-0.12} \times 10^8 M_{\odot}$, or $\log(M_{\text{BH}}/M_{\odot}) = 7.63$. Four years later, we determined a longer lag of $28.9^{+10.2}_{-10.0}$ days and a larger v_{line} of 2009 ± 102 km s $^{-1}$. Our black hole mass measurement of $1.03^{+0.41}_{-0.36} \times 10^8 M_{\odot}$, or $\log(M_{\text{BH}}/M_{\odot}) = 8.01$, is more than twice as large as the result from De Rosa et al. (2018), but our measurement's uncertainty is substantial such that there is statistical agreement within 1.2 σ .

A.3. *Mrk 110*

Like Ark 120, Mrk 110 was another RM target monitored by Peterson et al. (1998) and Peterson et al. (2004) within the time period and had a rest-frame H β lag of $20.4^{+10.5}_{-6.3}$, $24.3^{+5.5}_{-8.3}$, and $33.3^{+14.9}_{-10.0}$ days with v_{line} of 1115 \pm 103, 1196 \pm 141, 755 \pm 29 km s $^{-1}$ from three different epochs, respectively. Adopting the same f of 4.47 from Woo et al. (2015) as in this study, the black hole mass calculated correspond to $2.2^{+1.5}_{-1.1} \times 10^7 M_{\odot}$, $3.0^{+1.4}_{-1.8} \times 10^7 M_{\odot}$, and $1.7^{+0.9}_{-0.6} \times 10^7 M_{\odot}$, or a combined value of $2.3^{+0.8}_{-0.7} \times 10^7 M_{\odot}$ ($\log(M_{\text{BH}}/M_{\odot}) = 7.36$).

With a measured H β lag of $27.8^{+4.3}_{-5.1}$ days and a v_{line} of 1314 ± 69 km s $^{-1}$, we obtained a larger black hole mass of $3.5^{+0.6}_{-0.7} \times 10^7 M_{\odot}$ ($\log(M_{\text{BH}}/M_{\odot}) = 7.54$), consistent with the previous estimate within statistical uncertainties (1 σ) and ~ 0.2 dex.

A.4. *SBS 1518+593*

SBS 1518+593 was one target from the MAHA campaign carried out by Du et al. (2018a) during a similar time frame as our LAMP study. Du et al. (2018a) established an H β lag of $19.7^{+9.9}_{-6.0}$ days and v_{line} of 1038 ± 20 km s $^{-1}$. The inferred black hole mass adjusting for f is $1.98^{+1.00}_{-0.61} \times 10^7 M_{\odot}$, or $\log(M_{\text{BH}}/M_{\odot}) = 7.30$. Measuring a similar lag of $20.1^{+8.4}_{-8.9}$ days and larger v_{line} of 2249 ± 255 km s $^{-1}$, our black hole mass for this AGN was $5.5^{+2.2}_{-2.5} \times 10^7 M_{\odot}$, or $\log(M_{\text{BH}}/M_{\odot}) = 7.74$. Our measurements agree statistically within 1.3 σ given the large uncertainties, with a difference roughly within ~ 0.2 dex. However, our velocity-resolved lag spectra disagree qualitatively – Du et al. (2018a) found the BLR kinematics to be in virialized motion while we see infalling behavior. This discrepancy may be due to differences in our respective campaigns' monitoring durations, where they observed SBS 1518+593 for approximately 180 days while our coverage spanned 360 days. This discrepancy may also be due to the fact that our H β light curve exhibits much noise and scatter, which may in turn affect our velocity-resolved measurement.

A.5. *Zw 535-012*

The black hole mass in Zw 535-012 was previously determined to be $2.6 \times 10^7 M_{\odot}$ (Wang et al. 2009), or $\log(M_{\text{BH}}/M_{\odot}) = 7.41$. It was computed using the scaling law from Greene & Ho (2005), assuming $\log L(\text{H } \beta) =$

42.1 erg s^{-1} and $\text{FWHM}(H\beta) = 2555 \text{ km s}^{-1}$ from a single-epoch spectroscopic observation in 2006. In our campaign, we obtained a black hole mass of $3.8^{+1.6}_{-0.8} \times 10^7 M_{\odot}$, or $\log(M_{\text{BH}}/M_{\odot}) = 7.58$, which is 1.4 times larger than the previous estimate but consistent within statistical uncertainties (0.6σ) and 0.2 dex .

REFERENCES

- Arévalo, P., Uttley, P., Kaspi, S., et al. 2008, *MNRAS*, 389, 1479, doi: 10.1111/j.1365-2966.2008.13719.x
- Bachev, R., Strigachev, A., Semkov, E., & Mihov, B. 2008, *A&A*, 488, 887, doi: 10.1051/0004-6361:200810030
- Barth, A. J., Pancoast, A., Thorman, S. J., et al. 2011, *ApJL*, 743, L4, doi: 10.1088/2041-8205/743/1/L4
- Barth, A. J., Pancoast, A., Bennert, V. N., et al. 2013, *ApJ*, 769, 128, doi: 10.1088/0004-637X/769/2/128
- Barth, A. J., Bennert, V. N., Canalizo, G., et al. 2015, *ApJS*, 217, 26, doi: 10.1088/0067-0049/217/2/26
- Bennert, V. N., Treu, T., Auger, M. W., et al. 2015, *ApJ*, 809, 20, doi: 10.1088/0004-637X/809/1/20
- Bentz, M. C., & Katz, S. 2015, *PASP*, 127, 67, doi: 10.1086/679601
- Bentz, M. C., Peterson, B. M., Netzer, H., Pogge, R. W., & Vestergaard, M. 2009a, *ApJ*, 697, 160, doi: 10.1088/0004-637X/697/1/160
- Bentz, M. C., Peterson, B. M., Pogge, R. W., Vestergaard, M., & Onken, C. A. 2006, *ApJ*, 644, 133, doi: 10.1086/503537
- Bentz, M. C., Williams, P. R., Street, R., et al. 2021, arXiv e-prints, arXiv:2108.00482, <https://arxiv.org/abs/2108.00482>
- Bentz, M. C., Walsh, J. L., Barth, A. J., et al. 2009b, *ApJ*, 705, 199, doi: 10.1088/0004-637X/705/1/199
- . 2010a, *ApJ*, 716, 993, doi: 10.1088/0004-637X/716/2/993
- Bentz, M. C., Horne, K., Barth, A. J., et al. 2010b, *ApJL*, 720, L46, doi: 10.1088/2041-8205/720/1/L46
- Bentz, M. C., Denney, K. D., Grier, C. J., et al. 2013, *ApJ*, 767, 149, doi: 10.1088/0004-637X/767/2/149
- Bian, W.-H., Fang, L.-L., Huang, K.-L., & Wang, J.-M. 2012, *MNRAS*, 427, 2881, doi: 10.1111/j.1365-2966.2012.22123.x
- Blandford, R. D., & McKee, C. F. 1982, *ApJ*, 255, 419, doi: 10.1086/159843
- Boroson, T., Brown, T., Hjelmstrom, A., et al. 2014, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 9149, *Observatory Operations: Strategies, Processes, and Systems V*, 91491E, doi: 10.1117/12.2054776
- Boroson, T. A., & Green, R. F. 1992, *ApJS*, 80, 109, doi: 10.1086/191661
- Brown, T. M., Baliber, N., Bianco, F. B., et al. 2013, *PASP*, 125, 1031, doi: 10.1086/673168
- Bruzual, G., & Charlot, S. 2003, *MNRAS*, 344, 1000, doi: 10.1046/j.1365-8711.2003.06897.x
- Butler, N., Klein, C., Fox, O., et al. 2012, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 8446, *Ground-based and Airborne Instrumentation for Astronomy IV*, ed. I. S. McLean, S. K. Ramsay, & H. Takami, 844610, doi: 10.1117/12.926471
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245, doi: 10.1086/167900
- Chatterjee, R., Jorstad, S. G., Marscher, A. P., et al. 2008, *ApJ*, 689, 79, doi: 10.1086/592598
- Croton, D. J., Springel, V., White, S. D. M., et al. 2006, *MNRAS*, 365, 11, doi: 10.1111/j.1365-2966.2005.09675.x
- Dalla Bontà, E., Peterson, B. M., Bentz, M. C., et al. 2020, *ApJ*, 903, 112, doi: 10.3847/1538-4357/abbc1c
- De Rosa, G., Fausnaugh, M. M., Grier, C. J., et al. 2018, *ApJ*, 866, 133, doi: 10.3847/1538-4357/aadd11
- Denney, K. D., Peterson, B. M., Pogge, R. W., et al. 2009, *ApJL*, 704, L80, doi: 10.1088/0004-637X/704/2/L80
- . 2010, *ApJ*, 721, 715, doi: 10.1088/0004-637X/721/1/715
- Drake, A. J., Djorgovski, S. G., Mahabal, A., et al. 2009, *ApJ*, 696, 870, doi: 10.1088/0004-637X/696/1/870
- Du, P., & Wang, J.-M. 2019, *ApJ*, 886, 42, doi: 10.3847/1538-4357/ab4908
- Du, P., Hu, C., Lu, K.-X., et al. 2014, *ApJ*, 782, 45, doi: 10.1088/0004-637X/782/1/45
- . 2015, *ApJ*, 806, 22, doi: 10.1088/0004-637X/806/1/22
- Du, P., Lu, K.-X., Zhang, Z.-X., et al. 2016a, *ApJ*, 825, 126, doi: 10.3847/0004-637X/825/2/126
- Du, P., Lu, K.-X., Hu, C., et al. 2016b, *ApJ*, 820, 27, doi: 10.3847/0004-637X/820/1/27
- Du, P., Brotherton, M. S., Wang, K., et al. 2018a, *ApJ*, 869, 142, doi: 10.3847/1538-4357/aaed2c
- Du, P., Zhang, Z.-X., Wang, K., et al. 2018b, *ApJ*, 856, 6, doi: 10.3847/1538-4357/aaae6b
- Fausnaugh, M. M. 2017, *PASP*, 129, 024007, doi: 10.1088/1538-3873/129/972/024007
- Ferrarese, L., & Merritt, D. 2000, *ApJL*, 539, L9, doi: 10.1086/312838
- Filippenko, A. V. 1982, *PASP*, 94, 715, doi: 10.1086/131052

- Filippenko, A. V., Li, W. D., Treers, R. R., & Modjaz, M. 2001, in *Astronomical Society of the Pacific Conference Series*, Vol. 246, IAU Colloq. 183: Small Telescope Astronomy on Global Scales, ed. B. Paczynski, W.-P. Chen, & C. Lemme, 121
- Fonseca Alvarez, G., Trump, J. R., Homayouni, Y., et al. 2019, arXiv e-prints, arXiv:1910.10719. <https://arxiv.org/abs/1910.10719>
- Gebhardt, K., Bender, R., Bower, G., et al. 2000, *ApJL*, 539, L13, doi: 10.1086/312840
- Greene, J. E., & Ho, L. C. 2005, *ApJ*, 630, 122, doi: 10.1086/431897
- Grier, C. J., Peterson, B. M., Pogge, R. W., et al. 2012, *ApJ*, 755, 60, doi: 10.1088/0004-637X/755/1/60
- Grier, C. J., Peterson, B. M., Horne, K., et al. 2013, *ApJ*, 764, 47, doi: 10.1088/0004-637X/764/1/47
- Grier, C. J., Trump, J. R., Shen, Y., et al. 2017, *ApJ*, 851, 21, doi: 10.3847/1538-4357/aa98dc
- Grier, C. J., Shen, Y., Horne, K., et al. 2019, *ApJ*, 887, 38, doi: 10.3847/1538-4357/ab4ea5
- Guo, H., Wang, J., Cai, Z., & Sun, M. 2017, *ApJ*, 847, 132, doi: 10.3847/1538-4357/aa8d71
- Henden, A. A., Templeton, M., Terrell, D., et al. 2016, *VizieR Online Data Catalog*, II/336
- Hoormann, J. K., Martini, P., Davis, T. M., et al. 2019, *MNRAS*, 487, 3650, doi: 10.1093/mnras/stz1539
- Horne, K., Peterson, B. M., Collier, S. J., & Netzer, H. 2004, *PASP*, 116, 465, doi: 10.1086/420755
- Horne, K., De Rosa, G., Peterson, B. M., et al. 2021, *ApJ*, 907, 76, doi: 10.3847/1538-4357/abce60
- Hu, C., Du, P., Lu, K.-X., et al. 2015, *ApJ*, 804, 138, doi: 10.1088/0004-637X/804/2/138
- Joshi, R., Chand, H., Wiita, P. J., Gupta, A. C., & Srikanand, R. 2012, *MNRAS*, 419, 3433, doi: 10.1111/j.1365-2966.2011.19985.x
- Kasliwal, V. P., Vogeley, M. S., & Richards, G. T. 2015, *MNRAS*, 451, 4328, doi: 10.1093/mnras/stv1230
- Kaspi, S., Maoz, D., Netzer, H., et al. 2005, *ApJ*, 629, 61, doi: 10.1086/431275
- Kaspi, S., Smith, P. S., Netzer, H., et al. 2000, *ApJ*, 533, 631, doi: 10.1086/308704
- Kelly, B. C., Bechtold, J., & Siemiginowska, A. 2009, *ApJ*, 698, 895, doi: 10.1088/0004-637X/698/1/895
- Kelly, B. C., Becker, A. C., Sobolewska, M., Siemiginowska, A., & Uttley, P. 2014, *ApJ*, 788, 33, doi: 10.1088/0004-637X/788/1/33
- Kilerci Eser, E., Vestergaard, M., Peterson, B. M., Denney, K. D., & Bentz, M. C. 2015, *ApJ*, 801, 8, doi: 10.1088/0004-637X/801/1/8
- Kovačević, J., Popović, L., & Dimitrijević, M. S. 2010, *ApJS*, 189, 15, doi: 10.1088/0067-0049/189/1/15
- Landsman, W. B. 1993, in *Astronomical Society of the Pacific Conference Series*, Vol. 52, *Astronomical Data Analysis Software and Systems II*, ed. R. J. Hanisch, R. J. V. Brissenden, & J. Barnes, 246
- Lang, D., Hogg, D. W., Mierle, K., Blanton, M., & Roweis, S. 2010, *AJ*, 139, 1782, doi: 10.1088/0004-6256/139/5/1782
- Li, S.-S., Yang, S., Yang, Z.-X., et al. 2021, arXiv e-prints, arXiv:2106.05655. <https://arxiv.org/abs/2106.05655>
- Lu, K.-X., Bai, J.-M., Zhang, Z.-X., et al. 2019, *ApJ*, 887, 135, doi: 10.3847/1538-4357/ab5790
- Lu, K.-X., Wang, J.-G., Zhang, Z.-X., et al. 2021, arXiv e-prints, arXiv:2106.10589. <https://arxiv.org/abs/2106.10589>
- MacLeod, C. L., Ivezić, J., Kochanek, C. S., et al. 2010, *ApJ*, 721, 1014, doi: 10.1088/0004-637X/721/2/1014
- Markwardt, C. B. 2009, in *Astronomical Society of the Pacific Conference Series*, Vol. 411, *Astronomical Data Analysis Software and Systems XVIII*, ed. D. A. Bohlender, D. Durand, & P. Dowler, 251. <https://arxiv.org/abs/0902.2850>
- Martínez-Aldama, M. L., Czerny, B., Kawka, D., et al. 2019, *ApJ*, 883, 170, doi: 10.3847/1538-4357/ab3728
- Martínez-Aldama, M. L., Zajaček, M., Czerny, B., & Panda, S. 2020, *ApJ*, 903, 86, doi: 10.3847/1538-4357/abb6f8
- Marziani, P., Sulentic, J. W., Zamanov, R., et al. 2003, *ApJS*, 145, 199, doi: 10.1086/346025
- McHardy, I. M., Koeding, E., Knigge, C., Uttley, P., & Fender, R. P. 2006, *Nature*, 444, 730, doi: 10.1038/nature05389
- Miller, J. S., & Stone, R. P. S. 1994, *Lick Observatory Technical Reports*, 66
- Mushotzky, R. F., Edelson, R., Baumgartner, W., & Gandhi, P. 2011, *ApJL*, 743, L12, doi: 10.1088/2041-8205/743/1/L12
- Pancoast, A., Brewer, B. J., & Treu, T. 2011, *ApJ*, 730, 139, doi: 10.1088/0004-637X/730/2/139
- . 2014a, *MNRAS*, 445, 3055, doi: 10.1093/mnras/stu1809
- Pancoast, A., Brewer, B. J., Treu, T., et al. 2014b, *MNRAS*, 445, 3073, doi: 10.1093/mnras/stu1419
- Pei, L., Barth, A. J., Aldering, G. S., et al. 2014, *ApJ*, 795, 38, doi: 10.1088/0004-637X/795/1/38
- Pei, L., Fausnaugh, M. M., Barth, A. J., et al. 2017, *ApJ*, 837, 131, doi: 10.3847/1538-4357/aa5eb1
- Penton, A., Malik, U., Davis, T., et al. 2021, arXiv e-prints, arXiv:2101.06921. <https://arxiv.org/abs/2101.06921>
- Peterson, B. M. 1993, *PASP*, 105, 247, doi: 10.1086/133140

- Peterson, B. M., Wanders, I., Bertram, R., et al. 1998, *ApJ*, 501, 82, doi: 10.1086/305813
- Peterson, B. M., Ferrarese, L., Gilbert, K. M., et al. 2004, *ApJ*, 613, 682, doi: 10.1086/423269
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2016, *A&A*, 594, A13, doi: 10.1051/0004-6361/201525830
- Richards, G. T., Kruczek, N. E., Gallagher, S. C., et al. 2011, *AJ*, 141, 167, doi: 10.1088/0004-6256/141/5/167
- Sergeev, S. G. 2020, *MNRAS*, 495, 971, doi: 10.1093/mnras/staa1210
- Shen, Y. 2013, *Bulletin of the Astronomical Society of India*, 41, 61. <https://arxiv.org/abs/1302.2643>
- Shen, Y., Greene, J. E., Strauss, M. A., Richards, G. T., & Schneider, D. P. 2008, *ApJ*, 680, 169, doi: 10.1086/587475
- Shen, Y., & Liu, X. 2012, *ApJ*, 753, 125, doi: 10.1088/0004-637X/753/2/125
- Shen, Y., Richards, G. T., Strauss, M. A., et al. 2011, *ApJS*, 194, 45, doi: 10.1088/0067-0049/194/2/45
- Shen, Y., Brandt, W. N., Dawson, K. S., et al. 2015, *ApJS*, 216, 4, doi: 10.1088/0067-0049/216/1/4
- Shen, Y., Horne, K., Grier, C. J., et al. 2016, *ApJ*, 818, 30, doi: 10.3847/0004-637X/818/1/30
- Smith, C. E., & Nelson, B. 1969, *PASP*, 81, 74, doi: 10.1086/128742
- Smith, K. L., Mushotzky, R. F., Boyd, P. T., et al. 2018, *ApJ*, 857, 141, doi: 10.3847/1538-4357/aab88d
- Springel, V., Di Matteo, T., & Hernquist, L. 2005, *MNRAS*, 361, 776, doi: 10.1111/j.1365-2966.2005.09238.x
- Steele, I. A., Smith, R. J., Rees, P. C., et al. 2004, *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 5489, *The Liverpool Telescope: performance and first results*, ed. J. Oschmann, Jacobus M., 679–692, doi: 10.1117/12.551456
- Sun, J., & Shen, Y. 2015, *ApJL*, 804, L15, doi: 10.1088/2041-8205/804/1/L15
- Tody, D. 1986, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 627, *Instrumentation in astronomy VI*, ed. D. L. Crawford, 733, doi: 10.1117/12.968154
- Uttley, P., Edelson, R., McHardy, I. M., Peterson, B. M., & Markowitz, A. 2003, *ApJL*, 584, L53, doi: 10.1086/373887
- van Groningen, E., & Wanders, I. 1992, *PASP*, 104, 700, doi: 10.1086/133039
- vanden Berk, D., Yip, C., Connolly, A., Jester, S., & Stoughton, C. 2004, in *Astronomical Society of the Pacific Conference Series*, Vol. 311, *AGN Physics with the Sloan Digital Sky Survey*, ed. G. T. Richards & P. B. Hall, 21. <https://arxiv.org/abs/astro-ph/0310840>
- Véron-Cetty, M. P., Joly, M., & Véron, P. 2004, *A&A*, 417, 515, doi: 10.1051/0004-6361:20035714
- Vestergaard, M., & Peterson, B. M. 2006, *ApJ*, 641, 689, doi: 10.1086/500572
- Wandel, A., Peterson, B. M., & Malkan, M. A. 1999, *ApJ*, 526, 579, doi: 10.1086/308017
- Wang, J., Mao, Y. F., & Wei, J. Y. 2009, *AJ*, 137, 3388, doi: 10.1088/0004-6256/137/2/3388
- Wang, J.-M., Du, P., Hu, C., et al. 2014, *ApJ*, 793, 108, doi: 10.1088/0004-637X/793/2/108
- Watson, A. M., Richer, M. G., Bloom, J. S., et al. 2012, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 8444, *Ground-based and Airborne Telescopes IV*, ed. L. M. Stepp, R. Gilmozzi, & H. J. Hall, 84445L, doi: 10.1117/12.926927
- Welsh, W. F., & Horne, K. 1991, *ApJ*, 379, 586, doi: 10.1086/170530
- White, R. J., & Peterson, B. M. 1994, *PASP*, 106, 879, doi: 10.1086/133456
- Wilhite, B. C., Brunner, R. J., Grier, C. J., Schneider, D. P., & vanden Berk, D. E. 2008, *MNRAS*, 383, 1232, doi: 10.1111/j.1365-2966.2007.12655.x
- Williams, P. R., Pancoast, A., Treu, T., et al. 2018, *ApJ*, 866, 75, doi: 10.3847/1538-4357/aae086
- . 2020, *ApJ*, 902, 74, doi: 10.3847/1538-4357/abbad7
- Winter, L. M., Lewis, K. T., Koss, M., et al. 2010, *ApJ*, 710, 503, doi: 10.1088/0004-637X/710/1/503
- Woo, J.-H., Yoon, Y., Park, S., Park, D., & Kim, S. C. 2015, *ApJ*, 801, 38, doi: 10.1088/0004-637X/801/1/38
- Wright, E. L. 2006, *PASP*, 118, 1711, doi: 10.1086/510102
- Yuan, F., Lidman, C., Davis, T. M., et al. 2015, *MNRAS*, 452, 3047, doi: 10.1093/mnras/stv1507
- Zhang, Z.-X., Du, P., Smith, P. S., et al. 2019, *ApJ*, 876, 49, doi: 10.3847/1538-4357/ab1099
- Zu, Y., Kochanek, C. S., & Peterson, B. M. 2011, *ApJ*, 735, 80, doi: 10.1088/0004-637X/735/2/80