

DRAFT VERSION FEBRUARY 12, 2024
Typeset using L^AT_EX twocolumn style in AASTeX631

High-quality Extragalactic Legacy-field Monitoring (HELM) with DECam

MING-YANG ZHUANG (庄明阳) ¹, QIAN YANG ^{2,1}, YUE SHEN ^{1,3}, MONIKA ADAMÓW ⁴, DOUGLAS N. FRIEDEL ³,
R. A. GRUENDL ³, XIN LIU ^{1,3,5}, PAUL MARTINI ^{6,7}, TIMOTHY M. C. ABBOTT ⁸, SCOTT F. ANDERSON ⁹,
ROBERTO J. ASSEF ¹⁰, FRANZ E. BAUER ^{11,12,13}, RICH BIELBY ¹⁴, W. N. BRANDT ^{15,16,17}, COLIN J. BURKE ¹,
JORGE CASARES^{18,19}, YU-CHING CHEN ¹, GISELLA DE ROSA ²⁰, ALEX DRLICA-WAGNER ^{21,22,23}, TOM DWELLY ²⁴,
ALICE ELTVEDT¹⁴, GLORIA FONSECA ALVAREZ ²⁵, JIANYANG FU (傅健洋) ¹, CESAR FUENTES ^{26,27,28},
MELISSA L. GRAHAM ²⁹, CATHERINE J. GRIER ^{30,31}, NATHAN GOLOVICH ³², PATRICK B. HALL ³³, PATRICK HARTIGAN ³⁴,
KEITH HORNE ³⁵, ANTON M. KOEKEMOER ²⁰, MIRKO KRUMPE³⁶, JENNIFER I. LI ³⁷, CHRIS LIDMAN ^{38,39},
UMANG MALIK ³⁸, AMELIA MANGIAN ⁴⁰, ANDREA MERLONI ⁴¹, CLAUDIO RICCI ^{42,43}, MARA SALVATO ⁴⁴,
ROB SHARP ³⁸, ZACHARY STONE ^{1,4}, DAVID E. TRILLING ⁴⁵, BRAD E. TUCKER ^{38,46,47}, DI WEN ¹,
ZACHARY WIDEMAN ⁴⁸, YONGQUAN XUE ⁴⁹, ZHEFU YU ⁵⁰ AND CATHERINE ZUCKER ²

¹Department of Astronomy, University of Illinois Urbana-Champaign, Urbana, IL 61801, USA

²Center for Astrophysics | Harvard & Smithsonian, 60 Garden Street, Cambridge, MA 02138, USA

³National Center for Supercomputing Applications, University of Illinois Urbana-Champaign, Urbana, IL 61801, USA

⁴Center for AstroPhysical Surveys, National Center for Supercomputing Applications, University of Illinois Urbana-Champaign, Urbana, IL 61801, USA

⁵Center for Artificial Intelligence Innovation, University of Illinois at Urbana-Champaign, 1205 West Clark Street, Urbana, IL 61801, USA

⁶Department of Astronomy, The Ohio State University, Columbus, Ohio 43210, USA

⁷Center of Cosmology and Astro-Particle Physics, The Ohio State University, Columbus, Ohio 43210, USA

⁸CTIO, NSF's NOIRLab, Casilla 603, La Serena, Chile

⁹Astronomy Department, University of Washington, Box 351580, Seattle, WA 98195, USA

¹⁰Núcleo de Astronomía de la Facultad de Ingeniería y Ciencias, Universidad Diego Portales, Av. Ejército Libertador 441, Santiago, Chile

¹¹Instituto de Astrofísica, Facultad de Física, Pontificia Universidad Católica de Chile, Campus San Joaquín, Av. Vicuña Mackenna 4860, Macul Santiago, Chile, 7820436

¹²Centro de Astroingeniería, Facultad de Física, Pontificia Universidad Católica de Chile, Campus San Joaquín, Av. Vicuña Mackenna 4860, Macul Santiago, Chile, 7820436

¹³Millennium Institute of Astrophysics, Nuncio Monseñor Sótero Sanz 100, Of 104, Providencia, Santiago, Chile

¹⁴Centre for Extragalactic Astronomy, Department of Physics, Durham University, South Road, Durham, DH1 3LE, UK

¹⁵Department of Astronomy & Astrophysics, 525 Davey Lab, The Pennsylvania State University, University Park, PA 16802, USA

¹⁶Institute for Gravitation and the Cosmos, The Pennsylvania State University, University Park, PA 16802, USA

¹⁷Department of Physics, 104 Davey Laboratory, The Pennsylvania State University, University Park, PA 16802, USA

¹⁸Instituto de Astrofísica de Canarias, E-38205 La Laguna, Tenerife, Spain

¹⁹Departamento de Astrofísica, Universidad de La Laguna, E-38206 La Laguna, Tenerife, Spain

²⁰Space Telescope Science Institute, 3700 San Martin Dr., Baltimore, MD 21218, USA

²¹Fermi National Accelerator Laboratory, P.O. Box 500, Batavia, IL 60510, USA

²²Kavli Institute for Cosmological Physics, University of Chicago, Chicago, IL 60637, USA

²³Department of Astronomy and Astrophysics, University of Chicago, Chicago IL 60637, USA

²⁴Max-Planck Institute for Extraterrestrial Physics, Giessenbachstrasse 1, Garching, 85748, Germany

²⁵NSF's NOIRLab, 950 N. Cherry Ave., Tucson, AZ 85719, USA

²⁶Departamento de Astronomía, Universidad de Chile, Camino del Observatorio 1515, Las Condes, Santiago, Chile.

²⁷Centro de Excelencia en Astrofísica y Tecnologías Afines (CATA), Chile

²⁸Millennium Institute of Astrophysics (MAS), Chile

²⁹DIRAC Institute, Department of Astronomy, University of Washington, 3910 15th Avenue NE, Seattle, WA 98195, USA

³⁰Department of Astronomy, University of Wisconsin-Madison, Madison, WI 53706, USA

³¹Steward Observatory, The University of Arizona, 933 North Cherry Avenue, Tucson, AZ 85721, USA

³²Lawrence Livermore National Laboratory, 7000 East Ave, Livermore, CA 94550, USA

³³Department of Physics & Astronomy, York University, 4700 Keele St., Toronto, ON M3J 1P3, Canada

³⁴Department of Physics and Astronomy, Rice University, 6100 S. Main St., TX 77005-1892, USA

³⁵University of St Andrews, SUPA School of Physics & Astronomy, North Haugh, St Andrews KY16 9SS, Scotland, UK

mingyang@illinois.edu

arXiv:2402.06052v1 [astro-ph.GA] 8 Feb 2024

³⁶Leibniz Institute for Astrophysics Potsdam (AIP), An der Sternwarte 16, 14482 Potsdam, Germany

³⁷Department of Astronomy, University of Michigan, Ann Arbor, MI, 48109, USA

³⁸Research School of Astronomy and Astrophysics, Australian National University, Canberra, ACT 2611, Australia

³⁹Centre for Gravitational Astrophysics, College of Science, The Australian National University, Canberra ACT 2601, Australia

⁴⁰Information Trust Institute, University of Illinois Urbana-Champaign, Urbana, IL 61801, USA

⁴¹Max-Planck Institute for Extraterrestrial Physics, Giessenbachstrasse 1, 85748, Garching, Germany

⁴²Núcleo de Astronomía de la Facultad de Ingeniería, Universidad Diego Portales, Av. Ejército Libertador 441, Santiago, Chile

⁴³Kavli Institute for Astronomy and Astrophysics, Peking University, Beijing 100871, China

⁴⁴Max-Planck Institute for Extraterrestrial Physics, Giessenbachstrasse 1, 85748, Germany

⁴⁵Department of Astronomy and Planetary Science, Northern Arizona University, Flagstaff, AZ 86011, USA

⁴⁶National Centre for the Public Awareness of Science, Australian National University, Canberra, ACT 2601, Australia

⁴⁷The Australian Research Council Centre of Excellence for All-Sky Astrophysics in 3 Dimension (ASTRO 3D), Australia

⁴⁸Department of Physics and Astronomy, Texas A&M University, 4242 TAMU, College Station, TX 77843, USA

⁴⁹CAS Key Laboratory for Research in Galaxies and Cosmology, Department of Astronomy, University of Science and Technology of China, Hefei 230026, China

⁵⁰Kavli Institute for Particle Astrophysics and Cosmology, Stanford University, 452 Lomita Mall, Stanford, CA 94305, USA

ABSTRACT

High-quality Extragalactic Legacy-field Monitoring (HELM) is a long-term observing program that photometrically monitors several well-studied extragalactic legacy fields with the Dark Energy Camera (DECam) imager on the CTIO 4m Blanco telescope. Since Feb 2019, HELM has been monitoring regions within COSMOS, XMM-LSS, CDF-S, S-CVZ, ELAIS-S1, and SDSS Stripe 82 with few-day cadences in the $(u)gri(z)$ bands, over a collective sky area of $\sim 38 \text{ deg}^2$. The main science goal of HELM is to provide high-quality optical light curves for a large sample of active galactic nuclei (AGNs), and to build decades-long time baselines when combining past and future optical light curves in these legacy fields. These optical images and light curves will facilitate the measurements of AGN reverberation mapping lags, as well as studies of AGN variability and its dependences on accretion properties. In addition, the time-resolved and coadded DECam photometry will enable a broad range of science applications from galaxy evolution to time-domain science. We describe the design and implementation of the program and present the first data release that includes source catalogs and the first ~ 3.5 years of light curves during 2019A–2022A.

1. INTRODUCTION

Large-scale optical time-domain imaging surveys such as Sloan Digital Sky Survey (SDSS; York et al. 2000) supernova survey (Sako et al. 2018), Pan-STARRS1 Medium Deep Survey (Chambers et al. 2016; Jones et al. 2017), Zwicky Transient Facility (ZTF; Bellm et al. 2019), and Wide Field Survey Telescope (Wang et al. 2023) are rapidly shaping our understanding of the dynamic and variable universe. In addition to transients and explosive events from stellar systems, these time-domain surveys provide extremely valuable information to probe the physical mechanisms that drive the persistent variability from variable stars and active galactic nuclei (AGNs). Variability studies of AGNs also enable the measurements of their black hole masses via the reverberation mapping (RM) technique that measures the time lag between the driving continuum variability and the delayed response in the broad-line emission (e.g., Blandford & McKee 1982; Peterson 1993). In particular, high-quality photometric light curves with long time baselines and adequate cadences are necessary to secure a RM lag measurement and periodicity searches for binary supermassive black hole candidates when combined with dedicated spectroscopic monitoring data (e.g., Shen et al. 2015; Graham et al. 2015).

The SDSS-V Black Hole Mapper (BHM) program (Kollmeier et al. 2017) is conducting a multi-year (2020–2027), multi-object spectroscopic RM campaign (BHM-RM) in several legacy extragalactic fields. In order to provide the required photometric light curves that sample the AGN continuum variability, we are conducting a long-term photometric monitoring campaign with the Dark Energy Camera (DECam; Flaugher et al. 2015) on the 4m CTIO Blanco telescope. The fields chosen by the SDSS-V BHM-RM program coincide with several well-studied extragalactic fields, each with ample multi-wavelength data and/or earlier and future photometric light curves therein (e.g., Lacy et al. 2021; Zou et al. 2022). For example, most of these fields coincide with deep monitoring fields in the Pan-STARRS1 survey (Chambers et al. 2016) and the Dark Energy Survey supernovae program (Abbott et al. 2021). The combined optical light curves for most of these monitoring fields will span more than 25 years once combined with 10-yr light curves from the Legacy Survey of Space and Time (LSST) with the Vera C. Rubin Observatory. Therefore these DECam monitoring data will provide legacy value for broad science in these fields.

In this paper we provide an overview of the DECam monitoring program and its current status, and present the first data

release¹ from the first ~ 3.5 years of observations (2019-2022). In Section 2 we provide a technical overview of the High-quality Extragalactic Legacy-field Monitoring (HELM) program. In Section 3 we describe the data products, with examples to showcase the science applications. We summarize and provide an outlook in Section 4. We adopt AB magnitude system throughout the paper.

2. OBSERVATIONS AND DATA PROCESSING

HELM started observations in the 2019A semester (Feb-Jul, 2019) and is currently ongoing. During the early phase of the program (2019A-2020B), a cadence of ~ 6 days was adopted. Since 2021A, the cadence was increased to ~ 3 days and a new field (South-Continuous Viewing Zone; S-CVZ) was added to the field list. For each epoch, a set of consecutive exposures in the *gri* bands are taken, with 2 back-to-back exposures for each band. The nominal per-epoch exposure times are 160 s, 140 s, and 180 s in the three bands, respectively. A total of 14 DECam pointings (each with a 2.7 deg^2 field-of-view) received regular monitoring, most of which are for the BHM-RM program. However, some pointings are included in different programs (PIs: X. Liu, P. Martini) that have slightly different science goals, including periodicity searches of binary supermassive black hole candidates and RM lag measurement. We consolidated these different DECam monitoring programs with AGN science focuses and processed these data with the same pipeline. Table 1 summarizes the details of the monitoring fields. Three fields (CDF-S, XMM-LSS, and COSMOS) contain three pointings, while two fields (ELAIS-S1 and SDSS Stripe 82) contain two pointings. The layouts of the HELM fields are shown in Figure 1. These particular pointings are aligned with past monitoring within the DES supernovae program (e.g., Bernstein et al. 2012; Smith et al. 2020).

The first 2 semesters of HELM (2019A-2019B) were carried out on shared nights with other non-monitoring DECam programs. Since 2020A, HELM has been part of a collection of regular DECam monitoring programs that pool observing time and observers. In a few subsequent semesters, to improve the overall observing efficiency, some pointings of HELM were observed under the public DECam monitoring program that also targeted the COSMOS and ELAIS-S1 fields (Graham et al. 2023), but to greater depths than those of the HELM program. These public DECam data are included in our data reduction and data products. The monitoring program suffered significant time loss in 2020A due to

the COVID-19 pandemic, and substantial loss in the second half of 2020B due to DECam instrumental problems.

At the beginning of the program during 2019A, we observed some fields with additional filters to provide more wavelength coverage. The S-CVZ field was observed in the *ugriz* filters. For these additional imaging, *griz* has 61 available CCD chips (except CCD #61), while *u* has 60 chips with an additional CCD #2 not available. Table 1 summarizes the HELM observations including fields, pointings, observing program IDs, filters, and numbers of exposures up to 13 September 2022. Figure 2 illustrates the cadence of each field.

2.1. Data Reduction

The raw HELM observations were processed using the Dark Energy Survey (DES) “Final Cut” pipeline (see Morganson et al. 2018, for an overview) with the same configuration of `SExtractor` (Bertin & Arnouts 1996) as that used in the production campaigns for the second data release (DR2) of DES (DES-DR2; Abbott et al. 2021) and DECam Local Volume Exploration-DR2 (Drlica-Wagner et al. 2022). During the years that the DES Collaboration was actively taking data, the calibration products (e.g., bias, flat, sky-subtraction templates, secondary flat-field or “star-flats”, and the relative astrometric offsets among the CCDs) were monitored and updated as needed as part of that program. Processing of observations taken after the DES observations ended in January 2019 have, thus far, relied on the DES legacy calibration products from that period. In Section 2.2 we assess the overall systematic astrometric and photometric performance to understand the limitations this might have on the temporal monitoring by HELM.

2.1.1. Magnitude Zeropoint

A magnitude zeropoint (zp) pipeline is run separately for all of the CCD images. We use the `expCalib` (S. Allam & R. Gruendl, private communication) code to match objects in the Final Cut catalogs to the APASS 9 catalog (Henden et al. 2015). A $1''$ -radius match was performed for each CCD in one exposure. If the number of matched objects is low or a match was not found, the average zp value among all CCDs is adopted. The `expCalib` will not return a match (or will return a low number of matches) in two cases: - APASS catalog has no coverage in the requested region of the sky, or the magnitude range of stars in the APASS catalog does not overlap with the brightness of objects detected in the CCD (APASS sources are brighter and appear saturated in the DECam data). Among 7816 successfully reduced exposures (corresponding to 476743 CCD individual images), magnitude zero points are available from the `expCalib` for all but 124 exposures (2, 49, 28, 44, and 1 in the *ugriz* bands, respectively) and 7562 CCD images (1.6%).

¹ <https://ariel.astro.illinois.edu/helm/>

Table 1. HELM observation summary

Field	Pointing	Pointing center	ProgID	Filter	Nexp	5σ depth (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)
CDF-S	C1	03h37m06s -27d06m43s	C	<i>griz</i>	54-36-36-18	22.9-23.3-23.1-22.6
	C2	03h37m06s -29d05m18s	A-C	<i>griz</i>	54-36-36-18	
	C3	03h30m36s -28d06m01s	A-C	<i>griz</i>	248-190-187-34	
COSMOS	CO1	10h00m00s +03d05m56s	A-C-E-F-G	<i>gri</i>	362-295-295	22.8-23.1-22.9
	CO2	09h56m53s +01d44m58s	A-E-F-G	<i>gri</i>	278-232-235	
	CO3	10h03m07s +01d44m59s	A-E-F-G	<i>gri</i>	267-226-232	
ELAIS-S1	E1	00h31m30s -43d00m35s	A-C-E-F-G	<i>griz</i>	387-340-349-28	22.8-23.2-23.1-22.5
	E2	00h38m00s -43d59m53s	A-C-E-F-G	<i>griz</i>	365-332-322-302	
S-CVZ	S-CVZ	06h00m00s -66d33m39s	A-C	<i>ugriz</i>	33-223-196-196-24	21.1-22.8-22.9-22.7-22.0
Stripe 82	S1	02h51m17s -00d00m03s	A-B-C-H	<i>griz</i>	87-50-47-16	22.7-23.5-23.4-22.8
	S2	02h44m47s -00d59m20s	A-B-C-H	<i>griz</i>	82-47-47-17	
XMM-LSS	X1	02h17m54s -04d55m48s	A-C	<i>griz</i>	127-107-102-20	22.6-22.9-22.7-22.4
	X2	02h22m40s -06d24m46s	A-C	<i>griz</i>	127-107-102-20	
	X3	02h25m48s -04d36m04s	A-C	<i>griz</i>	127-107-102-20	

NOTE—Col. (1) Field name. Col. (2) Pointing name. Col. (3) Median coordinate of pointing center. Typical pointing accuracy of each exposure is $\sim 10''$. Col. (4) Program ID. A: 2019A-0065; B: 2019B-0219; C: 2019B-0910; D: 2021A-0037; E: 2021A-0113; F: 2021B-0149; G: 2022A-724693; H: 2022B-175073. Col. (5) Filter used. Col. (6) Number of successfully reduced exposures in each band, separated by dashes. Most of the nightly epochs in the HELM program consist of two back-to-back exposures per pointing per filter. Col. (7) Median 5σ depth of individual exposures for all pointings of a given field estimated from m_{PSF} .

Using four fields associated with 10 pointings of our HELM DECam monitoring program (CDF-S, ELAIS-S1, SDSS Stripe 82, and XMM-LSS) that are covered by the footprint of 6 years of DES science operations of ~ 5000 deg² from DES-DR2 (Abbott et al. 2021), we derive independent magnitude zeropoints for all exposures by cross-matching sources detected in our observations with those from the DES-DR2 source catalog. The DES-DR2 catalog is constructed using coadded images, which has typically ~ 10 times longer exposure time than our single-exposure images. Based on the extent of the source, we derive two versions of magnitude zeropoints, one based on point sources using m_{PSF} and another based on all the sources using m_{AUTO} to increase the number of available objects. We first select all the reliably extracted (IMAFLAGS_ISO=0, FLAGS<4), high signal-to-noise ratio (SNR; m_{PSF} error < 0.1 mag, which corresponds to an SNR of 10) sources from our observations. We use a relatively loose criterion of CLASS_STAR > 0.8 to select point-like sources, which ensures we do not miss a significant number of point sources and avoids including many extended sources. For the DES-DR2 catalog, we apply the same selection criteria of object quality (IMAFLAGS_ISO=0, FLAGS<4 in i band) and select point-like sources using $0 \leq \text{EXTENDED_COADD} \leq 1$ following Abbott et al. (2021).

The magnitude zp of each CCD chip is derived using the 3σ clipped median difference between magnitudes of objects from our single-exposure catalog and those from DES-DR2 catalog, with associated error equal to the standard deviation divided by the root of the number of the remaining objects. Figure 3 compares the number of objects used to derive the magnitude zps. Magnitude zps derived from all sources make use of on average 2.7 times more objects compared to those derived from point sources only (295 versus 108). Limiting to CCD images with at least 10 applicable objects for zp calibration, we find a small difference of 0.009 mag with 16th and 84th percentiles of the distribution of -0.008 and 0.019 mag between the two sets of zp, respectively (Figure 4). We find that the difference between two zps is correlated with seeing, with larger m_{PSF} -based zp in exposures with better seeing. This may be related to slight difference in aperture loss at different seeing conditions. We also compare the zp from the DECam pipeline with that from DES-DR2 using all sources (Table 2). The median differences are < 0.01 mag (slightly larger with 0.014 mag in the E2 field) with a scatter of $\lesssim 0.04$ mag. The final zp of a CCD image is determined with the priority decreasing from DES-DR2 AUTO magnitude to DES-DR2 PSF magnitude to DECam pipeline. We only adopt DES-DR2-based zp if the number of objects used to calibrate zp is greater or equal to 10. Finally, we success-




Figure 1. The layout of HELM pointings in (a) CDF-S, (b) COSMOS, (c) ELAIS-S1, (d) S-CVZ, (e) SDSS Stripe 82, and (f) XMM-LSS fields. Pointings 1, 2, and 3 are indicated in black, blue, and red colors, respectively.

fully obtain zp for 99.9% of the CCD images, with DES-DR2 AUTO contributing to the vast majority of them (99.4%).

2.2. Photometry and Astrometry Cross-Calibration

Following the same recipe as in Section 2.1.1, we check the accuracy of the HELM pipeline photometry and astrometry independently for the fields that are not covered by the official DES data products (COSMOS and S-CVZ). The DES-DR2 catalog is based on coadded images, with a photometric accuracy of ~ 0.01 mag and astrometric precision of $\sim 0''.03$. We select all the point sources detected in our observations and crossmatch with the DES-DR2 catalog for all the reliably extracted ($\text{IMAFLAGS_ISO}=0$, $\text{FLAGS}<4$), high SNR (m_{PSF} error < 0.1 mag) sources. We select point sources from the DES-DR2 catalog using $0 \leq \text{EXTENDED_COADD} \leq 1$. For our observations, we use a relatively loose criterion of $\text{CLASS_STAR} > 0.8$, which ensures we do not miss a significant number of point sources and avoids including many extended sources.

Figure 5 presents an example of comparison of m_{PSF} and coordinate difference (Δd) as a function of m_{PSF} error in

pointing C1. The statistics for all 10 pointings are presented in Table 3. Although various degrees of differences are found in different fields, the source magnitude (median $\Delta m_{\text{PSF}} \approx 0.02$ mag) and astrometry ($\Delta d \approx 0''.06$) of our HELM pipeline are in good agreement with DES-DR2 for high SNR sources. The overall good performances in zp, photometry, and astrometry lend support to the reliability of the HELM pipeline for use in fields not covered by the footprint of DES-DR2 (COSMOS and S-CVZ).

3. DATA PRODUCTS

For this first HELM data release, we include the source catalog covering all compiled fields, as well as the light curves for these sources from the first 3.5 years (2019A-2022A). Our source catalog is built by cross-matching detected sources in our individual images with the DES-DR2 (Abbott et al. 2021) source catalog for the CDF-S, XMM-LSS, ELAIS-S1, and SDSS Stripe 82 fields, and with the source catalog from the third data release (Aihara et al. 2022) of the Hyper Suprime-Cam (HSC) Subaru Strategic Program (SSP; Aihara et al. 2018) for the COSMOS field. In total, there are 5,404,721, 4,714,784, and 4,059,093 sources with detections in at least 3, 5, and 10 exposures in any band, re-




Figure 2. Cadence of the HELM in different fields, with vertical red lines indicating individual exposures. Moon illumination (illuminated fraction of the moon) is indicated by the blue gradient (white=no moonlight; dark blue=maximum moonlight) in the background.




Figure 3. Histograms of objects used to calibrate the magnitude zeropoint for each CCD image of every exposure. Histogram for point sources using m_{PSF} is in red (PSF), while histogram for all sources using m_{AUTO} is in black.

spectively. Figure 6 shows histograms of inverse variance-weighted mean m_{AUTO} ($m_{\text{AUTO-w}}$; see Section 3.1 for definition) in the *gri* bands for objects in all 6 fields with detections in at least 5 individual exposures. The peak $m_{\text{AUTO-w}}$ is ~ 24.0 , 23.6 , and 23.4 mag for the *gri* bands, respectively, which are deeper compared to those of typical single exposures (Table 1) and representing the best seeing condi-




Figure 4. Comparison of magnitude zeropoints (zp) calibrated against the DES-DR2 catalog using point sources (PSF) and all sources (AUTO). Only CCD images with at least 10 sources used for calibration are included. Red vertical dashed lines indicate 16th, 50th, and 84th percentiles of the distribution, with statistics shown in the lower-left corner.

tion and longest exposure time. In future data releases, we will provide individual images, deeper photometric catalogs built from the coadded images, and extended light curves (see Section 4). Below we describe in detail our source catalog and light curve compilation.

Table 2. Comparison of HELM zp, m_{PSF} , and astrometry to those from DES-DR2

Pointing	Δz_p	$\Delta m_{\text{PSF}} [g]$	$\Delta m_{\text{PSF}} [r]$	$\Delta m_{\text{PSF}} [i]$	$\Delta m_{\text{PSF}} [z]$	$\Delta d [g]$	$\Delta d [r]$	$\Delta d [i]$	$\Delta d [z]$
	mag	(mag)	(mag)	(mag)	(mag)	($''$)	($''$)	($''$)	($''$)
C1	$-0.007^{+0.017}_{-0.032}$	$-0.019^{+0.038}_{-0.042}$	$-0.026^{+0.036}_{-0.042}$	$-0.011^{+0.026}_{-0.029}$	$-0.015^{+0.024}_{-0.029}$	$0.059^{+0.054}_{-0.032}$	$0.052^{+0.050}_{-0.028}$	$0.050^{+0.047}_{-0.027}$	$0.050^{+0.045}_{-0.027}$
C2	$-0.006^{+0.015}_{-0.024}$	$-0.015^{+0.035}_{-0.036}$	$-0.020^{+0.034}_{-0.039}$	$-0.013^{+0.031}_{-0.028}$	$-0.015^{+0.028}_{-0.028}$	$0.059^{+0.054}_{-0.031}$	$0.052^{+0.050}_{-0.028}$	$0.051^{+0.047}_{-0.027}$	$0.051^{+0.046}_{-0.027}$
C3	$-0.005^{+0.018}_{-0.031}$	$-0.016^{+0.036}_{-0.041}$	$-0.018^{+0.032}_{-0.041}$	$-0.012^{+0.028}_{-0.030}$	$-0.018^{+0.025}_{-0.030}$	$0.066^{+0.063}_{-0.036}$	$0.060^{+0.060}_{-0.033}$	$0.058^{+0.057}_{-0.032}$	$0.052^{+0.048}_{-0.028}$
E1	$0.004^{+0.029}_{-0.043}$	$-0.029^{+0.037}_{-0.042}$	$0.020^{+0.039}_{-0.042}$	$0.007^{+0.032}_{-0.033}$	$-0.004^{+0.028}_{-0.028}$	$0.075^{+0.072}_{-0.040}$	$0.068^{+0.066}_{-0.036}$	$0.065^{+0.061}_{-0.035}$	$0.053^{+0.048}_{-0.028}$
E2	$-0.014^{+0.024}_{-0.035}$	$-0.037^{+0.036}_{-0.039}$	$-0.005^{+0.040}_{-0.034}$	$-0.014^{+0.032}_{-0.034}$	$-0.011^{+0.026}_{-0.025}$	$0.073^{+0.069}_{-0.039}$	$0.067^{+0.065}_{-0.036}$	$0.064^{+0.061}_{-0.034}$	$0.054^{+0.048}_{-0.029}$
S1	$-0.008^{+0.044}_{-0.029}$	$-0.023^{+0.045}_{-0.041}$	$-0.009^{+0.047}_{-0.032}$	$0.028^{+0.051}_{-0.043}$	$0.009^{+0.037}_{-0.026}$	$0.066^{+0.063}_{-0.036}$	$0.058^{+0.061}_{-0.032}$	$0.057^{+0.058}_{-0.031}$	$0.050^{+0.047}_{-0.027}$
S2	$0.000^{+0.026}_{-0.024}$	$0.009^{+0.052}_{-0.053}$	$-0.010^{+0.038}_{-0.033}$	$-0.009^{+0.032}_{-0.044}$	$-0.013^{+0.030}_{-0.037}$	$0.066^{+0.064}_{-0.036}$	$0.058^{+0.062}_{-0.032}$	$0.058^{+0.059}_{-0.032}$	$0.052^{+0.049}_{-0.028}$
X1	$-0.002^{+0.020}_{-0.035}$	$-0.008^{+0.041}_{-0.048}$	$-0.015^{+0.039}_{-0.045}$	$-0.008^{+0.031}_{-0.035}$	$-0.012^{+0.030}_{-0.034}$	$0.068^{+0.066}_{-0.036}$	$0.062^{+0.064}_{-0.034}$	$0.061^{+0.059}_{-0.033}$	$0.055^{+0.051}_{-0.030}$
X2	$0.004^{+0.019}_{-0.033}$	$-0.005^{+0.044}_{-0.051}$	$-0.008^{+0.041}_{-0.044}$	$-0.009^{+0.032}_{-0.037}$	$-0.015^{+0.029}_{-0.035}$	$0.070^{+0.068}_{-0.038}$	$0.064^{+0.064}_{-0.035}$	$0.062^{+0.061}_{-0.034}$	$0.056^{+0.051}_{-0.030}$
X3	$0.003^{+0.019}_{-0.030}$	$-0.009^{+0.042}_{-0.047}$	$-0.006^{+0.038}_{-0.041}$	$-0.003^{+0.031}_{-0.033}$	$-0.011^{+0.028}_{-0.029}$	$0.070^{+0.066}_{-0.038}$	$0.062^{+0.063}_{-0.034}$	$0.060^{+0.059}_{-0.033}$	$0.053^{+0.049}_{-0.028}$

NOTE— $\Delta \equiv$ our DECam data – DES-DR2. Statistics are 50th percentile with upper (84th – 50th) and lower (16th – 50th) 1σ of the distribution as the superscript and subscript, respectively. Δd represents the angular distance difference between coordinates.

3.1. Source catalogs and light curves

The majority of our DECam pointings in the COSMOS field are covered by Deep- and UltraDeep-depth images of HSC SSP. For regions outside the Deep and UltraDeep coverage, we use instead the source catalog from Wide-depth images of HSC SSP. We extract the source catalog with i band Kron magnitude brighter than 25.5 mag from the Hyper Suprime-Cam Legacy Archive² (`pdr3_dud_rev` for the Deep and UltraDeep catalogs and `pdr3_wide` for the Wide catalog) using example script `PDR3 example 1a`. For the Wide catalog, we further apply the criteria presented in `PDR3 example 6` script to the i band to keep only “clean” sources. For all the other fields except S-CVZ (i.e., CDF-S, ELAIS-S1, Stripe 82, and XMM-LSS), we use DES-DR2 as the base source catalog for catalog construction, as it is deeper than individual exposures of our program. A radius of $0''.5$ is adopted for crossmatching.

For the S-CVZ field, we construct the source catalog from scratch as no deep optical catalog in this field is currently available. We first randomly pick an individual exposure catalog as the reference catalog and crossmatch other catalogs in the *griz* bands to it one by one using a matching radius of $0.5''$. We then iteratively build the preliminary master catalog by adding new sources whose separations from existing sources are larger than $0.5''$. The coordinates of the preliminary master catalog obtained after the first run are determined as the median value of those classified as the same source in individual catalogs. We then perform a second run by crossmatching individual exposure catalogs to the prelim-

inary master catalog and obtain a final catalog in the *griz* bands. Finally, we crossmatch the u band individual exposure catalogs with the *griz* catalog, as the u band depth is much shallower compared to that in the *griz* bands.

Table 3 presents the table format of our median source catalog, including source index, coordinate, magnitudes (PSF and AUTO), extraction flags, star-galaxy classifier (`spread_model` and `class_star`), and number of exposures. For magnitude, we also provide inverse variance weighted mean (`mag_w`), defined as

$$\text{mag_w} \equiv \frac{\sum(m_i/e_i^2)}{\sum(1/e_i^2)}, \quad (1)$$

where m_i and e_i represent an individual measurement and its error in the i th exposure, respectively. Its corresponding uncertainty is

$$\text{mag_w_err} = \frac{\sqrt{\sum(e_i/e_i^2)^2}}{\sum(1/e_i^2)} = \frac{1}{\sqrt{\sum(1/e_i^2)}}, \quad (2)$$

after error propagation. For the S-CVZ field, we provide separate source indices and coordinates for *griz* and u bands, as the u band is much shallower compared to the other bands. Separation between the median coordinate in the *griz* bands and that in the u band is also provided. Table 4 describes the format of the light curve files of individual objects. The light curves of individual sources can be extracted from our online database³ using source indices provided in Table 3.

3.2. Example light curves

² <https://hsc.la.mtk.nao.ac.jp/doc/>

³ <https://ariel.astro.illinois.edu/helm/>




Figure 5. Differences in m_{PSF} (top) and coordinates (middle) between our measurements of all observations and those from the DES-DR2 catalog ($\Delta \equiv \text{HELM} - \text{DES-DR2}$) as a function of m_{PSF} error for all high SNR (m_{PSF} error < 0.1 mag) point sources in pointing C1. Normalized source number density (N/N_{max}) of m_{PSF} error is shown in the bottom panel. m_{PSF} error is the propagated error by combining our measurement error and DES-DR2 measurement error in quadrature sum. Median values in the $griz$ filters are shown as blue, green, orange, and red solid curves, respectively. Dashed curves in top and middle panels indicate values at 84th and 16th percentiles. Statistics of all 10 pointings are shown in Table 2.




Figure 6. Histograms of inverse variance-weighted mean m_{AUTO_w} in all detected exposures (m_{AUTO_w} ; see Equation 1) in g (blue), r (green), and i (orange) bands for objects in all 6 fields with detection in at least 5 individual exposures.

Figure 7 shows example light curves of three RR Lyrae stars and three quasars in the COSMOS field. For RR Lyrae stars, we have modeled their light curves using `astropy` and presented phase-folded light curves. The high-quality photometry and dense time sampling ensure excellent recovery of period information. The overall clean phase-folded light curves of RR Lyrae stars also demonstrate the good performance of our zp calibration (Section 2.2). The long time baseline, decent ~ 3 -day cadence, and multi-band coverage make it possible to qualitatively investigate both long and short timescale, as well as multi-band AGN variability. In particular, as shown in Figure 8 and its inset, the epoch magnitude uncertainty is ~ 5 times smaller in HELM than in the ZTF DR19 – median uncertainty of 0.014 mag from HELM r band versus 0.068 mag from ZTF r band. Therefore the resulting optical light curves from HELM are of much better quality than ZTF light curves, albeit with sparser cadences.

3.3. Caveats

In this paper, our source catalogs for the CDF-S, ELAIS-S1, Stripe 82, and XMM-LSS fields are constructed based on the DES-DR2 source catalog, which has a median coadded catalog depth for a $1''.95$ diameter aperture at SNR=10 of $g=24.7$, $r=24.4$, $i=23.8$, and $z=23.1$ mag. However, we may miss transients that were not detected in previous surveys. Given the number of exposures (up to 387 s) and total exposure time (up to 4480 s) in these fields, together with on-going exposures that are not included in this data release, we would eventually achieve much deeper source detection by coadding all individual exposures. For the S-CVZ field, our source catalog is constructed based on catalogs of individual exposures. Therefore, the depth is limited by the deepest exposure. For the COSMOS field, we notice a source splitting issue for saturated objects in the HSC catalog. As we are using deeper catalogs for source cross-matching (in particular for the COSMOS field), our source catalog may contain some spurious sources with low N_{exp} and close to or even below the typical depth of DECam individual exposures (relatively steep increase of source number toward the faint end in the g and i bands as seen in Figure 6). Moreover, as the saturation magnitudes are quite faint for the HSC survey (18.0, 18.2, and 18.6 mag for the gri bands; Aihara et al. 2018), this issue may affect the sources that are above their saturation limits. Another major caveat is that there is no forced photometry or upper limits, which may give a false sense of the overall variability especially for sources close to the detection limit. The caveats mentioned above would be largely mitigated by building source catalogs directly from coadded DECam images and performing forced photometry on each source in our future data releases. On the other hand, any variable / transient sources that are not associated with a static object in the matched catalogs will be missed (e.g., off-axis transients and




Figure 7. Example light curves of RR Lyrae stars (left; in folded phase) and AGNs (right) in the COSMOS field. Blue, green, and red dots represent the g , r , and i bands, respectively, with error bar indicating the certainty. The periods of RR Lyrae stars are shown at the top of the left panels.




Figure 8. Comparison of DECAM r band (open blue circle) and ZTF r band (open red square) light curves of a quasar in the COSMOS field. The inset shows a zoom-in view around MJD=59300 day. The error bars indicate 1σ uncertainty.

moving objects). These objects will be included in our future data releases after implementing dedicated pipeline.

4. SUMMARY

In this work we present an overview of the HELM long-term monitoring program in extragalactic legacy fields, with the DECAM imager on the 4m CTIO Blanco telescope. With a multi-year baseline and relatively high cadence, HELM aims to provide high-quality optical light curves for a large sample of AGNs in these fields, as part of the effort to ob-

tain RM measurements for distant AGNs and quasars in the SDSS-V BHM-RM program. This long-term photometric monitoring program will also bridge past light curves and future LSST/WFST observations to compile decades-long light curves to facilitate investigations of long-term variability of astronomical objects (particularly AGNs). Given the ample multi-wavelength data in these legacy fields, the HELM light curve data will also provide legacy value to time-domain studies. At the time of writing, HELM has been extended to

continue DECam monitoring in these fields through at least 2025B, given the delayed start of the LSST.

In this work we also present the first data release from HELM, which includes source catalogs and light curves from the first 3.5 years of observations. In future data releases, we will provide fully calibrated individual and coadded images, as well as extended light curves.

1 We thank Tamara M. Davis for initial discussions. This work
2 is supported by NSF grants AST-2206499 and AST-2308077.
3 F.E.B. acknowledges support from ANID - Millennium
4 Science Initiative Program - ICN12.009, CATA-BASAL -
5 FB210003, and FONDECYT Regular - 1200495. Y.Q.X. ac-
6 knowledges support from the National Key R&D Program
7 of China (2023YFA1608100), NSFC grants (12025303,
8 12393814), and the Strategic Priority Research Program of
9 the Chinese Academy of Sciences (XDB0550300).

Facilities: CTIO Blanco telescope

Software: `astropy` (Astropy Collaboration et al. 2013, 2018), `matplotlib` (Hunter 2007), `Numpy` (Harris et al. 2020), `SExtractor` (Bertin & Arnouts 1996)

REFERENCES

- Abbott, T. M. C., Adamów, M., Aguena, M., et al. 2021, *ApJS*, 255, 20, doi: [10.3847/1538-4365/ac00b3](https://doi.org/10.3847/1538-4365/ac00b3)
- Aihara, H., Arimoto, N., Armstrong, R., et al. 2018, *PASJ*, 70, S4, doi: [10.1093/pasj/psx066](https://doi.org/10.1093/pasj/psx066)
- Aihara, H., AlSayyad, Y., Ando, M., et al. 2022, *PASJ*, 74, 247, doi: [10.1093/pasj/psab122](https://doi.org/10.1093/pasj/psab122)
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33, doi: [10.1051/0004-6361/201322068](https://doi.org/10.1051/0004-6361/201322068)
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, *AJ*, 156, 123, doi: [10.3847/1538-3881/aabc4f](https://doi.org/10.3847/1538-3881/aabc4f)
- Bellm, E. C., Kulkarni, S. R., Graham, M. J., et al. 2019, *PASP*, 131, 018002, doi: [10.1088/1538-3873/aaecbe](https://doi.org/10.1088/1538-3873/aaecbe)
- Bernstein, J. P., Kessler, R., Kuhlmann, S., et al. 2012, *ApJ*, 753, 152, doi: [10.1088/0004-637X/753/2/152](https://doi.org/10.1088/0004-637X/753/2/152)
- Bertin, E., & Arnouts, S. 1996, *A&AS*, 117, 393, doi: [10.1051/aas:1996164](https://doi.org/10.1051/aas:1996164)
- Blandford, R. D., & McKee, C. F. 1982, *ApJ*, 255, 419, doi: [10.1086/159843](https://doi.org/10.1086/159843)
- Chambers, K. C., Magnier, E. A., Metcalfe, N., et al. 2016, *arXiv e-prints*, arXiv:1612.05560, doi: [10.48550/arXiv.1612.05560](https://doi.org/10.48550/arXiv.1612.05560)
- Drlica-Wagner, A., Ferguson, P. S., Adamów, M., et al. 2022, *ApJS*, 261, 38, doi: [10.3847/1538-4365/ac78eb](https://doi.org/10.3847/1538-4365/ac78eb)
- Flaugher, B., Diehl, H. T., Honscheid, K., et al. 2015, *AJ*, 150, 150, doi: [10.1088/0004-6256/150/5/150](https://doi.org/10.1088/0004-6256/150/5/150)
- Graham, M. J., Djorgovski, S. G., Stern, D., et al. 2015, *MNRAS*, 453, 1562, doi: [10.1093/mnras/stv1726](https://doi.org/10.1093/mnras/stv1726)
- Graham, M. L., Knop, R. A., Kennedy, T. D., et al. 2023, *MNRAS*, 519, 3881, doi: [10.1093/mnras/stac3363](https://doi.org/10.1093/mnras/stac3363)
- Harris, C. R., Millman, K. J., van der Walt, S. J., et al. 2020, *Nature*, 585, 357, doi: [10.1038/s41586-020-2649-2](https://doi.org/10.1038/s41586-020-2649-2)
- Henden, A. A., Levine, S., Terrell, D., & Welch, D. L. 2015, in *American Astronomical Society Meeting Abstracts*, Vol. 225, American Astronomical Society Meeting Abstracts #225, 336.16
- Hunter, J. D. 2007, *Computing in Science & Engineering*, 9, 90, doi: [10.1109/MCSE.2007.55](https://doi.org/10.1109/MCSE.2007.55)
- Jones, D. O., Scolnic, D. M., Riess, A. G., et al. 2017, *ApJ*, 843, 6, doi: [10.3847/1538-4357/aa767b](https://doi.org/10.3847/1538-4357/aa767b)
- Kollmeier, J. A., Zasowski, G., Rix, H.-W., et al. 2017, *ArXiv e-prints*. <https://arxiv.org/abs/1711.03234>
- Lacy, M., Surace, J. A., Farrah, D., et al. 2021, *MNRAS*, 501, 892, doi: [10.1093/mnras/staa3714](https://doi.org/10.1093/mnras/staa3714)
- Morganson, E., Gruendl, R. A., Menanteau, F., et al. 2018, *PASP*, 130, 074501, doi: [10.1088/1538-3873/aab4ef](https://doi.org/10.1088/1538-3873/aab4ef)
- Peterson, B. M. 1993, *PASP*, 105, 247, doi: [10.1086/133140](https://doi.org/10.1086/133140)
- Sako, M., Bassett, B., Becker, A. C., et al. 2018, *PASP*, 130, 064002, doi: [10.1088/1538-3873/aab4e0](https://doi.org/10.1088/1538-3873/aab4e0)
- Shen, Y., Brandt, W. N., Dawson, K. S., et al. 2015, *ApJS*, 216, 4, doi: [10.1088/0067-0049/216/1/4](https://doi.org/10.1088/0067-0049/216/1/4)
- Smith, M., D’Andrea, C. B., Sullivan, M., et al. 2020, *AJ*, 160, 267, doi: [10.3847/1538-3881/abc01b](https://doi.org/10.3847/1538-3881/abc01b)
- Wang, T., Liu, G., Cai, Z., et al. 2023, *Science China Physics, Mechanics, and Astronomy*, 66, 109512, doi: [10.1007/s11433-023-2197-5](https://doi.org/10.1007/s11433-023-2197-5)
- York, D. G., Adelman, J., Anderson, John E., J., et al. 2000, *AJ*, 120, 1579, doi: [10.1086/301513](https://doi.org/10.1086/301513)
- Zou, F., Brandt, W. N., Chen, C.-T., et al. 2022, *ApJS*, 262, 15, doi: [10.3847/1538-4365/ac7bdf](https://doi.org/10.3847/1538-4365/ac7bdf)

Table 3. FITS table format for the median source catalog

Column Name	Format	Units	Description
Index	LONG		Source index
ra	DOUBLE	degree	Right Ascension (J2000)
dec	DOUBLE	degree	Declination (J2000)
mag_psf	DOUBLE	mag	SExtractor PSF magnitude
mag_auto	DOUBLE	mag	SExtractor AUTO magnitude
flags	LONG		SExtractor extraction flags
spread_model	DOUBLE		SExtractor model-based star-galaxy classifier
class_star	DOUBLE		SExtractor neural network-based star-galaxy classifier
Nexp_psf	LONG		Number of exposures with successful mag_psf measurement
Nexp_auto	LONG		Number of exposures with successful mag_auto measurement
Separation	DOUBLE	arcsec	Distance between median coordinate in <i>griz</i> bands and <i>u</i> band (only in S-CVZ field)

NOTE—All columns except Separation can have prefix, while all columns except Index, Nexp_psf, Nexp_auto, and Separation have suffix. Prefix: “*_” indicates the associated band for the parameter, with * = one or a combination of *ugriz* bands (such as *g_* in all fields or *griz_* only for S-CVZ). If not specified (for Index, ra, and dec in all fields except S-CVZ), it represents the value for all bands combined. Suffix: “_m” and “_err” indicate unweighted median (50%) and its corresponding error of Nexp measurements with *_err* = $0.5 \times (84\text{th percentile} - 16\text{th percentile})/\sqrt{N_{\text{exp}}}$; “_w” indicates inverse variance weighted mean ($\Sigma(m_i/e_i^2)/\Sigma(1/e_i^2)$, where m_i and e_i represent individual measurement and its error at *i*th exposure, respectively); “_w_err” indicates the error associated with *_w* after error propagation ($1/\sqrt{\Sigma(1/e_i^2)}$). Specifically, “_err” = “_w_err” = error of individual measurement when Nexp=1. Value = “-99” indicates no data or non-detection.

Table 4. FITS table format for the light curve

Column Name	Format	Units	Description
ra	DOUBLE	degree	Right Ascension (J2000)
dec	DOUBLE	degree	Declination (J2000)
exnum	LONG		DECam exposure number
mjd	DOUBLE	day	Modified Julian date of exposure
band	Char		Filter (one of <i>ugriz</i>)
mag_psf	DOUBLE	mag	SExtractor PSF magnitude
mag_e_psf	DOUBLE	mag	Uncertainty of SExtractor PSF magnitude
mag_auto	DOUBLE	mag	SExtractor AUTO magnitude
mag_e_auto	DOUBLE	mag	Uncertainty of SExtractor AUTO magnitude
flags	LONG		SExtractor extraction flags
spread_model	DOUBLE		SExtractor model-based star-galaxy separation
spread_model.e	DOUBLE		Uncertainty of SExtractor model-based star-galaxy separation
class_star	DOUBLE		SExtractor neural network-based star-galaxy separation