



SMALL ANGLE ELASTIC PROTON-PROTON SCATTERING  
FROM 25 TO 200 GeV

V. Bartenev, R. Carrigan, I-Hung Chiang, R. L. Cool,  
K. Goulianos, D. Gross, A. Kuznetsov,  
E. Malamud, A. C. Melissinos, B. Morozov,  
V. Nikitin, S. L. Olsen, Y. Pilipenko,  
V. Popov, R. Yamada, and L. Zolin

October 1972



SMALL ANGLE ELASTIC PROTON-PROTON SCATTERING FROM 25 TO 200 GeV

V. Bartenev, R. Carrigan, I-Hung Chiang, R. L. Cool,  
K. Goulianos, D. Gross, A. Kuznetsov,  
E. Malamud, A. C. Melissinos, B. Morozov,  
V. Nikitin, S. L. Olsen, Y. Pilipenko,  
V. Popov, R. Yamada, and L. Zolin

The State Committee for Utilization of Atomic Energy of the USSR, Moscow

National Accelerator Laboratory, Batavia, Illinois

Rockefeller University, New York, New York<sup>†</sup>

University of Rochester, Rochester, New York<sup>‡</sup>

ABSTRACT

We have measured the differential cross section for small angle p-p scattering from 25 to 200 GeV incident energy and in the momentum transfer range  $0.015 < |t| < 0.080 \text{ (GeV/c)}^2$ . We find that the slope of the forward diffraction peak,  $b(s)$ , increases with energy and can be fit by the form  $b(s) = b_0 + 2 \alpha' \ln s$  where  $b_0 = 8.3 \pm 1.3$  and  $\alpha' = 0.28 \pm 0.13 \text{ (GeV/c)}^{-2}$ . Such dependence is compatible with the data existing both at higher and lower energies. We have also obtained the energy dependence of the p-p total cross section in the energy range from 48 to 196 GeV. Within our errors which are  $\pm 1.1 \text{ mb}$  the total cross section remains constant.

---

<sup>†</sup> Work supported in part by the U.S. Atomic Energy Commission under Contract Number AT(11-1)-2232.

<sup>‡</sup> Work supported in part by the U.S. Atomic Energy Commission under Contract Number AT(11-1)-3065.

We have measured the differential cross section for p-p elastic scattering at small angles for incident energies from 25 to 200 GeV, in the momentum transfer range from 0.015 to 0.080 (GeV/c)<sup>2</sup>. From the data we have determined the slope of the forward diffraction peak and the energy dependence of the elastic differential cross section extrapolated to  $t = 0$ .

The experimental method<sup>(1)</sup> makes use of the fact that the kinetic energy,  $T$ , of the recoil proton from elastic p-p scattering is directly related to the momentum transfer  $|t|$ , through  $|t| = 2mT$  where  $m$  is the mass of the proton. Furthermore, for a fixed value of  $|t|$  the angle of emission of the recoil proton is almost independent of the incident beam energy. We have measured the energy and angle of the recoil protons by using lithium drifted silicon solid state detectors. Eight such detectors were placed so as to cover the angular range from  $90^\circ$  to  $80^\circ$  in the laboratory. The detectors were collimated to an area of  $0.6 \text{ cm}^2$  and were located 3.5m from the scattering target. The energy response of the detection system was linear and had a resolution of 0.5%. The energy calibration was continuously monitored using 5.47 MeV  $\alpha$ -particles from an Americium source.<sup>(2)</sup>

The measurements were performed at the National Accelerator Laboratory using a polyethylene target in the internal beam of the accelerator. The target consisted of a thin (typically 3 microns) strip of polyethylene foil which rotated at 60 cps through the internal beam. The axis of rotation was oriented at an angle of  $20^\circ$  from the beam direction.<sup>(3)</sup> The tip of the foil was at a radius of 62 mm from the axis, and the foil was 6 mm wide at the tip. The foil was cut radially so as to render the interaction rate independent of beam position. The thickness of the foil was uniform to  $\pm 4$  percent.

The detectors were mounted on a carriage so that their angular position

could be changed. Data were obtained at six different carriage positions which provides (1) for cross-calibration of detector acceptance and efficiency, and (2) for a measurement of the shape of the background under the peaks. It was observed that the background is independent of detector position to a high degree of accuracy. One of the detectors was kept at a fixed angle to serve as a normalization monitor. The beam intensity was also monitored using a toroidal coil encircling the beam pipe.<sup>(4)</sup> The signals from each detector were routed through appropriate circuitry to a single ADC and then written on magnetic tape. The signals were gated in synchronism with the foil passing through the beam; an asynchronous gate provided simultaneous target out data. Each event recorded on tape contained information on detector position, the recoil proton energy, the primary beam energy and the target status. In addition, events occurring during the ADC readout time were counted and recorded for use in correcting for dead time losses.

Clear peaks corresponding to elastic scattering were observed in the energy spectrum of each detector. At low values of  $|t|$  the peak is due to protons stopping in the detector whereas at high  $|t|$ -values the peaks were due to the energy loss ( $\int dE/dx$ ) of protons traversing the detector. The elastic peaks were superimposed on a background arising mainly from the proton-carbon interactions in the target. To extract the yield of elastic events a background envelope, obtained from several positions of the same detector (but separately for each given energy), was subtracted from the data histogram. That this procedure selects only elastic events from p-p interactions in the target, was verified by comparing data for the same  $|t|$ -value (and same energy) but obtained from different detectors as their angular position was varied. Finally, the elastic yield was corrected for the following effects: (a) Nuclear interactions in the detector;<sup>5</sup> 0.2 to

2.6 percent. (b) Interference of the Coulomb and nuclear amplitudes; 1.7 to 7.0 per cent.<sup>6</sup> (c) Geometric acceptance correction; 2.5 to 12.3 per cent.

The differential cross sections extracted from the data in the interval  $0.015 < |t| < 0.080$  were fitted to a functional form

$$\frac{d\sigma}{dt}(s,t) = \frac{d\sigma}{dt}(s,t=0) e^{-b(s)|t|} \quad (1)$$

and the slope parameter,  $b(s)$ , was determined for eight different energy intervals. The width of the energy bins was 10 GeV centered at 25 GeV intervals. Data from all detectors and all positions were combined, with appropriate errors to account for the statistical fluctuations of the peak and background. Typically the fits gave  $\chi^2/\text{degree of freedom}$  in the range 0.7 - 1.5; the error on  $b(s)$  was incremented by  $\sqrt{\chi^2/\text{d.f.}}$  when the  $\chi^2$  exceeded 1.0. The differential cross sections are shown in Fig. 1 and contain approximately 60,000 elastic events at each energy.

The values of  $b(s)$  obtained are shown in Fig. 2 where data from other experiments are also included.<sup>1,7,8,9,10</sup> Our data are, in general, in agreement with that of the other authors. If  $b(s)$  is parametrized in the usual way as

$$b(s) = b_0 + a \alpha' \ln s$$

we find, using only our data

$$b_0 = 8.3 \pm 1.3 \quad \alpha' = 0.28 \pm 0.13 \quad (\text{GeV}/c)^{-2}$$

We refit the data to Eq. (1) using the best fit values of  $b(s)$ , to determine  $d\sigma/dt(s, t=0)$ . We note that the forward cross section depends only weakly on  $b(s)$ ; a 10 per cent variation in  $b(s)$  results in a 3 per cent change in  $d\sigma/dt(s, t=0)$ . The data were examined for possible variations with target position, target deterioration, and beam intensity; within statistical uncertainties no effects of this kind were observed. We have scaled the

statistical error by a factor of two to allow for possible systematic effects.

Using the optical theorem and the usual assumption about spin-independence of the p-p interaction, the forward differential cross section can be related to the total cross section

$$\sigma_T(s) = \frac{d\sigma}{dt}(s, t = 0) \frac{4\sqrt{\pi}}{(1 + \alpha^2/2)}$$

Our data on the energy dependence of  $\sigma_T$  are presented in Table I where the values of  $\alpha = \text{Re}(f)/\text{Im}(f)$  are taken from Ref. 6. Finally, if the total cross section at  $E = 48$  GeV is normalized to the value  $\sigma_T = 38.5$  mb given by Denisov et al.,<sup>11</sup> we obtain the p-p total cross section at the remaining six energies.<sup>12</sup> The error on each point is  $\pm 1.1$  mb and within these errors the total cross section remains constant in the energy interval from 48 to 196 GeV. This observation is in agreement with total cross section data reported by Holder et al.<sup>13</sup> at an equivalent energy of  $E = 500$  GeV and by Charlton et al.<sup>14</sup> at  $E = 200$  GeV.

We are indebted to many individuals who have contributed at various stages of this experiment, and in particular we wish to acknowledge the assistance and support given us by the NAL accelerator operations staff.

TABLE I

Energy Dependence of p-p Total Cross Section

| $E_{\text{lab}}$<br>(GeV) (a) | $b(s)$<br>(GeV/c) <sup>-2</sup> (b) | $\alpha = \frac{\text{Re}A}{\text{Im}A}$ (c) | $\left. \frac{d\sigma}{dt} \right _{t=0}$<br>Relative units (d) | $\sqrt{\frac{d\sigma}{dt}} / (1+\alpha^2/2)$<br>Relative units | $\sigma_{\text{total}}$ (e) |
|-------------------------------|-------------------------------------|--|---|--|-----------------------------|
| 48                            | 10.9                                | - 0.16                                       | (1.03)  | (1.00)   | (38.5 ± 0.1)                |
| 73                            | 11.1                                | - 0.09                                       | 0.96 ± 0.06   | 0.98 ± 0.03  | 37.7 ± 1.1                  |
| 98                            | 11.3                                | - 0.09                                       | 0.96 ± 0.06   | 0.98 ± 0.03  | 37.7 ± 1.1                  |
| 122                           | 11.5                                | - 0.09                                       | 0.93 ± 0.06   | 0.96 ± 0.03  | 37.0 ± 1.1                  |
| 147                           | 11.6                                | - 0.09                                       | 0.96 ± 0.06   | 0.98 ± 0.03  | 37.6 ± 1.1                  |
| 172                           | 11.7                                | - 0.09                                       | 0.95 ± 0.06   | 0.97 ± 0.03  | 37.4 ± 1.1                  |
| 196                           | 11.7                                | - 0.09                                       | 1.01 ± 0.06   | 1.00 ± 0.03  | 38.5 ± 1.2                  |

(a) The energy bins are 10 GeV wide centered at the indicated energies.

(b) The slope parameters are obtained from the fit shown in Fig. 2.

(c) The values of  $\alpha$  are taken from Ref. 6 for the lowest energy and assumed to be -0.09 for  $E \geq 73$  GeV. Note that if  $\alpha = 0$ , the change in  $\sigma_T$  is 0.5 per cent.

(d) The forward cross section is given in relative units with the 48 GeV point having been set equal to 1.03.

(e) The total cross section at 48 GeV was normalized to the point given by Denisov et al. (Ref. 11).

References and Notes

1. This technique was first successfully used by G. Beznogikh et al. Phys. Letters 30B, 274 (1969).
2. The detectors were also calibrated using proton beams from the Princeton Cyclotron. We are indebted to E. Cecil for his assistance in these runs.
3. We wish to thank Messrs. T. Haelan and C. Wilson for the construction of the target and vacuum chamber.
4. We thank Dr. D. Sutter for designing and providing the beam intensity monitor. This instrument had a time constant of 500  $\mu$ sec. and its linearity and stability were better than 1 per cent at a circulating beam current of 5 mA.
5. The correction for nuclear interactions in the silicon detectors was taken from report CERN 69-17 by D. F. Measday and C. Richard-Serre.
6. To perform the correction for the contribution of Coulomb interference to the nuclear scattering, the data of G. Beznogikh et al., Phys. Letters 39B, 411 (1972) were used and extrapolated up to 200 GeV/c.
7. G. Barbiellini et al., Phys. Letters 39B, 663 (1972).
8. U. Amaldi et al., Phys. Letters 36B, 504 (1971).
9. G. Bellettini et al., Phys. Letters 14, 164 (1965).
10. K. Chernev et al., Phys. Letters 36B, 266 (1971).
11. S. Denisov et al., Phys. Letters 36B, 415 (1971).
12. The internal beam intensity varied considerably in the first energy bin (centered at 23 GeV); therefore, this point has been excluded from the data.
13. M. Holder et al., Phys. Letters 35B, 361 (1971). These authors quote  $\sigma_T = 40.3 \pm 2$  mb at a center of mass energy of 30 GeV.



14. G. Charlton et al., Phys. Rev. Letters 29, 515 (1972).

Figure Captions

Figure 1. The elastic p-p differential cross section  $d\sigma/dt$  at the eight measured energy intervals. The data points have been corrected as discussed in the text. The normalization of the data is based on the value of  $\sigma_T$  at an energy of 48 GeV given in reference (12).

Figure 2. The slope of the diffraction peak,  $b(s)$  as a function of the square of the c.m. energy. The straight line fit is made only to the eight points measured in this experiment.



