

A NOTE ON PHASE 1  $\mu e$  SCATTERING AND  
MUON BREMSSTRAHLUNG EXPERIMENTS AT WESTON

W. A. Blanpied  
Case Western Reserve University

May 1, 1968

In a memorandum dated April 1968, Richard Wilson has outlined the essential features of a facility at Weston that could be used for many different experiments with a 100 GeV muon beam, and in particular discusses (1) measuring the total cross section for interactions of virtual photons with nucleons by means of muon inelastic scattering, and (2) the production of  $\rho$ -mesons by virtual photons. The proposed facility extrapolates from the experiments performed by Perl, et al, at SLAC, and by the Case-Harvard-McGill-Stanford collaboration at the AGS. In addition to a 1 meter hydrogen target it would include magnets to analyze the momenta of the incident muons, the inelastically scattered muon and other final state charged particles, trigger counters and wire plane spark chambers, and large total absorption photon counters with a few percent resolution such as have been studied by Hofstadter and others.

Two major sources of background triggers in any hadron production experiment using this facility would be (1)  $\mu e$  scattering events (knock-on electrons), and (2) muon bremsstrahlung events ( $\mu + N \rightarrow \mu' + N + \gamma$ ). Of course both of these processes are of considerable

interest in their own right. It is obvious that since they are important sources of background in hadron measurements, the proposed facility could also be modified to study both of them, though probably not simultaneously, since the knock-on cross section is orders of magnitude larger than the bremsstrahlung cross section. Both experiments are in the Phase I category in the sense that they could probably be performed prior to installation of the complete facility. In fact, an interesting region of the bremsstrahlung spectrum could be studied using conventional magnets.

In the muon inelastic scattering experiment at the AGS, knock-on triggers were largely eliminated by the kinematic requirement that an incident, 12 GeV muon lose at least  $3\frac{1}{2}$  GeV by interacting in the target before triggering the system. Except for very high energy loss processes, the kinematic elimination of knock-on triggers will not be feasible at Weston, since a 100 GeV muon can lose more than 80% of its energy to an electron. Therefore, as Wilson points out, a non-coplanarity requirement will have to be included in the trigger for any hadron production experiment. Conversely, a coplanarity requirement could be used to insure that most triggers resulted from scattering events. If the incident muon momentum, the opening angle and momenta of the final state muon and electron were measured, and the electron and muon identified as such, the kinematics would be

overdetermined. Therefore, it appears that  $\mu e$  scattering could be studied even if the incident muon energy were not very well known.

The existence of total absorption photon counters with energy resolution on the order of 5% would imply the possibility of carrying out muon bremsstrahlung measurements without measuring the incident muon momentum too well. In this case the trigger would include a maximum two-charged particle requirement (the inelastic muon and, for high momentum transfer events, the recoil proton). Measurement of the  $\mu\gamma$  opening angle and the energies of both final state particles would then determine the kinematics for any "elastic" event, i. e., events with no other particles save the recoil proton in the final state.

The major advantage in planning experiments that do not require momentum analysis of both the incoming and outgoing particles is obvious. Large superconducting magnets are going to be required in the facility. (For example, a 72 inch, 100 kilogauss dipole gives  $\frac{\Delta\theta}{\Delta p} = 6.4 \times 10^{-4}$  rad/GeV for a 90 GeV particle and  $2.2 \times 10^{-3}$  rad/GeV at 50 GeV.) Clearly, it will be easier to obtain one magnet of this description than two during Phase I operations at Weston. Certainly their design and construction should be pushed as soon as possible so that in fact the full facility can be set up soon after the muon beam is installed. However, performing the  $\mu e$  and/or the bremsstrahlung experiments at an even earlier stage would provide

an opportunity to study the 100 GeV,  $10^5/\text{sec}$ , muon beam we hope will exist soon after the accelerator is available to experimenters, as well as to do some worthwhile physics.

It is worth noting that an experiment to measure the very high energy part of the muon bremsstrahlung spectrum might well be performed even before high field magnets are available, since momentum analysis of the accompanying 10-15 GeV final state muons could be performed using more conventional low field magnets. In addition, the high trigger background in other parts of the spectrum could be largely eliminated by imposing a kinematic cutoff on the scattered muon (as in the AGS experiment) in a high energy loss, wide angle region. In fact, the most interesting physics is here, since these high energy loss events offer the most sensitive probe of the muon propagator. If one superconducting magnet were available its most advantageous role would be to analyze the incident muon momentum and thus to overdetermine the kinematics.

Low counting rates might prove to be the major reason for postponing the bremsstrahlung experiment beyond Phase I. Assume  $10^5$  100 GeV muons incident per second upon a 1 meter long liquid hydrogen target, and consider final photons with energies in the range  $90 \pm 2$  GeV and angles within  $\pm 0.2^\circ$  of the forward direction.

Then the Bethe-Heitler differential equation for a point proton gives 1 or 2 per hour for scattered muons between  $2^\circ$  and  $3^\circ$ . This angular range is quite interesting, corresponding to effective masses of the virtual muon ranging from 1.0 to 1.6 GeV.\* Rates for smaller scattering angles are expected to be much higher -- a few per minute in the range from  $1^\circ$  to  $2^\circ$ . However, there is considerably less physics interest here, since the corresponding effective muon masses are smaller than 1 GeV.

---

\*For one of the two dominant diagrams contributing to the process. The effective mass of the other is on the order of  $M^2 = (-200 \text{ GeV})^2$ .