


Jefferson Laboratory Overview



Anthony W Thomas

Neutrino Meeting: May 4th 2006

Thomas Jefferson National Accelerator Facility



U.S. DEPARTMENT OF ENERGY
S&T, September 11, 2005 1


JLab: Unique Forefront Capabilities for Science




Cryomodules in the accelerator tunnel



Superconducting radiofrequency (SRF) cavities undergo vertical testing.



An aerial view of the recirculating linear accelerator and 3 experimental halls.




CEBAF Large Acceptance Spectrometer (CLAS) in Hall B



JLab is Central to Nuclear Science

Quark-Gluon Structure Of Nucleons and Nuclei

Nature of Confinement



Correlations
n-radii: $N \neq Z$
Hypernuclei

Hadrons in- medium
Effective NN (+ HN) force


Precise
few-nucleon
calculations

Exotic mesons
and baryons

12 GeV Upgrade Project

- Status
 - CD-1 approved!!!
 - R&D providing valuable information
 - ACD effort on track for CD-2A mid-year
- Planning
 - Project is supported in all funding scenarios
 - CD-1 approval opens the door for getting firm commitments of resources from non-DOE funding sources
 - February '06 project funding guidance:
 - Reduced construction funds in FY08 and FY09, increased in FY10 and FY11
 - Best current guess: 6 GeV program to end CY10

CEBAF at 12 GeV



Highlights of the 12 GeV Program

- **Revolutionize Our Knowledge of Spin and Flavor Dependence of Valence PDFs**
- **Revolutionize Our Knowledge of Distribution of Charge and Current in the Nucleon**
- **Totally New View of Hadron (and Nuclear) Structure: GPDs**
 - **Determination of the quark angular momentum**




Highlights of the 12 GeV Program....2

- **Exploration of QCD in the Nonperturbative Regime:**
 - **Existence and properties of exotic mesons**
- **New Paradigm for Nuclear Physics:**
Nuclear Structure in Terms of QCD
 - **Spin and flavor dependent EMC Effect**
 - **Study quark propagation through nuclear matter**
- **Precision Tests of the Standard Model**
 - **Factor 20 improvement in $(2C_{2u}-C_{2d})$**



Distribution of Charge and Current in the Nucleon



HP 2010

Apparatus almost complete: new scattering Chamber, 1700 block calorimeter, new focal plane polarimeter


- Perdrisat *et al.* E01-109 — will increase range of Q^2 by 50% in 2007 (range of Q^2 for n will double over next 3-4 years)
- With 12 GeV and SHMS in Hall C

JLab data on the EM form factors provide a testing ground for theories constructing nucleons from quarks and glue


Planned Extensions w/ 6 GeV beams

Electric


Proton




Neutron



Magnetic



Selected World Data





Thomas Jefferson National Accelerator Facility



Two-Photon Exchange Experiment

- Measurement of proton *electric* form factor differs by factor of 4 in two different measurements
- Two-photon exchange is only known explanation



- Unambiguous determination of this process can be made by comparing positron-proton to electron-proton elastic scattering

- CLAS experiment can determine this with 1% systematic error

Strangeness Widely Believed to Play a Major Role – Does It?

- As much as 100 to 300 MeV of proton mass??

$$M_N = \langle N(P) | -\frac{9\alpha_s}{4\pi} \text{Tr}(G_{\mu\nu}G^{\mu\nu}) + m_u \bar{\psi}_u \psi_u + m_d \bar{\psi}_d \psi_d + m_s \bar{\psi}_s \psi_s | N(P) \rangle$$


$$\Delta M_N^{s\text{-quarks}} = \frac{ym_s}{m_u + m_d} \sigma_N$$

Through proton spin crisis:

- as much as 10% of the spin of the proton??

HOW MUCH OF THE ELECTRIC & MAGNETIC FORM FACTORS ?

Strange Quark Form Factors at $Q^2 = 0.1$ GeV²



$$G_E^S = -0.013 \pm 0.028$$

$$G_M^S = +0.62 \pm 0.31 \mu_N$$

Theories

1. Leinweber, et al.
PRL **94** (05) 212001
2. Lyubovitskij, et al.
PRC **66** (02) 055204
3. Lewis, et al.
PRD **67** (03) 013003
4. Silva, et al.
PRD **65** (01) 014016

1. Latest Lattice QCD

UIUC G0 web page:

“Excluded at 95.5% confidence level”



Thomas Jefferson National Accelerator Facility




Significance & Comparison with Lattice QCD

Size and sign of the strange magnetic moment
is astounding!

- For the deuteron, this result (G0) gives - $0.54 \mu_N$
- i.e. - 60% of its experimental magnetic moment!!
- Also remarkable versus lattice QCD which gives
 $+0.03 \pm 0.01 \mu_N$ (Leinweber et al., PRL 94 (2005) 212001)
- Sign would require violation of universality of
valence quark moments by $\sim 70\%$!



Modern Lattice QCD Result for G_M^s



Yields : $G_M^s = -0.046 \pm 0.019 \mu_N$

Leinweber et al., (PRL June '05) hep-lat/0406002



Thomas Jefferson National Accelerator Facility



U.S. DEPARTMENT OF ENERGY
S&T, September 11, 2005

January 2006: G_E^s by same technique

In this case only know Σ^- radius (and p and n)

$$2p + n = u^p + 3 O_N \qquad p + 2n = d^p + 3 O_N$$

$$\Rightarrow \langle r^2 \rangle_s = 0.000 \pm 0.006 \pm 0.007 \text{ fm}^2 ; 0.002 \pm 0.004 \pm 0.004 \text{ fm}^2$$

(c.f. using Σ^- : $-0.007 \pm 0.004 \pm 0.007 \pm 0.021 \text{ fm}^2$)

$$G_E^s(0.1 \text{ GeV}^2) = +0.001 \pm 0.004 \pm 0.004$$

(up to order Q^4)

Note consistency and level of precision!

Leinweber, Young et al., hep-lat/0601025: Jan 2006




Thomas Jefferson National Accelerator Facility




U.S. DEPARTMENT OF ENERGY
S&T, September 11, 2005

Ross Young: Why not use ALL the data?

No theoretical constraint (other than charge symmetry); use systematic Taylor expansion of $G_{E,M}^s$ in powers Q^2






HAPPEX-III and G0 Backward Angle

E05-109, HAPPEX-III, together with Backward angle G0, will provide an unprecedented precision on a measurement of **all three** strange form factors at $Q^2 = 0.64 \text{ GeV}^2$


HP 2010



SAMPLE and HAPPEX II Q^2
(final HAPPEX II error will be 1/3 of prelim)

G0 Backward Angle run Q^2 Values

HAPPEX III Q^2



Nuclear Physics: The Core of Matter, The Fuel of Stars

(NAS/NRC Report, 1999)

Science Chapter Headings:

The Structure of the Nuclear Building Blocks

The Structure of Nuclei

Matter at Extreme Densities

The Nuclear Physics of the Universe

Symmetry Tests in Nuclear Physics



Thomas Jefferson National Accelerator Facility




U.S. DEPARTMENT OF ENERGY
S&T, September 11, 2005

PREX : ^{208}Pb Radius Experiment

Low Q^2 elastic e-nucleus scattering

($E = 850 \text{ MeV}$, $\Theta=6^\circ$)

Z^0 (Weak Interaction) : **couples mainly to neutrons**



$$\frac{dA}{A} = 3\% \rightarrow \frac{dR_n}{R_n} = 1\%$$

Measure a Parity Violating Asymmetry

$$A = \frac{G_F Q^2}{2\pi\alpha\sqrt{2}} \left[1 - 4 \sin^2 \theta_w - F_n(Q^2) \right]$$

Applications:


- Fundamental check of Nuclear Theory
- Input to Atomic PV Expts
- Neutron Star Structure



Nuclear Structure

After more than 70 years, the neutron density of a heavy nucleus is a fundamental nuclear-structure observable that remains elusive!

- As fundamental as the charge density of a heavy nucleus
 - ★ *cf.* proton and neutron electromagnetic structure
- Reflects a poor understanding of the symmetry energy of NM
 - ★ Symmetry energy penalty imposed for breaking $N = Z$ balance
- Pure neutron matter well constrained at $\rho \approx (2/3)\rho_0$
- Slope is completely unconstrained by available nuclear data!



Adding the neutron radius of a single heavy nucleus to the database will eliminate the large dispersion in the plot!


Major Challenges for Nuclear Physics

- Origin of Nuclear Saturation

- EOS ... as $\rho \uparrow$; as $T \uparrow$
as $S \uparrow$; as $N-Z \uparrow$

- Phase Transition to:

- quark matter (QM)
- superconducting QM, strange condensate
- related to nuclear astrophysics; n-stars....







Neutron Star Composition

Hyperons enter at
just 2-3 ρ_0



Hence need effective
 Σ -N and Λ -N forces
in this density region!

Ξ - Hypernuclear data is
important input: we have none!





E01-011 (HKS) in Hall C: the Next Generation

Improved Trues to Accidentals Ratio



Reduced Backgrounds



Enhanced Luminosity

210 Lambdas



1390 Lambdas

HKS has demonstrated improved performance in its first run (completed 9/05)

Net factor of 50 in Figure of Merit, and resolution < 500 keV

New electron spectrometer under construction in Japan for final configuration




Thomas Jefferson National Accelerator Facility



U.S. DEPARTMENT OF ENERGY
S&T, September 11, 2005


Generalized Parton Distribution (GPDs) & Nucleon Tomography (06-003)

Begin the exploration of a major **New Direction** in **Hadron Physics** at 6 GeV, and **prepare for the full exploration at 12 GeV.**




Deeply Virtual Compton Scattering

3D mapping of the quark structure of the nucleon via Deeply Virtual Compton Scattering (DVCS)




Quark distributions in protons at different momentum fractions.



The shape of the proton varies with momentum.

Study of the nucleon orbital angular momentum.

HP 2008 +
Extract accurate information on generalized parton distributions in measurements of deeply virtual Compton scattering.



$$J_q = \frac{1}{2} A_q(0, \mu^2) + \frac{1}{2} B_q(0, \mu^2) = \frac{1}{2} \int dx x [H(x, \xi, 0, \mu^2) + E(x, \xi, 0, \mu^2)]$$

Thomas Jefferson National Accelerator


At 12 GeV: Exclusive ρ^0 with transverse target

$$A_{UT} = - \frac{2\Delta (\text{Im}(AB^*))/\pi}{|A|^2(1-\xi^2) - |B|^2(\xi^2+t/4m^2) - \text{Re}(AB^*)2\xi^2}$$

ρ^0

$$A \sim (2H^u + H^d)$$

$$B \sim (2E^u + E^d)$$



Asymmetry depends linearly on the GPD E , which enters Ji's sum rule.


K. Goeke, M.V. Polyakov,
M. Vanderhaeghen, 2001

Flavor Decomposition: semi-inclusive DIS

“semi-SANE” : **HP 2011**

DIS probes only the sum of quarks and anti-quarks → requires assumptions on the role of sea quarks


$$\sum e_q^2 (q + \bar{q})$$



Solution: Detect a final state hadron in addition to scattered electron

→ Can ‘tag’ the flavor of the struck quark by measuring the hadrons produced: ‘**flavor tagging**’

$$\sum e_q^2 q(x) D_{q \rightarrow M}(z)$$



(e,e') $W^2 = M^2 + Q^2 (1/x - 1)$

For M_m small, \vec{p}_m collinear with $\vec{\gamma}$, and $Q^2/v^2 \ll 1$


(e,e'm) $W'^2 = M^2 + Q^2 (1/x - 1)(1 - z)$

$z = E_m/v$

Flavor asymmetry of proton sea: Vacuum structure inside proton?


HP 2013

...New: Polarization!




pion cloud

$$\Delta\bar{u} - \Delta\bar{d} = 0$$



qq̄ pair
(Pauli blocking)

$$\Delta\bar{u} - \Delta\bar{d} > 0$$



...Can be answered by
initial SIDIS measurements
in 2009 and fully with 12 GeV

[Thomas 83; Schreiber *et al.*, 90;
Diakonov *et al.* 96; Fries, Schaefer, Weiss 03]



FROST: Frozen-Spin Target Experiments at CLAS (E02-112: 2007)

HP 2009

Determine basic symmetry properties of baryon matter at the core of the visible universe.

➤ Discover excited baryon states if produced in photoproduction of pseudo-scalar mesons.

➤ Use high-energy photons with circular and linear polarization on **FROST** with longitudinal and transverse spin polarization.

➤ **Measure nearly complete set of single, double, and triple polarization observables including hyperon recoil polarization.**


One of the top 10 milestones in hadron physics:

❖ **Complete the combined analysis of available data on single π , η and K photoproduction of resonances and incorporate the analysis of 2π final states into the coupled channel analysis of resonances.**



FROST

Example (sample PWA using MC data)
▶ greatly reduced uncertainties!




Thomas Jefferson National Accel

QCD: Unsolved in Nonperturbative Regime



The Nobel Prize in Physics 2004

Gross, Politzer, Wilczek



- 2004 Nobel Prize awarded for “asymptotic freedom”
- BUT in nonperturbative regime QCD is still unsolved
- One of the top 10 challenges for physics!
- Is it right/complete?
- Do glueballs, exotics and other apparent predictions of QCD in this regime agree with experiment?

JLab at 12 GeV is uniquely positioned to answer!




Thomas Jefferson National Accelerator Facility



U.S. DEPARTMENT OF ENERGY
S&T, September 11, 2005

Glueballs and hybrid mesons




Initial search
FY07 –
G12 (CLAS)

Colin Morningstar:
Gluonic Excitations workshop, 2003
(Jlab)

The EMC Effect: Nuclear PDFs

- Observation **stunned and electrified** the HEP and Nuclear communities 20 years ago
- Nearly 1,000 papers have been generated.....

- What is it




J. Ashman *et al.*, *Z. Phys. C57*, 211 (1993)

J. Gomez *et al.*, *Phys. Rev. D49*, 4348 (1994)

$g_1(A)$ – “Polarized EMC Effect”

- New calculations indicate larger effect for polarized structure than unpolarized: scalar field modifies lower cpts of Dirac wave function
(Cloet, Bentz, AWT, Phys Rev Lett 95 (2005) 0502302)

- Spin




The Q_{weak}^p Experiment


The first measurement of the weak charge of the proton; a precision test of the Standard Model and a search for New Physics Beyond the Standard Model – at the TeV scale

Weak Mixing Angle

Scale dependence in $\overline{\text{MS}}$ scheme including higher orders



- Electroweak radiative corrections
 $\rightarrow \sin^2\theta_W$ varies with Q



- Extracted values of $\sin^2\theta_W$ **must** agree with Standard Model or **new** physics is indicated.


$$Q_{\text{weak}}^p = 1 - 4 \sin^2 \theta_W \sim 0.072$$

- A 4% Q_{Weak}^p measurement probes for new physics at energy scales to:

$$\frac{\Lambda}{g} \sim \frac{1}{\sqrt{\sqrt{2}G_F} |\Delta Q_W^p|} \approx 4.6 \text{ TeV}$$

- Q_{weak}^p (semi-leptonic) and E158 (pure leptonic) make a powerful program to search for and identify new physics.


Q_{weak} Apparatus



Time Frame for 12 GeV & Advances in Lattice QCD ⇒ Wonderful synergy!

That is: Our growing ability to use lattice QCD to calculate the unambiguous consequences of nonperturbative QCD is beautifully matched to the capacity of Jlab at 12 GeV to measure the corresponding observables with precision!

....and hence really test if QCD is the complete theory of the strong interaction



Advances in Lattice QCD


Inclusion of Pion Cloud

Actions with exact chiral symmetry

Precise computations at
Physical Pion Mass

Advances in high-performance computing

Octet Magnetic Moments




Leinweber et al., PRL 94 (2005) 212001

Moments of Flavor-NS PDFs and GPDs - I

- Lattice QCD can compute both moments of GPD's with respect to x , and t -dependence


$$A_{n0}^q(-\bar{\Delta}_\perp^2) = \int d^2b_\perp e^{i\bar{\Delta}_\perp \cdot \bar{b}_\perp} \int_{-1}^1 dx x^{n-1} q(x, \bar{b}_\perp)$$



Decrease slope : decreasing transverse size as $x \rightarrow 1$ -

Burkardt

Lattice data: $m_\pi = 740$ MeV



From: LHPC & SESAM


Thomas Jefferson National Accelerator Facility



Lattice QCD Simulation of Change of Vacuum Structure in a Nucleon

Leinweber, Signal et al.

$\langle r \rangle = 0.16$ fm



Axion Search : Recent Observation by PVALS

Polarization experiments


- Send linearly polarized laser beam through transverse magnetic field \Rightarrow measure changes in polarization state
- Real and virtual production induce
 - **rotation:** photons polarized $\parallel \mathbf{B}$ will disappear leading to apparent rotation of polarization plane by

$$\varepsilon_\phi = -N_r \left(\frac{gB\ell}{4} \right)^2 F(q\ell) \sin 2\theta$$

- **ellipticity:** virtual production causes retardation between \mathbf{E}_\parallel and $\mathbf{E}_\perp \Rightarrow$ elliptic polarization

$$\psi_\phi \approx \frac{N_r}{6} \left(\frac{gB\ell}{4} \right)^2 \frac{m_\phi^2 \ell}{\omega} \sin 2\theta$$

for small masses, $m_\phi^2 \ell / 4\omega \ll 1$.




Publication:

[Zavattini et al. '05]

$$1.7 \times 10^{-6} \text{ GeV}^{-1} \lesssim g \lesssim 1.0 \times 10^{-5} \text{ GeV}^{-1}$$

$$0.7 \text{ meV} \lesssim m_\phi \lesssim 2.0 \text{ meV}$$

JLab FEL Power from THz to UV



For information: www.jlab.org/FEL




Thomas Jefferson National Accelerator Facility




U.S. DEPARTMENT OF ENERGY
S&T, September 11, 2005

Long-term Landscape : ELIC/eRHIC



Luminosity vs CM Energy



- ELIC at Jlab
 - 3-7 GeV e^- on
30-150 GeV p
(both polarized)
 - 20-65 GeV CM Energy
 - Polarized light ions
 - Luminosity as high as $0.8 \times 10^{35} \text{ cm}^{-2} \text{ sec}^{-1}$
- eRHIC at BNL
 - 5-10 GeV e^- on
50-150 GeV p
(both polarized)
 - 30-100 GeV CM Energy
 - Polarized light ions
 - Heavy ion beams available
 - Luminosity from 10^{33} to
perhaps as high as 10^{34}

AIM: Establish a New Paradigm for Nuclear Physics

In the 21st Century we have the challenge to unify our understanding of nuclear systems over otherwise impossible ranges of density and strangeness in terms of THE best candidate for a fundamental theory of the strong force: QCD

- **Precision electron scattering is essential to guide this unification**
- **On world scene JLab will beautifully complement the work in this area by J-PARC and GSI as well as RIA**
- **12 GeV will play a crucial role in solving one of the 10 outstanding problems in modern physics: origin of confinement**



Thomas Jefferson National Accelerator Facility



U.S. DEPARTMENT OF ENERGY
S&T, September 11, 2005



Thomas Jefferson National Accelerator Facility



U.S. DEPARTMENT OF ENERGY
S&T, September 11, 2005



Thomas Jefferson National Accelerator Facility



U.S. DEPARTMENT OF ENERGY
S&T, September 11, 2005
47