


Event reconstruction in LAr TPC



From “easy” to progressively more complicated topologies reconstruction



NuFact09 - IIT - Chicago

O. Palamara

Gran Sasso National Laboratory
(ITALY)

The Liquid Argon TPC Working Principle

- A charged particle crossing LAr produces e^- - Ar^+ pairs along its path.
- An Electric Field applied to the LAr volume makes ionization electrons to drift toward the TPC anode (made of 3 parallel wire planes: 1 grid and 2 read-out planes, wire pitch ~ 3 -4mm)
- Electrons drift over very long distances if Argon is very pure (1 meter drift requires purity level at 0.1 ppb)
- e^- -charges induce an electronic signal on the wires.
- Signals are acquired through low noise charge amplifiers and fast ADC waveform recording.
- Multiple non-destructing read-out wire signals can be assembled for 3D event reconstruction




Event reconstruction procedure in LAr TPC (I)

The purpose of the reconstruction procedure is to extract physical information provided by the wire output signals (multiple non-destructing read-out planes), i.e. the energy deposited by the different particles and the space coordinates where such a deposition has occurred (HIT)

→ to build a complete 3D (imaging) and calorimetric picture of the event

The offline reconstruction procedure consists of:

- hit identification*: the hits are independently searched for in every wire as signal regions of a certain width above the baseline;
- hit reconstruction*: the parameters defining the hit (position, height, area), which contain the physical information, are determined;



Event reconstruction procedure in LAr TPC (II)

3. *cluster reconstruction*: hits are **grouped** based on their position in the wire/drift coordinate plane (**2D reconstruction**);

4. *3D hit reconstruction*: the hit spatial coordinates are reconstructed by the association of hits from different views into **common track segments**;


5. *calorimetric reconstruction*: the determination of the **energy release** in LAr is performed in two steps:


- accounting for the charge loss due to the attachment by electro-negative impurities

$$Q_{\text{corr}} = Q e^{td/\tau_e}$$

- charge to energy conversion with correction for the quenching effect on the ionization charge in LAr (Birks law).


6. *Particle ID*: with dE/dx measurement vs. range





Observation of long ionizing muon tracks with the ICARUS T600 LAr TPC (test on surface)

About 18 m long c.r. muon tracks (~2000 collection wires)




Raw images from the Collection plane


ICARUS Coll. NIM A 508 (2003) 287


7/22/09

O. Palamara



Reconstructed energy deposition







Measurement of the μ decay spectrum with the ICARUS T600 LAr TPC (test on surface)


Fully reconstructed stopping muon event

Right chamber:
muon decay event views (Collection and Induction II)




7/22/09

O. Palamara



Michel Electron Spectrum

From the calorimetric reconstruction:
Energy spectrum of the electrons from
muon decay



ICARUS Coll. Eur. Phys. J. C 33 (2004), 233


- Study of stopping muon sample
 - 3000 events analyzed and fully reconstructed in 3D
- ρ parameter measurement (from comparison with MC simulation)

$$\rho = 0.72 \pm 0.06 (stat) \pm 0.08 (sys)$$


➤ Standard Model $\rho = 0.75$

- Energy resolution for electrons below ~ 50 MeV



$$\frac{\sigma(E)}{E} = \frac{11\%}{\sqrt{E}} \oplus 2\%$$



Energy reconstruction of e.m. showers from π^0 decays with the ICARUS T600 LAr TPC (test on surface)



π^0 signature is given by the presence of two e.m. showers coming from the two photons of the $\pi^0 \rightarrow \gamma\gamma$ decay, pointing to the hadronic interaction vertex



Measurement of the shower energy and shower direction


Selected sample
(after a fiducial volume cut):
196 π^0 candidates

Average mass:

$$m_{\gamma\gamma} = 134.4 \pm 3.0 \text{ MeV}/c^2$$


+ a contribution of 7.1% from systematics

Reconstructed invariant mass
of the photon pairs (a)



The ICARUS 50 lt Detector

(1st exposure of a LAr TPC to a neutrino beam $\langle E_\nu \rangle = 28$ GeV, 1998)




50 lt Detector backed by NOMAD exp. (magnetic spectrometer)


Trigger: $(SPS\ beam\ spill \cdot AND \cdot ScintCount \cdot AND \cdot (T1 + T2)) \cdot NOT \cdot (Chorus \cdot AND \cdot Veto)$

7/22/09

O. Palamara






The ICARUS 50 lt Detector




2D views and

3D reconstruction
of DIS event






2D views and




3D reconstruction
of QE event

- Collection of around 10 000 CC events
- Selection of 86 “golden sample” events with:
 - an identified proton of kinetic energy >40 MeV fully contained in the TPC and one muon whose direction extrapolated from NOMAD matches the outgoing track in the TPC.



The ICARUS 50 lt Detector





Ptcl. ID

Proton track
recognition


Proton reconstruction
and momentum
measurement
performed using only
TPC information

Energy Reconstruction for Muon and proton

Kinematic reconstruction of the outgoing muon performed using the tracking capability of NOMAD and traced back to the TPC.




Proton kinetic energy calculated from range



The ArgoNeuT Experiment (see M. Antonello talk on Friday)

- ArgoNeuT is a 175 l (active) **Liquid Argon TPC** exposed to the NuMI low energy neutrino beam at FNAL (Commissioning run: **May-June 2009**)
- ArgoNeuT detector is located between Minerva and the MINOS near detector at NuMI Tunnel – 100m underground. Muons escaping the TPC are reconstructed in MINOS ND.




- Collecting events in the 0.1 to 10 GeV range, ArgoNeuT is producing the first ever data for low energy neutrino interactions within a LArTPC.



External Trigger

for ν -beam operation or horizontal c.r. muons




Three main ν -beam topologies:


1. **through-going μ** from ν -int. in the rock upstream
 2. ν -int in LAr (good event to be selected)
 3. Empty event: No interaction in coincidence with beam spill
- or
1. **Horizontal c.r. muons**




3D reconstruction of a sample of cosmic muons




TOP VIEW



46 cm




2D and 3D reconstruction







Calorimetric reconstruction of cosmic muons: *detector performance validation test*



*Reasonable median cosmic ray
muon energy in the NuMI
underground location*





Reconstruction of neutrino CC interactions

ArgoNeuT expected rate at NuMi :



~ 19 QE events/day + ~ 15 RES events/day + ~ 83 DIS events/day

*Main aim of the experiment: QE cross section measurement with Ar target
in the few GeV energy range.*

QE process:



when nucleon bound in the nuclear target **nuclear effects** must be taken into account:




These products are usually neglected because not detectable,
unless...

.... a high quality imaging detector is in use !!

Reconstruction procedure (proton and muon) as for the ICARUS 50 lt




Sensitivity to nuclear effects: as an example, a particularly interesting event






2D and 3D reconstruction of the event






Final State Muon and Proton reconstruction




Proton reconstruction and momentum meas. performed using only TPC information. Kinetic energy calculated from range.


Proton kinetic energy = 80 MeV



Particle ID: Proton track recognition



A closer inspection of the event topology (thanks to the imaging capabilities of the LAr TPC) shows:




1) large activity near the vertex

$\mu^- + p + X$ (X = additional “short track” [2 wires] associated with high energy density deposition)

X compatible with a second 25 MeV p track from nuclear evaporation (FSI in nucleus) or pion re-absorption

2) an extra energy deposition (37 MeV) possibly associated with the event (e^+e^- pair), induced by a neutral particle.



This event reconstruction is still preliminary.

A full and detailed MC simulation including nuclear effects is required for validation.

A preliminary FLUKA MC simulation support the possibility to detect such nuclear effects in LAr TPC.

Conclusions

- Next-generation neutrino physics experiments require precision particle identification and **fine grained 3D imaging**. **Liquid Argon (LAr)** is recognised as an ideal detection medium, allowing the possibility of simultaneous ionisation charge, scintillation and Cerenkov light signals collection in large volumes.
- The LAr TPC is a detector particularly suitable to study low energy neutrino interactions due to its high energy resolution and its robust particle identification capability down to the **"few GeV range"**.
- **A big effort is under way to improve the event reconstruction procedures exploiting the full imaging and calorimetric capabilities of the LAr TPC technique.**